nature geoscience

LETTERS

PUBLISHED ONLINE: 25 NOVEMBER 2012 | DOI: 10.1038/NGEO1635

Extensive dissolution of live pteropods in the Southern Ocean

N. Bednaršek[1,2,3], G. A. Tarling[1]*, D. C. E. Bakker[2], S. Fielding[1], E. M. Jones[4], H. J. Venables[1], P. Ward[1], A. Kuzirian[5], B. Lézé[2], R. A. Feely[6] and E. J. Murphy[1]

Copyright / Ex. 4

[1]British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge, CB3 0ET, UK, [2]School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK, [3]University of Nova Gorica, Vipavska 13, 5000 Nova Gorica, Slovenia, [4]Royal Netherlands Institute for Sea Research, PO Box 59, 1790 AB Den Burg, Texel, The Netherlands, [5]Department of Geology and Geophysics, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA, [6]NOAA, Pacific Marine Environmental Laboratory, Seattle, Washington 98115, USA. *e-mail: gant@bas.ac.uk.

© 2012 Macmillan Publishers Limited. All rights reserved.

ED_002660K_00032655-00001

Copyright / Ex. 4

© 2012 Macmillan Publishers Limited. All rights reserved.

ED_002660K_00032655-00002

Copyright / Ex. 4

© 2012 Macmillan Publishers Limited. All rights reserved.

ED_002660K_00032655-00003

Copyright / Ex. 4

© 2012 Macmillan Publishers Limited. All rights reserved.

ED_002660K_00032655-00004

Copyright / Ex. 4

© 2012 Macmillan Publishers Limited. All rights reserved.

ED_002660K_00032655-00005