Ecological Indicators 76 (2017) 240–244

Contents lists available at ScienceDirect

Ecological Indicators

journal homepage: www.elsevier.com/locate/ecolind

Original Articles

# New ocean, new needs: Application of pteropod shell dissolution as a biological indicator for marine resource management

N. Bednaršek[a,*], T. Klinger[a], C.J. Harvey[b], S. Weisberg[c], R.M. McCabe[d], R.A. Feely[e], J. Newton[f], N. Tolimieri[b]

[a] School of Marine and Environmental Affairs, University of Washington, 3707 Brooklyn Avenue NE, Seattle, WA 98105 USA
[b] Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Boulevard E, Seattle, WA 98112 USA
[c] Southern California Coastal Water Research Project Authority, 3535 Harbor Blvd, Costa Mesa, CA 92626 USA
[d] University of Washington, Joint Institute for the Study of the Atmosphere and Ocean (JISAO), 3737 Brooklyn Avenue NE, Box 355672, Seattle, WA 98195 USA
[e] Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, 7600 Sand Point Way NE, Seattle, WA 98115 USA
[f] University of Washington, Applied Physics Lab, 1013 NE 40th Street, Box 355640, Seattle, WA 98105 USA

ARTICLE INFO

*Article history:*
Received 12 August 2016
Received in revised form 22 January 2017
Accepted 22 January 2017

*Keywords:*
Ocean acidification
Water quality assessment
Pteropod
Indicator
Sentinel species
Ocean acidification management

Copyright / Ex. 4

## 1. Introduction

Copyright / Ex. 4

Copyright / Ex. 4

* Corresponding author.
*E-mail address:* nina.bednarsek@gmail.com (N. Bednaršek).

http://dx.doi.org/10.1016/j.ecolind.2017.01.025
1470-160X/© 2017 Elsevier Ltd. All rights reserved.

Copyright / Ex. 4

Copyright / Ex. 4

Copyright / Ex. 4

Copyright / Ex. 4