Seagrass habitat metabolism increases short-term extremes and long-term offset of CO_2 under future ocean acidification

Stephen R. Pacella[a,b,1], Cheryl A. Brown[a], George G. Waldbusser[b], Rochelle G. Labiosa[c], and Burke Hales[b]

[a]Western Ecology Division, National Health and Environmental Effects Research Laboratory, Office of Research and Development, United States Environmental Protection Agency, Newport, OR 97365; [b]College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR 97331; and [c]Region 10, United States Environmental Protection Agency (EPA), Seattle, WA 98101

Edited by David M. Karl, University of Hawaii, Honolulu, HI, and approved February 16, 2018 (received for review March 7, 2017)

Copyright / Ex. 4

Copyright / Ex. 4

Author contributions: S.R.P., C.A.B., G.G.W., R.G.L., and B.H. designed research; S.R.P. and C.A.B. performed research; B.H. contributed new reagents/analytic tools; S.R.P., G.G.W., and B.H. analyzed data; and S.R.P. and G.G.W. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

Data deposition: This dataset has been published via the EPA Office of Research and Development Environmental Dataset Gateway (https://edg.epa.gov; doi: 10.23719/1407616).

[1]To whom correspondence should be addressed. Email: spacella@coas.oregonstate.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1703445115/-/DCSupplemental.

ED_002660_00002484-00001

Copyright / Ex. 4

Copyright / Ex. 4

Copyright / Ex. 4

Copyright / Ex. 4

Copyright / Ex. 4