Message

From: White, Roshanna [White.Roshanna@epa.gov]
Sent: 8/5/2019 1:35:22 PM
To: Wahlstrom-Ramler, Meghan [Wahlstrom-Ramler.Meghan@epa.gov]; Ferry, Rol [Ferry.Roland@epa.gov]
CC: Holliman, Daniel [Holliman.Daniel@epa.gov]
Subject: RE: Velella Epsilon EA Changes

No, I assumed after the round of discussions you all had last week that Rol would answer me this week with his answer.

Roshanna

From: Wahlstrom-Ramler, Meghan <Wahlstrom-Ramler.Meghan@epa.gov>
Sent: Monday, August 5, 2019 9:33 AM
To: White, Roshanna <White.Roshanna@epa.gov>; Ferry, Rol <Ferry.Roland@epa.gov>
Cc: Holliman, Daniel <Holliman.Daniel@epa.gov>
Subject: RE: Velella Epsilon EA Changes

Did you ever get an answer for this?

Thanks,

Meghan

Meghan Wahlstrom | Environmental Engineer | Permitting & Grants Branch | Water Division
Special Emphasis Program Manager for the Federal Women's Employment Program (FWP)
Region 4 | Atlanta Federal Center | 61 Forsyth Street SW | Atlanta, GA 30303-8960
(404) 562-9672 | Wahlstrom-ramler.meghan@epa.gov

From: White, Roshanna <White.Roshanna@epa.gov>
Sent: Tuesday, July 30, 2019 10:19 AM
To: Ferry, Rol <Ferry.Roland@epa.gov>
Cc: Wahlstrom-Ramler, Meghan <Wahlstrom-Ramler.Meghan@epa.gov>; Holliman, Daniel <Holliman.Daniel@epa.gov>
Subject: Velella Epsilon EA Changes

Rol:
The buffer for the cage will be changed to 500 m in the permit and other documents. How would you like to proceed with the change in the buffer in regard to these statements in the EA?

4.2.2 Sediment Quality

"Studies of offshore aquaculture operations in the Mediterranean showed that the severe effects of organic inputs from fish farming on benthic macrofauna are limited to up to 25 m from the edge of the cages (Lampadariou, Karakassis, & Pearson, 2005) although the influence of carbon and nitrogen from farm effluents in sea floor can be detected in a wide area about 1,000 m from the cages (Sara, Scilipoti, Mazzola, & Modica, 2004)."

5.4.2 Sediment Quality

"As discussed in *Section 4.2.2 Sediment Quality*, numerous studies within the Mediterranean have shown that organic inputs from fish farms on benthic macrofaunal are only limited up to 25 m from the edge of the cages (Lampadariou, Karakassis, & Pearson, 2005) and carbon and nitrogen produced by fish farm effluents on the sea floor is detected in an area about 1,000 m from the cages (Sara, Scilipoti, Mazzola, & Modica, 2004)."

"...Additionally, waste from ships could contribute to cumulative impacts associated with organic and inorganic pollution. It is unlikely that organic and nitrogen from land-based discharges would reach the proposed facility 45 miles offshore. Conversely, the effluent from the cages would not travel past 1,000 m to incrementally combine with these other organic and nitrogen laden discharges to cause a cumulative impact."

5.5.6 Essential Fish Habitat

"Additionally, waste from ships could contribute to cumulative impacts associated with organic and inorganic pollution. It is unlikely that organic and nitrogen from land-based discharges would reach the proposed facility 45 miles off shore. Conversely, the effluent from the cages would not travel past 1,000 m to incrementally combine with these other organic and nitrogen laden discharges to cause a cumulative impact."

Roshanna
Voice: 404-562-9035 | Email: white.roshanna@epa.gov