EPA

MRID NUMBER

40798001

A. Craven
ESB

Manual of Acute Toxicity: Interpretation and Data Base for 410 Chemicals and 66 Species of Freshwater Animals



BEST AVAILABLE COPY

UNITED STATES DEPARTMENT OF THE INTERIOR
Fish and Wildlife Service / Resource Publication 160

ED_005172C_00000035-00002

RESOURCE PUBLICATIONS

This publication of the Fish and Wildlife Service is one of a series of semitechnical or instructional materials dealing with investigations related to wildlife and fish. Each is published as a separate paper. The Service distributes a limited number of these reports for the use of Federal and State agencies and cooperators. A list of recent issues appears on inside back cover.

Copies of this publication may be obtained from the Publications Unit, U.S. Fish and Wildlife Service, Washington, DC 20240 or may be purchased from the Government Printing Office, Superintendent of Documents, Washington, DC 20402 and National Technical Information Service (NTIS), 5285 Port Royal Road, Springfield, VA 22161

Library of Congress Cataloging-in-Publication Data

Mayer, F. L. (Foster Lee), 1942-
Manual of acute toxicity.

(Resource publication / United States Department of the Interior, Fish and Wildlife Service ; no. 160)
Bibliography: p.
1. Aquatic animals—Effect of water pollution on. 2. Pesticides and wildlife. 3. Pesticides—Toxicology. I. Ellersieck, Mark R. II. Title. III. Series: Resource publication (U.S. Fish and Wildlife Service) ; no. 160.
S914.A3 no. 160 333.95'4'0973 s [591.2'4] 86-11312
[QH545.P4]

ED_005172C_00000035-00003

Manual of Acute Toxicity: Interpretation and Data Base for 410 Chemicals and 66 Species of Freshwater Animals

By Foster L. Mayer, Jr.
Mark R. Ellersieck

UNITED STATES DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE
Resource Publication 160
Washington, D.C. • 1986

3

ED_005172C_00000035-00004

Contents

	Page
Acknowledgments	iii
Abstract	1
Materials and Methods	3
General	3
Statistical Analyses	4
Results and Discussion	5
Comparative Toxicity	5
Factors Affecting Toxicity	18
Conclusions	28
References	30
Appendix	33

ED_005172C_00000035-00005

Acknowledgments

We thank the many scientists and technicians who assisted in the work, as well as the numerous companies that provided the chemicals for testing. Keypunching was done by J. English; L. C. Sappington managed the formating, verifying, and analyzing of data with assistance by J. D. Lutz and J. A. Thomas; F. J. Borgers, Jr., helped manage the data base; the Environmental Effects Staff (Office of Toxic Substances, U.S. Environmental Protection Agency, Washington, D.C.) and K. S. Mayer provided constructive reviews; and R. C. Bernhardt and E. F. Head typed the manuscript. This manuscript was reviewed for scientific content by T. W. Duke, E. E. Kenaga, A. J. McErlean, D. I. Mount, J. M. Neuhold, and R. L. Spehar. Funding for this project was provided by the U.S. Environmental Protection Agency under Task 1B of Interagency Agreement EPA-IAG-AD-14-F-1-556-0.

ED_005172C_00000035-00006

Manual of Acute Toxicity: Interpretation and Data Base for 410 Chemicals and 66 Species of Freshwater Animals

By

Foster L. Mayer, Jr.[1]

Columbia National Fisheries Research Laboratory
U.S. Department of the Interior
Fish and Wildlife Service
Route 1
Columbia, Missouri 65201

and

Mark R. Ellersieck

Department of Food Science and Nutrition
University of Missouri
Columbia, Missouri 65211

Abstract

All acute toxicity data developed by the Columbia National Fisheries Research Laboratory, U.S. Fish and Wildlife Service, since 1965 were evaluated for quality, and a data base was established for 4,901 tests with 410 chemicals (mainly pesticides) and 66 species of aquatic animals. The data were also analyzed by various statistical approaches to make taxonomic comparisons, and to assess the degree to which various factors affect toxicity. Insects were the most sensitive group, followed by crustaceans, fishes, and amphibians. Among the four most commonly tested forms, daphnids were the most sensitive 58% of the time, followed by rainbow trout, *Salmo gairdneri* (35%), bluegills, *Lepomis macrochirus* (5%), and fathead minnows, *Pimephales promelas* (2%). In comparisons with other species tested with the same chemical, two observations were evident: (1) the lowest of the two LC50's for daphnids and rainbow trout was <15× that of the most sensitive species 95% of the time, and <25 × 100% of the time; and (2) testing of three species (*Daphnia*, *Gammarus*, and rainbow trout) provided the lowest toxicity value 88% of the time, and could not be improved more than 2.5% by adding any other single species. Interspecies correlation models can be used to predict acute toxicity values, but the confidence limits were smallest when extrapolations were within families. Flow-through toxicity values can be obtained by multiplying static values by 0.51, but the error is likely to be large because ratios of static to flow-through tests differed by 1.8 orders of magnitude. Toxicity of aged test solutions increased by a factor of 2 or more only 11% of the time and never increased by more than a factor of 4; toxicity decreased in 22% of the tests and remained unchanged in 69%. Although pH affected the toxicity of only 20% of the chemicals tested, it caused a greater average

[1]Present address: Environmental Research Laboratory, U.S. Environmental Protection Agency, Sabine Island, Gulf Breeze, Fla. 32561.

ED_005172C_00000035-00007

change in toxicity than did any other factor examined. Temperature affected the toxicity of 40% of the chemicals tested, toxicity generally increasing with increases in temperature—although the toxicity of a few chemicals (DDT, dimethrin, and methoxychlor) decreased. The effect of temperature on toxicity conformed to the Q_{10} concept and was predictable. Temperature increased the toxicity of most chemicals by a factor of 3.1 per 10°C rise in temperature, but the factor for organophosphate insecticides was higher (5.1)—possibly due to the simultaneous increase in acetylcholinesterase (AChE) activity and rate of AChE inhibition. Regression slopes were consistent among species within a chemical for both pH and temperature, indicating chemical rather than biological differences in toxicity. Water hardness had little if any effect on the toxicity of organic chemicals; any effects observed were probably due to pH differences in solutions of differing hardness. Formulations of pesticides increased the toxicity of technical material 32% of the time, and decreased it 11% of the time. The effect of formulation on toxicity was within a factor of 0.2 to 5× 85% of the time, but was as high as 318×. The diet fed to fish before they were tested altered toxicity by as much as 5.7×, but factors of 1.1 to 3.8× would be anticipated within available commercial diets. Source of rainbow trout contributed to variation in toxicity data, differences ranging from 1.8 to 4.7× 88% of the time. Sensitivity of invertebrates and fish decreased as development of the animals progressed and size increased. Differences in toxicity among different life stages of fish were 5× or less 83% of the time. The change in toxicity was usually small within the sizes normally tested.

Of the factors affecting toxicity, the differences were within 5× or less 80% or more of the time, and this generalization included pH and temperature, evaluated on the basis of a 1.0 unit or 10°C change. Generalizations and predictions can be made for comparative toxicity and factors affecting toxicity ≥80% of the time. However, due to inherent variation and frequent exceptions in acute toxicity data, it remains best to test the chemical and species of concern under environmental conditions of interest when precise information is required.

The Federal Insecticide, Fungicide, and Rodenticide Act (PL 80-104), as amended by the Federal Environmental Pesticide Control Act of 1972 and others (7 U.S.C. 136-136y), requires that all formulators and producers of pesticides register their products and establishments with the U.S. Environmental Protection Agency (EPA) on the basis of safety data regarding health and environment. Furthermore, the Toxic Substances Control Act of 1976 (PL 94-469) and the Federal Insecticide, Fungicide, and Rodenticide Act authorize EPA to obtain data from industry on the health and environmental effects of chemical substances and mixtures. Acute toxicity testing plays a major role in the testing requirements involved in both laws, for good reason. In terms of utility for the assessment of hazard to aquatic environments, acute toxicity tests were rated highest of 15 tests evaluated by Brungs and Mount (1978) and Macek et al. (1978). Six criteria were used in the evaluations: ecological significance of effects, scientific and legal defensibility, availability of acceptable methodology, utility of test results in predicting effects in aquatic environments, general applicability to all classes of chemicals, and simplicity and cost of the test.

Toxicologists are well aware of the merits and limitations of the acute toxicity test; yet, there are probably few measurements for evaluating hazard or safety of a chemical to aquatic life that have been as grossly misunderstood. Users of acute toxicity data must bear in mind that the LC50 generally measures only one biological response—a lethal one. Its main value lies in its provision of a relative starting point for hazard assessment, along with the expected environmental concentration, and other measurements (e.g., water solubility, partition coefficient, and degradation rate) of the chemical in question. In addition, the acute toxicity test provides a rapid, cost-efficient way to measure relative toxicity of different forms and formulations of a chemical, toxicity in different types of water, and toxicity to organisms representing different trophic levels (Buikema et al. 1982; Sprague 1970).

One of the greatest needs in acute toxicity testing today is an evaluation of the data already available for its utility and applicability in the assessment of chemical hazard. At present, there is a tendency to dwell on the selection of species for toxicity tests and the biological, chemical, and physical factors affecting acute toxicity, without fully defining the

ED_005172C_00000035-00008

variation in species sensitivity, the extent to which various physicochemical and biological factors affect toxicity, and the probabilities associated with such variation. Assessments of the probability that adverse ecological effects will result from environmental releases of a given contaminant are of primary importance in hazard evaluation. Probabilities should be derived that will reasonably represent acute toxicity to most species—especially the more sensitive ones. However, the probabilities that can be developed today are, for the most part, based on species easily and economically tested, and therefore may or may not include a reasonable portion of the very sensitive species. Furthermore, a hazard assessment is not a one-time estimate. Additional evaluations must be made as the data base and knowledge expand; and at the present state of the art, no scheme or procedure can eliminate the need for sound scientific judgment.

The large acute toxicity data base (Appendix Table A5) used in the analyses presented here is unique in that the research was conducted within one laboratory system and by methods that, for the most part, are the consensus methods in use today (American Society for Testing and Materials 1980; Committee on Methods for Toxicity Tests with Aquatic Organisms 1975). The objectives were as follows:

1. Make available the data base, especially unpublished data.
2. Develop the general statistical relation of relative sensitivities within species, families, and classes.
3. Determine the frequency with which EC50's or LC50's determined with commonly tested species (daphnids; rainbow trout, *Salmo gairdneri*; fathead minnows, *Pimephales promelas*; or bluegills, *Lepomis macrochirus*) are lower than the EC50's or LC50's for all other species tested.
4. Determine if there is a group of three or four test species that would yield the lowest EC50 or LC50 at least 95% of the time.
5. Develop the statistical relation between static and flow-through LC50's.
6. Determine how often the toxicity of an aged test solution is less than that of a fresh solution.
7. Assess the effects of physical, chemical, and biological factors on acute toxicity.

We hope that these data will contribute to the many assumptions, comparisons, and extrapolations required in an ecotoxicology testing scheme for hazard assessment and for other purposes.

Materials and Methods

General

The data base used to prepare this report (Appendix Table A5) was compiled from the results of tests conducted at the Columbia National Fisheries Research Laboratory and its field laboratories in 1965–1984, and judged acceptable according to good laboratory practices. It consists of data from 4,901 tests performed with 66 species of aquatic organisms and 410 chemicals. Not all of the species were tested with every chemical. The species of invertebrates, fishes, and amphibians tested are listed in phylogenetic order (Appendix Tables A1 and A2), and chemicals are arranged alphabetically by common name or trade name where common names do not exist (Appendix Table A4). Toxicity values are expressed as ng/L, μg/L, mg/L, or ml/L. Duration of the toxicity tests was either 48 h (fairy shrimp, *Streptocephalus seali*; daphnids, *Daphnia magna*, *D. pulex*, and *Simocephalus serrulatus*; seed shrimp, *Cypridopsis vidua*; and larval midges, *Chironomus plumosus*) or 96 h (all other species). The chemicals represent all major pesticides, as well as several industrial and inorganic chemicals. Most of the data are on organic chemicals and fishes. Data for invertebrates were often lacking, and are therefore discussed only when adequate data exist.

Portions of the data base (static acute tests on 271 chemicals, 58 species, and 1,587 tests) were previously published by Johnson and Finley (1980). The present data base includes data from tests on 139 additional chemicals. In many of the tests, the following factors were assessed: test type (static or flow-through), aging of test solutions, temperature, pH, hardness, formulation, diet, fish source, and size and life stage. Where the few discrepancies occur between our data and those of Johnson and Finley (1980), the present data should be used. Because we used original data records to prepare this report, the absence of data that were previously published does not necessarily mean that the data were unacceptable: they may have been lost or were unavailable. Furthermore, although some data on certain chemicals did not appear to conform with other test results, the tests were included in the data base because they could not be eliminated on the basis of good laboratory practice.

Static toxicity test procedures with invertebrates and fishes were previously described by Johnson and Finley (1980). Static and flow-through test techniques

ED_005172C_00000035-00009

were generally those of the American Society for Testing and Materials (1980) and the Committee on Methods for Toxicity Tests with Aquatic Organisms (1975). Test procedures for amphibians (tadpoles) were those described by Sanders (1970). The level of statistical significance in all tests was $P < 0.05$.

Statistical Analyses

Comparative Toxicity

Taxonomic comparisons. The data set analyzed included most of the data in Appendix Table A5, but did not include data from tests on eggs, flow-through tests, or aged test solutions, or tests yielding "less-than" or "more-than" values. The data were analyzed by one-way analyses of variance (Snedecor and Cochran 1967) that included the effect of category (species, family, or class) for each chemical. Log_{10} transformation of toxicity data was used to meet the assumptions for normally distributed data in analysis of variance. The first step in statistical analyses of the data set was to determine the number of repeated measurements (48-h EC50's and 96-h LC50's) within a category for each chemical. This step was necessary to enable derivation of an error term for testing differences among species, families, or classes. The design was not balanced because the numbers of observations in each category were unequal. To determine mean differences among EC50's or LC50's, we used Fisher's least significance difference (LSD), and sum of ranks/*n* analyses, based on the log of the values or means that were used for comparisons of sensitivity. Frequency analysis by inspection—determination of how often the EC50 or LC50 of one taxonomic group was greater or less than that of others—was also used.

Interspecies comparisons and correlations. Interspecies comparisons were made by summing ranks/*n* and frequency analysis by inspection—determining how often the EC50 or LC50 for one species was greater or less than that for others. The values used for comparisons were log values; their derivation is described in the materials and methods for taxonomic comparisons.

We conducted interspecies correlations, using Model II least squares methodology (Snedecor and Cochran 1980), on daphnids (*Daphnia magna* or *D. pulex*, or both), rainbow trout, fathead minnows, and bluegills with other species. A standardized data set was established from Appendix Table A5 for these analyses, to reduce variation. The data set was based mainly on technical materials, and water temperature, pH, and hardness conformed to the requirements of American Society for Testing and Materials (1980) and the Committee on Methods for Toxicity Tests with Aquatic Organisms (1975); data were not included from tests on eggs, flow-through tests, or aged test solutions, or tests having "less-than" or "more-than" values. Acceptable fish weights were 0.2 to 1.5 g. The coefficient of variation for a species with three or more tests on a chemical (all fishes) averaged 37% (range, 3–115%) for toxicity values in the standardized data set, whereas the data used in the comparative toxicity section on species averaged 65% (range, 13–215%). Slopes and intercepts were derived from the equation $\log Y = a + b(\log X)$, where X equaled toxicity values for one of the four species previously mentioned and Y equaled toxicity values for other species. Paired tests on five or more chemicals were the minimum requirement for inclusion in the analyses. When either one of the paired values included more than one EC50 or LC50 value, the logs of the individual values were averaged.

Frequency distribution of toxicity. Estimates of frequency distributions of toxicity of all chemicals for the categories of species, class, and all taxa combined were performed with the standardized data set. Each chemical was classified by the following 48-h EC50 or 96-h LC50 concentrations (μg/L): <0.01, 0.01 < 0.1, 0.1 < 1.0, 1.0 < 10, 10 < 100, 100 < 1,000, 1,000 < 10,000, 10,000 < 100,000, or ≥100,000. Frequency distributions of the toxicity classification were then made for each species and class, and for all taxa tested. Only results for 14 of the individual species are presented graphically, but 75% of the tests were conducted with those species.

Static and flow-through tests and aged test solutions. Formulation and categorization of the ratios were used in analyzing data on static and flow-through tests and aged test solutions. Ratios of static to flow-through tests (96-h LC50's) were calculated for 123 paired tests that included 41 chemicals and 19 species. The geometric mean was used when two or more toxicity values existed under similar conditions. The percent incidence of the ratios occurring in three categories was determined: at <0.5, the chemical was more toxic under static than under flow-through conditions; at 0.5–1.5, there was no difference in toxicity between the two types of tests; and at >1.5, the chemical was more toxic under flow-through than under static conditions.

ED_005172C_00000035-00010

Aged test solutions were compared with fresh test solutions (46 paired tests), under static conditions, for 22 pesticides and 12 species. Solutions were aged for 4 to 28 days, and 96-h LC50's determined on both fresh and aged test solutions. Most comparisons were made at 7 days, for consistency. Deactivation indices (Marking and Dawson 1972) were obtained by dividing the 96-h LC50 of an aged test solution by the 96-h LC50 of the paired fresh solution. Further details about the deactivation index were described by Marking (1972). The percent incidence of the indices was determined for three categories: at <0.5, the chemical was more toxic after aging; at 0.5–1.5, aged and fresh solutions were equally toxic; and at >1.5, aged solutions were the less toxic.

No differences were considered to exist at ratios of 0.5 to 1.5 in comparisons between static and flow-through tests, and between aged and fresh test solutions. Although this ratio range is not statistically exact, it appears to encompass most of the range observed within normal experimental variation. This approach was also used in comparing the toxicity of technical materials with that of formulated pesticides.

Factors Affecting Toxicity

Temperature, pH, and hardness of test solution, and size of test animal. We conducted regression analyses, using Model II least squares methodology, to determine relations between 48-h EC50's or 96-h LC50's and pH, temperature (°C), and hardness (mg/L as $CaCO_3$) of the test water, or size (weight, g) of the test animals. Slopes and intercepts were derived from the equation $\log Y = a + bX$, where Y = the toxicity value and X = pH, temperature, hardness, or size. Only tests containing three or more data points (EC50's or LC50's) for a specific species and chemical combination were used. Analyses of covariance were computed to determine differences between linear slopes for treatment. Also, we tested the hypothesis that the individual linear slopes were equal to zero by using individual regression analyses, as well as by calculating a pooled estimate of the standard error in the analysis of covariance (Snedecor and Cochran 1980). If the slopes had been analyzed individually, they might have differed significantly from zero, but not necessarily with analysis of covariance. Only tests having slopes that differed significantly from zero are presented in tabular form.

Formulation, diet, source, and life stage. Ratios of technical to formulated pesticides (static 48-h EC50's or 96-h LC50's), based on active ingredients, were calculated for 161 paired tests that included 48 chemicals and 16 species. The percent incidence of the ratios occurring in three categories was determined: at <0.5, the formulated pesticide was less toxic than the technical form; at 0.5–1.5, technical and formulated pesticides were equally toxic; and at >1.5, the formulated pesticide was the more toxic.

Differences among 96-h LC50's attributable to diet were determined by comparing means within 95% confidence intervals (Mehrle et al. 1977). Frequency analysis by inspection—how often the 96-h LC50 of one category was greater or less than that of another category—was used for comparisons of fish source and life stage, owing to the small sample sizes and characteristics of the data.

Results and Discussion

Comparative Toxicity

Taxonomic Comparisons

Species. Acute static toxicity data were highly variable among the 63 species tested against 174 chemicals. Some species appeared to be very sensitive or very insensitive overall, but there were also frequent exceptions for the same species. The largest number of species (14 to 25) consistently tested in the total data base were tested with 14 insecticides (2 carbamates, 6 organochlorines, and 6 organophosphates). Data from these tests were compared by sum of ranks/n to provide a general sensitivity ranking. In descending order of sensitivity (not necessarily statistically significant), the species ranked as follows: stoneflies (*Claassenia sabulosa*, *Pteronarcys californica*, *Pteronarcella badia*), glass shrimp (*Palaemonetes kadiakensis*), amphipods (*Gammarus fasciatus*), daphnids (*Daphnia pulex*, *D. magna*, *Simocephalus serrulatus*), brown trout (*Salmo trutta*), rainbow trout (*S. gairdneri*), seed shrimp (*Cypridopsis vidua*), largemouth bass (*Micropterus salmoides*), cutthroat trout (*S. clarki*), bluegills (*Lepomis macrochirus*), sow bugs (*Asellus brevicaudus*), coho salmon (*Oncorhynchus kisutch*), yellow perch (*Perca flavescens*), channel catfish (*Ictalurus punctatus*), common carp (*Cyprinus carpio*), black bullheads (*I. melas*), green sunfish (*L. cyanellus*), fathead minnows (*Pimephales promelas*), goldfish (*Carassius auratus*), western chorus frog

BEST AVAILABLE COPY

ED_005172C_00000035-00011

(*Pseudacris triseriata*), and Fowler's toad (*Bufo woodhousei fowleri*). Analysis of the standardized data provided a slightly different ranking: stoneflies, glass shrimp, daphnids (*Daphnia magna*, *D. pulex*), amphipods, daphnids (*Simocephalus serrulatus*), rainbow trout, sow bugs, bluegills, largemouth bass, seed shrimp, coho salmon, cutthroat trout, yellow perch, brown trout, channel catfish, common carp, green sunfish, fathead minnows, goldfish, black bullheads, western chorus frog, and Fowler's toad. The rankings should be viewed only as general trends, since not all species were tested with all chemicals, and all the chemicals tested were insecticides. The overall trends for species sensitivity were, however, consistent with those found for family and class groupings.

Among 82 chemicals tested with 6 or more (up to 42) species, the highest toxicity values within a chemical averaged 256× the lowest, and ranged from 2.6 to 166,000×. Average differences were largest for insecticides (868×; range, 5.0 to 166,000×), followed by herbicides, fungicides, and other pesticides ($\bar{x}$, 33×; range, 2.6 to 438×); the few industrials were lowest ($\bar{x}$, 8.8×; range, 3.6 to 127×).

The four forms most commonly tested (*Daphnia magna* or *D. pulex*, rainbow trout, fathead minnows, and bluegills) were compared with each other in tests of 40 chemicals in which all four were tested together. *Daphnia* was the most sensitive 58% of the time, followed by rainbow trout (35%), bluegills (5%), and fathead minnows (2%). Sensitivity was greatly increased in combinations of *Daphnia* and rainbow trout only in that one of the two species was the most sensitive 93% of the time; in the remaining 7%, fathead minnows and bluegills were the most sensitive. However, the rainbow trout was always the next most sensitive species, and the LC50's were never significantly different from those of fathead minnows and bluegills.

Kenaga (1978) reported that no aquatic species was always the least or most sensitive for the 42 compounds he evaluated with rainbow trout, fathead minnows, and bluegills. Blanck et al. (1984), who tested 18 chemicals and an industrial wastewater with 13 green algae, showed that differences in sensitivity among species within a chemical may be as high as a factor of 2,000. Blanck (1982) concluded that no algal species of the 13 tested was generally sensitive or generally insensitive to all chemicals, and that even within a group of organisms such as algae, it is impossible to select a test species with a sensitivity representative of that group. Mount (1982) basically agreed with this assessment when, on the subject of surrogate species, he wrote:

> The point is that we have ample data to prove that a species can only represent itself consistently and not a group.
>
> However, species can be viewed as indicators of sensitivity in quite a different way. We can see that species sensitivity (LC50 or LD50) distributes itself in a rather consistent way for most chemicals. The distribution resembles a log normal one. We can then take the approach of sampling this distribution in order to predict the range about the mean. This is in reality our objective. Thus, each species we test is not representing any other species but is one estimate of general species sensitivity. With several such estimates, the overall range of sensitivity for all species can be determined. Our problem is to know how many species and what type of species to test to adequately represent the whole range.

Birge and Black (1982) also agreed with this concept, and further stated that with as few as three or no more than five aquatic species, it should be possible to establish a "biological response range." Blanck et al. (1984) reported that toxicity values for 3 algal species (randomly selected from the 13 studied) would be within two orders of magnitude of the most sensitive algae tested for 95% of the chemicals, and that a 5-species group would increase predictiveness to 99% of the chemicals. Although Mount (1982) maintained that species must be randomly chosen, he found that for most chemicals, the five most commonly tested species (*Daphnia magna*, *Gammarus*, rainbow trout, fathead minnows, and bluegills) represented a large part of the range of sensitivity of all species tested.

The many chemicals requiring hazard assessment and the lack of testing capabilities to quickly screen such an array of chemicals may require an effective toxicity screening process within a short time, as opposed to the "Surrogate Species Cluster Concept." This assessment is further complicated by the many new chemicals marketed each year (Maki et al. 1979). Kenaga (1978) proposed methods for early assessment of acute toxicity of chemicals to aquatic organisms, which were further expanded by Doherty (1983). Kimerle et al. (1983) analyzed an approach used by the Organization for Economic Cooperation and Development and the U.S. Environmental Protection Agency that includes only two or three species—a daphnid, a fish, and possibly an alga. This

ED_005172C_00000035-00012

latter approach is discussed first, based on our data.

Kimerle et al. (1983) compared the lowest LC50 of the four most commonly tested species to the LC50 of other species tested within 82 chemicals by orders of magnitude. Daphnids (*Daphia magna* or *D. pulex*, or both), rainbow trout, fathead minnows, or bluegills were within an order of magnitude of the most sensitive species 76, 74, 47, and 52% of the time, and within two orders of magnitude 93, 91, 74, and 70% of the time, respectively. Combinations of the three fish species did not improve the percent of chemicals included within one and two orders of magnitude. However, daphnids in combination with one of the three fish species (69 to 81 chemicals) greatly improved the frequency: 90% of the chemicals were within one order of magnitude of the lowest LC50 value and up to 98% were within two orders of magnitude.

The four main species were tested together with 40 chemicals (6 to 42 species per chemical) in our data, and the LC50 comparisons generally agreed with those of Kimerle et al. (1983). Daphnids (*Daphnia magna* or *D. pulex*), rainbow trout, fathead minnows, and bluegills were within an order of magnitude of the most sensitive species 75, 35, 28, and 38% of the time and within two orders of magnitude 90, 65, 58, and 72% of the time, respectively. The testing of daphnids in combination with one of the three fishes increased the frequency to 85% within one order of magnitude and 98 to 100% within two orders of magnitude. More specifically, the exact orders of magnitude to include the most sensitive species tested 100% of the time within the 40 chemicals we examined were 1.40 (25×) for daphnid-rainbow trout combinations, 2.05 (112×) for daphnid-fathead minnows, and 1.87 (74×) for daphnid-bluegills. In other words, one could divide the lowest of the two LC50's for daphnids and rainbow trout by 25 to obtain an estimate of an LC50 that would include the "most sensitive" species, based on the 6 to 42 species examined. The value for daphnid-fathead minnows was 112 and that for daphnid-bluegills was 74. The values that included the most sensitive species 95% of the time for the three combinations of species were 15 for daphnid-rainbow trout, 46 for daphnid-fathead minnows, and 28 for daphnid-bluegills.

As attractive as it may seem, the use of specific factors can lead to overestimation of the toxicity for the most sensitive species with some chemicals. The daphnid-rainbow trout combinations were examined thoroughly within the 40 chemicals, since in the few tests where LC50's for fathead minnows or bluegills were below those for daphnids or rainbow trout, the differences were not statistically significant. Other species were more sensitive than daphnids or rainbow trout to 27 of the 40 chemicals tested—only 0.5 order of magnitude difference in toxicity values included 59% of the 27 chemicals, and one order of magnitude included 85%. However, LC50's of the most sensitive species were significantly less than those of daphnids or rainbow trout in 11 of the 27 chemicals. No particular chemical type was represented by the 11 compounds (aldrin, benomyl, DDD, fenitrothion, fenvalerate, fluridone, heptachlor, lindane, permethrin, Pydraul 50E, and terbufos) except that eight were insecticides. The species having values (n) significantly less than those of *Daphnia* or rainbow trout were *Gammarus* (5), *Orconectes nais* (1), *Pteronarcys californica* (2), *Chironomus plumosus* (1), coho salmon (1), and channel catfish (1). If an amphipod (*Gammarus*) was included in the comparisons as a test species, as suggested by Mount (1982), the number of chemicals having a species LC50 significantly less than that for daphnids, amphipods, or rainbow trout was five (12%). The most sensitive species for those five chemicals (Aldrin, benomyl, fluridone, heptachlor, and Pydraul 50E) represented no one particular taxonomic group; they were *Orconectes nais*, *Pteronarcys californica*, *Chironomus plumosus*, coho salmon, and channel catfish.

An approach by Doherty (1983) and Kenaga (1978) was to correlate the LC50's of one species against another for all chemicals tested with those two species, by regression analyses. They each used four species of freshwater organisms (*Daphnia magna* or *D. pulex*, rainbow trout, fathead minnows, and bluegills) in their correlations. The correlations were good in Doherty's study ($r = 0.89$ to 0.97), indicating that the LC50 of one species can be estimated by the substitution of the LC50 of another species in appropriate regression equations. Kenaga found that correlations within the three fish species were also good ($r = 0.85$ to 0.99), but the correlation with *Daphnia* and rainbow trout was poor ($r = 0.56$) with 95% confidence limits of 2.2 to 3.3 orders of magnitude. We also found that the toxicity values for *D. magna* did not correlate well with those for other species (Table 1). This observation with daphnids confirms those by Kenaga (1978) and LeBlanc (1984)—all three data sets consisted of all or mostly

BEST AVAILABLE COPY

ED_005172C_00000035-00013

Table 1. *Interspecies correlations*[a] *for acute static LC50's (ng/L) with* Daphnia magna.

Species	n	Intercept (a)	Slope (b)	$\bar{Y}\pm$95% CL	Correlation coefficient (r)
Daphnia pulex	8	3.031	0.156	3.566±0.497	0.21
Simocephalus serrulatus	9	2.346	0.395[b]	3.690±0.322	0.70
Asellus brevicaudus	5	12.035	−1.963	4.581±1.231	0.83
Gammarus fasciatus	14	1.876	0.611[b]	4.289±0.457	0.82
Gammarus pseudolimnaeus	18	−0.462	1.010[b]	4.912±0.271	0.95
Claassenia sabulosa	5	6.949	−0.999	3.147±0.761	0.78
Pteronarcella badia	7	2.597	0.220	3.497±0.491	0.35
Pteronarcys californica	10	1.475	0.667[b]	4.151±0.938	0.65
Chironomus plumosus	25	0.802	0.846[b]	5.527±0.300	0.86
Coho salmon	8	7.056	−0.531	4.969±1.511	0.36
Cutthroat trout	7	4.943	0.265	6.061±1.159	0.43
Rainbow trout	31	3.208	0.428[b]	5.338±0.387	0.54
Goldfish	6	9.212	−0.941	5.981±1.578	0.43
Common carp	5	10.494	−1.480	5.590±2.032	0.67
Fathead minnow	26	4.743	0.236	5.890±0.478	0.32
Black bullhead	5	10.367	−1.402	5.723±2.529	0.56
Channel catfish	31	5.306	0.095	5.776±0.434	0.12
Green sunfish	6	7.652	−0.591	5.624±1.689	0.31
Bluegill	31	3.140	0.488[b]	5.480±0.432	0.60
Largemouth bass	6	9.548	−1.326	5.013±1.625	0.60
Fowler's toad	5	7.092	−0.438	5.405±0.495	0.65
Western chorus frog	5	7.165	−0.414	5.724±0.606	0.73

[a]$\text{Log } Y = a + b(\log X)$; Y = LC50 for other species, X = LC50 for *Daphnia magna*.
[b]Significantly different from zero ($P < 0.05$).

pesticides. The data of Doherty (1983) covered a broader range of chemical types.

For our data, correlations of toxicity values for rainbow trout, fathead minnows, or bluegills were good with those of some invertebrate species, better with those of other fish species, but best with those of other species within the same family (Tables 2 through 4). This general trend was also observed by LeBlanc (1984) and Suter and Vaughn (1985), who concluded that the more distant the relationship between two species, the more different their responses to chemical toxicity. Although toxicity of a chemical for one species can be predicted from the known toxicity of another, it must be remembered that correlation is not necessarily the same as causation in regression analyses, and that the error of estimation in correlation can therefore be large.

In initial hazard assessment testing tiers, Doherty (1983) concluded that *D. magna* should be the species of choice; Kenaga (1978) recommended a daphnid and a fish species; and Kimerle et al. (1983) recommended an alga, a daphnid, and a fish. Considering these studies and the present data, *Daphnia* and rainbow trout appear to be the first two taxa of choice for estimating freshwater animal sensitivity to chemicals in a first tier of testing. Other fish species could be used, but rainbow trout appeared to be the most sensitive overall. When a small margin of safety exists (less than one order of magnitude) between the expected or measured environmental concentration of a chemical and its acute toxicity, a more extensive data base is justified. As judged by our data, the inclusion of tests with amphipods and possibly stoneflies is indicated, to better approximate the toxicity for the most sensitive animal species.

On the basis of the work of Kimerle et al. (1983) and Blanck et al. (1984), algae should be further investigated as a third possible species for first tier testing. Additionally, interspecies correlations and other statistical and mathematical approaches must be expanded to include freshwater and marine species, and both the common and uncommon test species for which data exist.

ED_005172C_00000035-00014

Table 2. *Interspecies correlations*[a] *for acute static LC50's (ng/L) with rainbow trout.*

Species	n	Intercept (a)	Slope (b)	$\bar{Y} \pm 95\%$ CL	Correlation coefficient (r)
Daphnia magna	31	1.299	0.687[b]	4.961±0.493	0.54
Daphnia pulex	34	2.368	0.408[b]	4.502±0.492	0.35
Simocephalus serrulatus	35	2.979	0.298	4.555±0.502	0.25
Cypridopsis vidua	9	1.165	0.703[b]	4.390±0.311	0.85
Asellus brevicaudus	15	0.858	0.841[b]	4.854±0.398	0.80
Gammarus fasciatus	45	2.635	0.424[b]	4.855±0.397	0.36
Gammarus lacustris	19	2.219	0.357	4.121±0.489	0.44
Gammarus pseudolimnaeus	22	−0.968	1.067[b]	5.245±0.443	0.83
Claassenia sabulosa	10	1.296	0.434[b]	3.222±0.346	0.74
Pteronarcella badia	12	0.760	0.560[b]	3.729±0.329	0.90
Pteronarcys californica	42	0.990	0.632[b]	4.266±0.363	0.55
Chironomus plumosus	24	−0.135	0.960[b]	5.442±0.411	0.74
Coho salmon	24	0.022	0.985[b]	5.441±0.248	0.95
Chinook salmon	7	0.212	0.962[b]	6.047±0.284	0.98
Cutthroat trout	18	0.726	0.894[b]	5.803±0.178	0.96
Atlantic salmon	11	−0.163	1.026[b]	5.667±0.337	0.95
Brown trout	14	−0.023	1.019[b]	5.478±0.165	0.98
Brook trout	11	−0.804	1.130[b]	5.697±0.186	0.98
Lake trout	8	0.654	0.872[b]	5.627±0.184	0.99
Goldfish	15	1.258	0.878[b]	5.508±0.463	0.86
Common carp	8	1.511	0.832[b]	5.734±0.824	0.78
Fathead minnow	61	0.796	0.938[b]	5.961±0.134	0.94
Black bullhead	8	2.203	0.708[b]	5.553±0.972	0.79
Channel catfish	78	1.391	0.802[b]	5.906±0.160	0.82
Green sunfish	13	1.137	0.849[b]	5.408±0.255	0.96
Bluegill	101	0.135	1.005[b]	5.800±0.084	0.96
Largemouth bass	15	0.258	0.983[b]	5.611±0.339	0.96
Yellow perch	11	1.236	0.834[b]	5.391±0.366	0.94
Fowler's toad	12	3.563	0.456[b]	5.537±0.337	0.70
Western chorus frog	10	3.796	0.448[b]	5.904±0.300	0.85

[a]Log $Y = a + b(\log X)$; Y = LC50 for other species, X = LC50 for rainbow trout.
[b]Significantly different from zero ($P < 0.05$).

Families. The relative sensitivity of 34 families of aquatic organisms tested with 203 chemicals varied greatly. The maximum number of families (15) was tested with DDT, endrin, and malathion; they are listed here in descending order of sensitivity: Perlidae, Pteronarcidae, Gammaridae, Daphnidae, Centrarchidae, Palaemonidae, Salmonidae, Astacidae, Cypridae, Asellidae, Ictaluridae, Cyprinidae, Rhagionidae, Bufonidae, and Hylidae. However, some of the families were represented by only one species and others by many. A more complete representation might change the ranking. Although additional comparisons with smaller numbers of families and greater numbers of chemicals altered the ranking to some degree, the overall trend was maintained (e.g., insects were the most sensitive, followed by crustaceans, fishes, and amphibians).

Sensitivity of crustaceans was evaluated with six families and five insecticides, and four families and nine chemicals (eight insecticides and one herbicide). Gammaridae were the most sensitive to the five insecticides (carbaryl, DDT, endrin, malathion, and methoxychlor), followed by the Palaemonidae, Daphnidae, Astacidae, Cypridae, and Asellidae. Sanders (1970) found a different order of sensitivity with herbicides, reporting that Daphnidae were the most sensitive, followed in descending order by Cypridae, Palaemonidae, Asellidae, Gammaridae, and

BEST AVAILABLE COPY

ED_005172C_00000035-00015

Table 3. *Interspecies correlations*[a] *for acute static LC50's (ng/L) with fathead minnows.*

Species	n	Intercept (a)	Slope (b)	$\bar{Y} \pm 95\%$ CL	Correlation coefficient (r)
Daphnia magna	26	2.241	0.442	4.841±0.636	0.33
Daphnia pulex	22	5.741	−0.311	3.996±0.537	0.34
Simocephalus serrulatus	22	5.969	−0.336[b]	4.054±0.455	0.43
Cypridopsis vidua	8	2.649	0.314[b]	4.253±0.381	0.73
Asellus brevicaudus	11	1.446	0.567[b]	4.511±0.404	0.82
Gammarus fasciatus	24	4.744	−0.086	4.270±0.512	0.10
Gammarus lacustris	14	3.107	0.167	4.052±0.457	0.34
Gammarus pseudolimnaeus	14	−1.911	1.123[b]	5.060±0.690	0.80
Claassenia sabulosa	9	1.692	0.303[b]	3.274±0.407	0.71
Pteronarcella badia	11	1.859	0.286[b]	3.434±0.180	0.87
Pteronarcys californica	27	2.012	0.326[b]	3.837±0.413	0.42
Chironomus plumosus	18	−0.888	1.082[b]	5.651±0.551	0.73
Coho salmon	17	−0.856	1.038[b]	5.249±0.308	0.96
Cutthroat trout	13	−0.559	1.001[b]	5.731±0.471	0.86
Rainbow trout	61	−0.090	0.947[b]	5.555±0.126	0.95
Atlantic salmon	10	−1.772	1.198[b]	5.736±0.322	0.96
Brown trout	9	−2.225	1.259[b]	5.656±0.571	0.88
Brook trout	11	−1.329	1.134[b]	5.697±0.421	0.92
Goldfish	19	0.572	0.924[b]	5.732±0.189	0.97
Common carp	8	−0.040	0.988[b]	5.734±0.365	0.96
Black bullhead	8	1.411	0.767[b]	5.553±0.664	0.91
Channel catfish	48	0.954	0.832[b]	5.865±0.185	0.88
Green sunfish	12	0.695	0.842[b]	5.472±0.333	0.95
Bluegill	62	0.018	0.954[b]	5.675±0.148	0.93
Largemouth bass	15	−0.433	0.972[b]	5.370±0.423	0.93
Yellow perch	7	−0.110	0.967[b]	5.229±0.322	0.99
Fowler's toad	12	5.075	0.087	5.492±0.399	0.16
Western chorus frog	9	4.776	0.192	5.733±0.391	0.50

[a]Log $Y = a + b(\log X)$; Y = LC50 for other species, X = LC50 for fathead minnows.
[b]Significantly different from zero ($P < 0.05$).

Astacidae. The nine-chemical comparison resulted in a similar trend of decreasing sensitivity, in the following order: Gammaridae, Daphnidae, Cypridae, and Asellidae. The Cypridae and Asellidae were significantly more sensitive than Daphnidae only to lindane. The Streptocephalidae were tested only with phosmet; the LC50 was higher than that for the other invertebrates tested, but below the range of LC50's for fishes.

Among families, insects were generally the most sensitive, due to the high sensitivity of stoneflies (Perlidae, Perlodidae, and Pteronarcidae) and one family of mayflies (Baetidae). The LC50's for species in other families of insects (Ephemeridae, Coenagrionidae, Gomphidae, Hydropsychidae, Limnephilidae, Culicidae, Rhagionidae, Tendipedidae, and Tipulidae) were within the range of (or generally not statistically significant from) the invertebrate and vertebrate families more commonly tested. However, the LC50's for Tendipedidae were significantly less than those of other families tested within a chemical for fluoridone, metolachlor, oxamyl, piperonyl butoxide, and Santicizer 148.

Macek and McAllister (1970) reported on the sensitivity of fishes to nine insecticides: the Cyprinidae and Ictaluridae were the least sensitive, the Centrarchidae were intermediate, and the Salmonidae were the most sensitive; the Percidae ranked between the Centrarchidae and Salmonidae. In the present study, in tests with the four main fish

ED_005172C_00000035-00016

Table 4. *Interspecies correl~tions[a] for acute static LC50's (ng/L) with bluegills.*

Species	n	Intercept (a)	Slope (b)	$\bar{Y}\pm$95% CL	Correlation coefficient (r)
Daphnia magna	31	0.649	0.745[b]	4.785±0.529	0.61
Daphnia pulex	25	3.081	0.242	4.339±0.640	0.19
Simocephalus serrulatus	27	3.422	0.186	4.414±0.612	0.16
Cypridopsis vidua	7	1.767	0.593[b]	4.285±0.541	0.76
Asellus brevicaudus	13	0.607	0.887[b]	5.053±0.472	0.86
Gammarus fasciatus	47	1.863	0.546[b]	4.774±0.361	0.49
Gammarus lacustris	19	2.436	0.306	3.992±0.511	0.40
Gammarus pseudolimnaeus	22	−1.117	1.040[b]	5.220±0.386	0.87
Procambarus sp.	5	1.465	0.701	5.589±1.497	0.70
Claassenia sabulosa	10	1.580	0.368[b]	3.222±0.330	0.77
Pteronarcella badia	12	1.724	0.392	3.680±0.546	0.57
Pteronarcys californica	40	0.514	0.732[b]	4.430±0.398	0.63
Chironomus plumosus	25	0.649	0.811[b]	5.458±0.430	0.72
Coho salmon	24	0.222	0.933[b]	5.155±0.233	0.95
Chinook salmon	5	−0.511	1.083[b]	5.788±0.285	0.99
Cutthroat trout	22	0.692	0.876[b]	5.696±0.229	0.90
Rainbow trout	101	0.440	0.898[b]	5.635±0.079	0.96
Atlantic salmon	13	−0.229	1.028[b]	5.426±0.214	0.98
Brown trout	14	−0.691	1.141[b]	5.323±0.251	0.94
Brook trout	11	−0.886	1.139[b]	5.728±0.337	0.95
Lake trout	11	0.237	0.903[b]	5.004±0.197	0.98
Goldfish	17	1.295	0.846[b]	5.644±0.460	0.84
Common carp	8	1.951	0.746[b]	5.734±0.787	0.80
Fathead minnow	62	0.947	0.883[b]	5.931±0.152	0.92
Black bullhead	8	1.696	0.798[b]	5.553±0.883	0.85
Channel catfish	75	1.918	0.713[b]	5.870±0.171	0.78
Green sunfish	15	0.999	0.881[b]	5.486±0.195	0.97
Largemouth bass	15	0.051	1.003[b]	5.408±0.233	0.98
Yellow perch	12	0.078	1.029[b]	5.167±0.432	0.91
Fowler's toad	11	3.996	0.375	5.523±0.400	0.44
Western chorus frog	9	4.412	0.297	5.733±0.339	0.66

[a]Log $Y = a + b(\log X)$; Y = LC50 for other species, X = LC50 for bluegills.
[b]Significantly different from zero ($P \leq 0.05$).

families and 65 chemicals, Salmonidae were the most sensitive, followed by the Centrarchidae, Ictaluridae, and Cyprinidae—a ranking that confirms the observations of Macek and McAllister (1970). However, Percidae ranked between Centrarchidae and Ictaluridae when tests of all five families with 18 chemicals were compared. Five of the six LC50's for Esocidae were below those of five other families (Centrarchidae, Cyprinidae, Ictaluridae, Percidae, and Salmonidae), but the difference was significant only for azinphos methyl. In the few tests conducted with fish of other families, the LC50's for Poeciliidae were within the range for the other families tested and those for the Cichlidae were generally greater, but not significantly so. Only 38% of the fish tests with 65 chemicals showed significant differences among families, and 76% of those differences were between cyprinids and salmonids.

When test data from seven major families tested with 20 insecticides and 1 herbicide were compared, invertebrates were the most sensitive 95% of the time. The Pteronarcidae were the most sensitive, followed by Daphnidae, Gammaridae, Salmonidae, Centrarchidae, Ictaluridae, and Cyprinidae.

BEST AVAILABLE COPY

ED_005172C_00000035-00017

Classes. No one class was always the most sensitive or always the least sensitive. Overall, sensitivity to 163 chemicals decreased in the following order: Insecta, Crustacea, Osteichthyes, and Amphibia. These four classes occurred together in tests of 16 chemicals; Insecta were the most sensitive 50% of the time, followed by Crustacea (31%), Osteichthyes (19%), and Amphibia (0%). The comparisons were not balanced, since the Crustacea were represented by 12 species, Insecta by 21, Osteichthyes by 30, and Amphibia by 2. In other studies (Birge et al. 1980; Black et al. 1982), amphibians (pickerel frog, *Rana palustris*; African clawed frog, *Xenopus laevis*; Fowler's toad) were as tolerant as cyprinids, whereas the leopard frog (*R. pipiens*) and European common frog (*R. temporaria*) were as sensitive as salmonids. However, we believe that the overall trend would not be greatly changed by the addition of more species. In comparisons of Crustacea, Insecta, and Osteichthyes tested with 85 chemicals, the Insecta were the most sensitive 49% of the time, followed by Crustacea (27%) and Osteichthyes (24%). Crustacea were the second most sensitive 51% of the time, and Osteichthyes the least sensitive 55% of the time. Analysis of variance indicated that toxicity was not significantly different among classes for 32 of the 85 chemicals examined: 2 fungicides, 9 herbicides, 7 industrials, 12 insecticides, 1 molluscicide, and 1 surfactant.

Frequency Distribution of Chemical Toxicity

Frequency distributions of EC50's and LC50's tended to be bimodal for many species—the toxicities of insecticides were mainly in the <100 μg/L categories, and those of herbicides, fungicides, industrials, and other chemicals were in the >1,000 μg/L categories (Fig. 1*A*–1*D*). The lower mode contained almost all of the insecticides tested with invertebrates, whereas among fishes, insecticides were spread over more of the toxicity range. Insecticides occurring in the lower modes of fishes were mainly botanicals and organochlorines plus some carbamates and organophosphates. The distribution of chemical toxicities in larval frogs and toads was similar to that for fishes, within the confines of the few chemicals tested against amphibians. For all invertebrates and fishes, the higher mode consisted mainly of herbicides and industrials. The trends were the same as those found by Kenaga (1978); insecticides are generally more toxic to animals than are herbicides.

Frequency distributions of EC50's and LC50's among species (352 chemicals, 61 species) ranged over nine orders of magnitude, and 90% fell within five orders of magnitude (Fig. 1*E*). For all the compounds tested within a class, the ranges of EC50 or LC50 values were eight orders of magnitude for Crustacea, seven for Insecta, nine for Osteichthyes, and four for Amphibia (Fig. 1*E*). The frequency distributions for classes were similar to those for species, in that they were bimodal rather than normally distributed for Crustacea, Insecta, and Osteichthyes. The frequency was highest in the low mode (1.0–10 μg/L) for Insecta and in the high mode (1,000–10,000 μg/L) for fish; for Crustacea, frequencies in the two modes (10–100 and 1,000–10,000 μg/L) were about equal. The chemical distribution was similar to that for species, in that insecticides were mainly in the first mode and herbicides, industrials, and others in the second mode. However, insecticides were spread more evenly over the entire range of chemical toxicities for Insecta and Osteichthyes.

If large variations in species response to a variety of chemicals is desirable for differentiating between toxicities of compounds (Kenaga 1978), we can make recommendations for test species. Various chemical toxicities ranged over seven orders of magnitude for *Daphnia magna*, *D. pulex*, *Gammarus fasciatus*, and *G. pseudolimnaeus*; for all other invertebrates this range was six orders of magnitude or less. The range was nine orders of magnitude for rainbow trout and eight orders of magnitude for coho salmon, fathead minnows, and bluegills. The orders of magnitude for all other fish and amphibians ranged from three to seven. However, the data may be somewhat misleading, since not all species were tested with all chemicals. Also, an upper limit of testing (100–1,000 mg/L), though appropriate, may have reduced the full range of toxicities for the generally less sensitive species.

Static and Flow-through Tests

The data set consisted of 123 paired static and flow-through tests with 41 chemicals. Ratios of static to flow-through tests (96-h LC50's) used for analyses ranged from 0.12 to 8.5 (mean, 2.0). The ratios varied considerably among chemicals, but the variation was generally as great among species within a chemical. Static LC50's were more than 1.5× those of flow-through tests 53% of the time, and no differences (ratios of 0.5–1.5) were observed in 37%

ED_005172C_00000035-00018



Fig. 1. Distribution of various chemical toxicities within species and classes, and among all species. Vertical axis = percent frequency of EC50's or LC50's occurring within a toxicity range, and horizontal axis = toxicity range. *A.* Crustaceans: *Daphnia magna* (49 tests, 46 chemicals), *D. pulex* (42 tests, 42 chemicals), *Simocephalus serrulatus* (62 tests, 43 chemicals), *Gammarus fasciatus* (81 tests, 79 chemicals), *G. pseudolimnaeus* (28 tests, 28 chemicals). *B.* Insects: *Pteronarcys californica* (59 tests, 57 chemicals), *Chironomus plumosus* (36 tests, 35 chemicals). *C.* Fishes: rainbow trout (286 tests, 184 chemicals), fathead minnows (116 tests, 89 chemicals), channel catfish (128 tests, 107 chemicals), bluegills (277 tests, 163 chemicals), yellow perch (26 tests, 16 chemicals).

BEST AVAILABLE COPY

ED_005172C_00000035-00019



Fig. 1. *Continued. D.* Amphibians: Fowler's toad (15 tests, 14 chemicals), western chorus frog (10 tests, 10 chemicals). *E.* Classes and among all species: Crustacea (333 tests, 139 chemicals, 12 species), Insecta (152 tests, 91 chemicals, 20 species), Osteichthyes (1,144 tests, 253 chemicals, 26 species), Amphibia (25 tests, 17 chemicals, 2 species), among all species (1,656 tests, 352 chemicals, 61 species).

of the tests. The observation that 10% of the ratios were <0.5 could indicate that degradation or hydrolysis products more toxic than the parent compounds were being tested under static conditions.

The eight organochlorine insecticides had one of the highest average ratios (2.5; range, 0.66–6.4) as expected, but the six pyrethrins and pyrethroids had the same average ratio (2.5; range, 0.51–8.5)—which was not expected. However, since the toxicity of pyrethrins and pyrethroids did not decrease with aging (Tables 5 and 6), these chemicals may be somewhat persistent, having a higher static to flow-through ratio, as organochlorine insecticides do. The ratios for four carbamate insecticides (1.8; range, 0.12–3.8) and nine organophosphate insecticides (1.9; range, 0.27–7.5) were similar to those of the two fungicides (2.1; range, 0.75–3.6); the ratios for nine herbicides (1.4; range, 0.32–3.3) and three industrial compounds (1.4; range, 0.83–2.4) tested were not different from each other.

It was previously reported that static toxicity values can be multiplied by 0.71 to estimate values for flow-through tests (U.S. Environmental Protection Agency 1979). The factor of 0.71 was based on

ED_005172C_00000035-00020

the geometric mean of values for 24 pairs of tests on 12 chemicals. Our data for 123 paired tests and 41 chemicals indicate an average factor of 0.51, which varies with chemical type. The use of the factor of 0.51 would only be a crude estimate, however, since the ratios of static to flow-through 96-h LC50's varied by 1.8 orders of magnitude.

Aged Test Solutions

The biological activity of a toxicant, which is responsible for the killing power, decreases as the toxicant is deactivated biologically and chemically over time (Marking 1972). In the absence of adequate analytical methods to measure the half-life of some chemicals in water, Marking (1972) outlined an appropriate bioassay method. His method does not differentiate between chemical degradation, physical removal, and biological inactivation, but does estimate the total activity of a chemical and its degradation products, except as the chemical is removed. The deactivation index, derived by Marking and Dawson (1972) is obtained by dividing the LC50 of an aged test solution by the LC50 of a fresh test solution, and determines both decreases and increases in biological activity.

To determine how often the toxicity of an aged test solution differed from that of a fresh solution, we established a data set with 46 tests on 22 chemicals. Data for dinoseb, glyphosate, mexacarbate, picloram, pyrethrins, and pyrethroids (Table 5) were previously published by Folmar et al. (1979), Mauck et al. (1976, 1977), and Woodward (1976).

Although toxicity increased in some fish species and decreased in others within the same chemical (e.g., aminocarb, fenitrothion, and glyphosate), it appears that deactivation indices between about 0.5

Table 5. *Toxicity (96-h LC50, mg/L) of pesticides in fresh and aged solutions to aquatic organisms in soft water (40–44 mg/L $CaCO_3$) under static conditions.*

Chemical and species	pH	Toxicity of solution: Fresh	Toxicity of solution: Aged 7 days	Deactivation index[a]
Aminocarb				
Rainbow trout	7.2	0.14	0.12	0.86
Atlantic salmon	7.5	2.4	3.5	1.46
Brown trout	7.5	17	8.6	0.51
Bluegill	7.2	0.24	0.24	1.00
Carbaryl				
Isogenus sp.[b]	7.5	0.004	0.007	1.75
Cutthroat trout	7.3	6.8	4.0	0.59
2,4-D butyl ester				
Cutthroat trout	7.4	0.67	0.98	1.46
Lake trout	7.4	0.72	1.3	1.81
2,4-D dimethylamine salt				
Fathead minnow	7.4	760	760	1.00
2,4-D dodecyl/tetradodecyl amine salt				
Fathead minnow	7.4	9.1	2.2	0.24
2,4-D propylene glycol butyl ether ester				
Cutthroat trout	7.4	0.49	1.4	2.86
Lake trout	7.4	1.1	1.4	1.27
Dimethrin				
Bluegill	6.5	0.024	0.028	1.17
	7.5	0.030	0.028	0.93
	9.5	0.029	0.034	1.17
Dinoseb				
Cutthroat trout	7.4	0.071	0.16	2.25

BEST AVAILABLE COPY

ED_005172C_00000035-00021

Table 5. *Continued.*

Chemical and species	pH	Toxicity of solution: Fresh	Toxicity of solution: Aged 7 days	Deactivation index[a]
Diuron				
Cutthroat trout	7.0	1.5	12	8.00
Rainbow trout	7.5	3.5	4.2	1.20
Lake trout	7.0	2.7	3.2	1.18
d-trans allethrin				
Bluegill	6.5	0.056	0.040	0.71
	7.5	0.056	0.034	0.61
	9.5	0.060	0.074	1.23
EPTC				
Cutthroat trout	7.1	16	16	1.00
Fenitrothion				
Atlantic salmon	7.5	0.33	0.64[c]	1.94
Brown trout	7.5	2.4	1.7	0.71
Brook trout	7.5	1.9	1.7	0.89
Bluegill	7.2	4.3	4.4	1.02
Glyphosate				
Rainbow trout	7.4	9.0	7.6	0.84
Bluegill	7.4	4.0	5.6	1.40
Leptophos				
Rainbow trout	7.4	0.013	0.038	2.92
Methomyl				
Gammarus pseudolimnaeus[b]	7.2	1.0	0.34	0.34
Rainbow trout	7.2	1.4	1.5	1.07
Mexacarbate				
Coho salmon	7.5	23	15	0.65
Bluegill	6.5	23	18	0.78
	8.5	4.0	1.0	0.25
	9.5	1.0	0.92	0.92
Phosmet				
Chironomus plumosus[b]	7.4	3.3	>10[d]	> 3.03
Bluegill	7.2	0.39	>10[d]	>25.64
Phoxim				
Bluegill	7.5	1.0	1.0	1.00
Picloram				
Cutthroat trout	7.4	5.8	4.8	0.83
Pyrethrum				
Bluegill	6.5	0.041	0.066	1.61
	7.4	0.053	0.056	1.06
	9.5	0.087	0.090	1.03
s-Bioallethrin				
Bluegill	6.5	0.039	0.034	0.87
	7.5	0.033	0.044	1.33
Trichlorfon				
Cutthroat trout	7.5	1.7	0.47	0.28

[a]Deactivation index = $\frac{\text{LC50 aged solution}}{\text{LC50 fresh solution.}}$

[b]48-h LC50.
[c]Solutions aged 14 days.
[d]Solutions aged 4 days.

ED_005172C_00000035-00022

Table 6. *Distribution of toxicity change of pesticides in test solutions aged for 7 days under acute static conditions.*

Pesticide type	Pesticides (n)	Tests (n)	Toxicity change distribution (%)[a]		
			None	Decrease	Increase
All chemicals	22	46	69	22	9
All chemicals (pH effects removed)	22	37	68	24	8
Herbicides	9	14	64	29	7
Insecticides	13	23	69	22	9
Carbamates	4	10	80	10	10
Organophosphates	5	9	44	44	11
Pyrethrins and pyrethroids	4	4	100	0	0

[a]Change in toxicity due to aging was determined by the ratio of aged to fresh solutions, and the ratios were categorized by change: none, 0.5–1.5; increase, <0.5; decrease, >1.5.

and 1.5 are probably within the range of normal experimental variation. If this assumption is true, aging did not affect chemical toxicity in 69% of the tests, decreased it in 22%, and increased it in 9% (Table 6). Stability of the chemicals was indicated where no change occurred. The toxicity of aged solutions decreased most in herbicides and increased most in carbamate insecticides.

Toxicities of dinoseb, EPTC, glyphosate, and picloram solutions aged for 7 days remained essentially unchanged. Although the toxicity of dinoseb appeared to decrease slightly and that of picloram to increase, continued aging and testing beyond 7 days showed no real change in toxicity. After 4 weeks of aging, deactivation indices for dinoseb, EPTC, and picloram were 1.22, 1.06, and 1.02, respectively. Conversely, diuron and the 2,4-D group of herbicides continued to decrease in toxicity after 3 to 4 weeks of aging. The one exception was the dodecyl/tetradodecylamine salt of 2,4-D, which increased in toxicity after 7 days of aging.

Essentially no changes occurred in deactivation studies with the carbamate insecticides aminocarb and methomyl. However, tests on *Gammarus pseudolimnaeus* with methomyl indicated some increase in toxicity. Exposure of fish to carbaryl and mexacarbate tended to show that the toxicity of aged solutions exceeded that of fresh solutions. Mauck et al. (1977) reported that mexacarbate hydrolyzes to chemicals that are more toxic than the parent compound before further hydrolyzing to less toxic chemicals. This phenomenon may have been the same for cutthroat trout exposed to carbaryl, since carbaryl hydrolyzes at pH 7.0 and above in aqueous solutions (Aly and El-Dib 1971). However, if hydrolysis products were formed, their toxicity to the stonefly *Isogenus* decreased after carbaryl was aged.

Studies on pyrethrum deactivation demonstrated no change in toxicity with aging at pH 7.5 and 9.5, but deactivation occurred at pH 6.5. Toxicity of dimethrin, s-bioallethrin, and *d*-trans allethrin was unchanged.

The biological activity of the organophosphate insecticides leptophos (phosphonothioate) and phosmet (phosphorodithioate) decreased considerably, whereas that of trichlorfon (phosphonate) increased. The responses of fishes to aged phosphorothioates (fenitrothion and phoxim) varied, but tended toward no change in toxicity. However, Eisler (1970) reported that 4 days of aging increased the toxicity of another phosphorothioate (methyl parathion) by 6.6×.

Deactivation studies are useful for determining biological activity of aged solutions of chemicals to aquatic organisms, but toxicity changes should not be of great concern in acute static tests. Chemical toxicity increased (by a factor of 2 or more) only 11% of the time and decreased 13% of the time. Toxicity never increased by more than a factor of 4; however, it decreased by as much as a factor of 25 in one test with phosmet, and the solutions were aged for 4 days—the same duration as that for acute static toxicity tests.

BEST AVAILABLE COPY

ED_005172C_00000035-00023

Factors Affecting Toxicity

Temperature

The effect of temperature on toxicity of chemicals to aquatic organisms was reviewed by Cairns et al. (1975*a*,*b*), Johnson (1968), Sprague (1970), and Tucker and Leitzke (1979). Temperature and toxicity are positively correlated for most chemicals: toxicity increases as temperature increases (positive temperature coefficient). Conversely, toxicity of DDD, DDT, and methoxychlor are negatively correlated with temperature and thus have negative temperature coefficients (Tucker and Leitzke 1979). Mauck et al. (1976) demonstrated a negative temperature coefficient in fishes for pyrethrins and several of the pyrethroids as well.

Differences in toxicity due to temperature have been attributed to differences in respiration rate (Weiss and Botts 1957), chemical absorption (Wuhrmann 1952), and excretion and detoxification of chemicals (Sprague 1970). Gammon et al. (1981) reported that the neurophysiological effects of *d*-trans allethrin on the cockroach (*Periplaneta americana*) are excitatory at both high and low temperatures, the temperature-dependent differences being in the types of nerves directly affected. Both peripheral and central nervous systems are affected at 32°C, but only the peripheral system at 15°C. Gammon et al. (1981) concluded that the negative temperature coefficient of allethrin toxicity could be a result of increases in sensitivity of peripheral nerves at low temperatures.

To determine if temperature affects acute toxicity in this data base, we conducted covariance analysis on 48 chemicals and 90 temperature tests for fishes (Table 7). Slopes in 26 tests with 19 chemicals differed significantly from zero. The ratio of the highest to lowest LC50 within a temperature test averaged 9.8 (ratio range, 3.1 to 51) among the 19 chemicals affected. Temperature coefficients were negative for DDT, dimethrin, and methoxychlor. There were also negative temperature coefficients (though not significantly different from zero) for pyrethrins and pyrethroids (fenvalerate, permethrin, and pyrethrum); however, the coefficient was positive for *d*-trans allethrin.

The negative slopes (positive temperature coefficient) were variable, ranging from −0.0218 for diuron to −0.1002 for chlorpyrifos, but were much more consistent among species within a chemical and within chemical groupings. When the slopes were compared with each other (Table 7), they were never significantly different among species within a chemical. Thus, differences were chemically rather than biologically specific. The slopes for the carbamates (aminocarb, carbaryl, and methomyl) were statistically similar, as were those for the organochlorines chlordecone and endrin. Among slopes for the herbicides, only those for diuron and trifluralin differed significantly. The slopes for the organophosphate insecticides varied to some degree, but the variation was reduced by further categorization. Among the slopes for phosphorothioates or phosphorodithioates, only two (phosphorodithioate) differed; the slope for azinphos-methyl and yellow perch was significantly different from that for malathion and bluegills. The slopes for malathion and parathion (phosphorothioate) differed significantly from most tests with trichlorfon, a phosphonate. The slope for the one organophosphate herbicide, merphos (phosphorotrithioate), was not significantly different from the slopes of any of the organophosphate insecticides. The slopes of the three chemicals (DDT, dimethrin, and methoxychlor) having negative temperature coefficients were also not significantly different from each other.

The temperature coefficient (Q_{10}) of respiration generally ranges from 2 to 4, indicating that a 10°C rise in temperature increases the rate of the reaction twofold to fourfold in cold-blooded animals (Giese 1968). Although Cairns et al. (1975*a*,*b*) included this well-known generalization in their review, they did not relate Q_{10} to the effects of temperature on chemical toxicity to aquatic organisms. The application of the Q_{10} concept is appropriate and useful for biologic effects, but was originally applied only to rate constants for nonbiotic reactions and processes. Thus it should apply to rates of reaction of factors that affect bioavailability under static test conditions (e.g., hydrolysis, microbial degradation, photolysis, solubility, sorption–desorption, and vaporization). If test concentrations are not measured, as in the present data, bioavailability could contribute significantly to temperature effects. In regard to the test organisms, the Q_{10} concept may apply not only to the rate of reaction of the toxicant at the site(s) of action, but also to residue dynamics within the organism (e.g., uptake, metabolism, and depuration). We analyzed the slopes (Table 7) to determine if the effects of temperature on toxicity were predictable within chemicals having positive or negative temperature coefficients, and whether the effects were related to the Q_{10} concept.

BEST AVAILABLE COPY

ED_005172C_00000035-00024

Table 7. *Effect of temperature on acute static toxicity (96-h LC50) of organic chemicals to fishes.*[a]

Chemical and species	Temperature range (°C)	LC50's (n)	LC50 range (μg/L)	Intercept (a)	Slope[b] (b)	r^2 (%)
Aminocarb						
Brook trout	7–17	3	2,600–9,400	4.3611	−0.0558	100
Yellow perch	7–22	4	1,700–11,700	4.4306	−0.0518	94
Azinphos-methyl						
Yellow perch	7–22	3	2.4–40	2.1678	−0.0820	100
Carbaryl						
Brook trout	7–17	3	680–2,100	3.9842	−0.0645	92
Yellow perch	7–22	4	1,200–13,900	4.6053	−0.0678	98
Chlordecone						
Redear sunfish	7–29	5	29–140	2.3558	−0.0304	100
Chlorpyrifos						
Rainbow trout	2–18	4	1.0–51	1.9361	−0.1002	96
DDT						
Bluegill	7–29	5	1.6–5.8	0.0282	0.0278	89
Dimethrin						
Bluegill	12–22	3	21–85	0.6517	0.0601	94
Diuron						
Bluegill	7–29	5	2,800–9,500	4.2199	−0.0218	74
Endothall, Aquathol K						
Bluegill	7–22	4	343–1,740[c]	6.6683	−0.0467	86
Endrin						
Bluegill	7–29	5	0.19–0.73	0.1008	−0.0258	90
Malathion						
Bluegill	7–29	5	20–87	2.2149	−0.0285	92
Merphos						
Bluegill	12–22	3	8,000–32,000	5.1043	−0.0546	100
Methomyl						
Bluegill	12–27	9	430–2,000	3.6418	−0.0394	75
Methoxychlor						
Brook trout	7–17	12	7.0–30	0.7625	0.0282	42
Parathion						
Bluegill	7–29	5	18–400	2.7551	−0.0421	37
Phosmet						
Bluegill	10–25	8	60–560	3.3188	−0.0664	92
Phoxim						
Coho salmon	7–17	3	215–1,160	3.5472	−0.0732	98
Bluegill	12–22	3	82–432	3.5074	−0.0722	100
Trichlorfon						
Rainbow trout[d]	7–17	21	190–11,400	3.7995	−0.0696	41
Atlantic salmon	7–17	3	580–4,100	4.0731	−0.0849	77
Brook trout	7–17	3	1,100–9,400	4.5889	−0.0932	98
Bluegill[d]	7–17	9	1,720–50,000	4.8566	−0.0867	50
Trifluralin						
Rainbow trout	2–18	7	14–330	2.6653	−0.0777	93
Bluegill	7–29	8	8.4–400	3.2144	−0.0635	70

[a]Log 96-h LC50 in μg/L = $a+b$ (temperature in °C).
[b]Significantly different from zero ($P < 0.05$).
[c]mg/L.
[d]Combined temperature effects at four pH's (6.5, 7.5, 8.5, and 9.0).

BEST AVAILABLE COPY

ED_005172C_00000035-00025

The antilog of the slope in the equation log LC50 = $a + b$ (temperature in °C) is, in essence, a factor of change in 96-h LC50 per 1°C. The slopes (Table 7) were multiplied by 10 to derive the log change per 10°C. The average for the 23 chemicals having positive temperature coefficients was 0.6082 (95% confidence limits [CL], 0.5140–0.7024). Thus, the log 96-h LC50 at a 10°C higher temperature can be predicted by subtracting 0.6082 from the log 96-h LC50 of an acute test previously conducted, or 0.6082 would be added for a 10°C decrease in temperature. This is equal to the LC50 increasing by a factor of 4.06 per 10°C decrease in temperature, or decreasing by a factor of 0.25 per 10°C increase. Similar factors for chemicals having negative temperature coefficients could be derived if adequate data were available.

The factor 4.06 (95% CL, 3.26–5.04) tends to be above the coefficients of 2 to 4 mentioned for the Q_{10} concept. Cairns et al. (1975*a*) stated that temperature interactions with organophosphates may be very complex due to the change in acetylcholinesterase (AChE) activity from temperature alone. The results of Hogan (1970) confirmed the increase in AChE activity with increased temperature for bluegills (log AChE = 2.4539 + 0.0165 [°C] or a factor of 1.46 increase per 10°C), and the slopes for the organophosphates tended to be greater than those for other chamicals (Table 7). The average slope was 0.7113 (95% CL, 0.5836–0.8390) for organophosphates alone, and 0.4956 (95% CL, 0.3775–0.6137) for the remaining chemicals.

Increased temperature, by accelerating AChE activity, may simultaneously increase the rate of AChE inhibition by organophosphates. By inspection, the interaction appears to be slightly more than additive, but further research is required to better delineate that interaction. The carbamates did not seem to be affected as much as the organophosphates, on the average, and this decreased effect may be due to differences in modes of action between carbamates and organophosphates in AChE inhibition (O'Brien 1967).

Most reports addressing temperature effects showed that most chemicals, including inorganics, were within or approximated a 2× to 4× increase or decrease in LC50 per 10°C change (Cairns et al. 1975*a*,*b*; DeGraeve et al. 1980; Hashimoto and Nishiuchi 1981; Macek et al. 1969; Mauck et al. 1976; Sanders 1969; Sanders et al. 1983; Schoettger 1970; Schoettger and Mauck 1978; Smith and Heath 1979; Sullivan 1977; Woodward 1976). This observation implies that the effect of temperature on 96-h LC50's for fish generally conforms to the Q_{10} concept.

Although the interactions between temperature and toxicity are not completely understood, interim approaches can be used to estimate effects. When a 96-h LC50 for a chemical has been determined for a fish species (96-h $LC50_1$), the following equations can be used for estimating the effect of a ±10°C change on the 96-h LC50's of chemicals having positive temperature coefficients:

Organophosphates:

$$\log 96\text{-h } LC50_2 = \log 96\text{-h } LC50_1 \pm 0.7113$$

Other chemicals:

$$\log 96\text{-h } LC50_2 = \log 96\text{-h } LC50_1 \pm 0.4956$$

Temperature effects on chemicals with negative temperature coefficients can also be estimated if the temperature coefficient is known. Additional research on temperature and toxicity of chemicals to invertebrates and structure–activity relations of chemicals with temperature is strongly indicated.

Hydrogen Ion Concentration (pH)

Depending on the chemical, the 96-h LC50 either increases or decreases with changes in pH (Johnson 1968; Sprague 1970; Tucker and Leitzke 1979). The toxicity of ionizable organic chemicals (weak acids and bases) is greatly affected by pH—increases in pH decrease the toxicity of acids but increase that of bases (Crandall and Goodnight 1959; Folmar et al. 1979; Goodnight 1942; Holcombe et al. 1980; Kaila and Saarikoski 1977; Kobayashi and Kishino 1980; Konemann and Musch 1981; Levy and Gucinski 1964; Marking 1975; Marking and Olson 1975; McLeay et al. 1979; Saarikoski and Vilukela 1981; Schoettger and Julin 1969; Sills and Allen 1971; Woodward 1976; Zanella 1983). The toxicities differ because un-ionized forms diffuse more readily across fish gills (Maren et al. 1968). Ionic pesticides such as diquat and paraquat might also be influenced by pH (Weber 1972).

It has been generally believed that carbamates and organophosphates undergo rapid hydrolysis under alkaline conditions; therefore, toxicity decreases due to the presence of less toxic hydroxylated products of the parent compounds. O'Brien (1967) reported that organophosphate hydrolysis products are generally poor inhibitors of acetylcholinesterase (AChE); however, inhibitory potency is not reduced in some hydrolysis products, and may even be increased (O'Brien 1967; Schoettger and Mauck 1978; Tucker and Leitzke 1979; Woodward and Mauck

BEST AVAILABLE COPY

ED_005172C_00000035-00026

Table 8. *Effect of pH on acute static toxicity (96-h LC50) of organic chemicals to aquatic organisms.*[a]

Chemical and species	pH range	LC50's (n)	LC50 range (μg/L)	Intercept (a)	Slope[b] (b)	r^2 (%)
Aminocarb						
Rainbow trout	7.5–9.5	3	5,000–21,000	6.8928	−0.3116	84
Bluegill	6.5–9.5	4	1,000–10,500	6.4465	−0.3272	84
Yellow perch	6.5–9.0	4	425–7,000	7.5024	−0.5023	69
Benomyl						
Rainbow trout	6.5–8.5	3	160–880	−0.4856	0.3702	82
Bluegill	6.5–8.5	3	1,200–6,400	0.5642	0.3461	71
Carbaryl						
Cutthroat trout	6.5–8.5	3	970–7,100	6.3612	−0.3561	
Yellow perch	6.5–9.0	4	350–4,200	7.5810	−0.5416	80
2,4-D dodecyl/tetradodecyl amine salt						
Fathead minnow	6.5–8.5	6	1,900–8,000	5.9166	−0.2930	86
Dinoseb						
Cutthroat trout	6.5–8.5	3	41–1,350	−3.7844	0.7588	96
Lake trout	6.5–8.5	3	32–1,440	−4.4295	0.8266	91
Diquat						
Bluegill	6.5–9.5	4	115,000–498,000	7.3810	−0.2312	92
Glyphosate						
Rainbow trout	6.5–9.5	4	1,400–7,600	5.2671	−0.2262	66
Mexacarbate						
Coho salmon	6.5–9.0	3	4,000–23,000	7.0636	−0.3702	73
Cutthroat trout	6.0–9.0	3	630–15,800	6.2523	−0.3364	46
Bluegill	6.5–9.5	5	600–22,900	8.5980	−0.5746	86
Phosmet						
Rainbow trout	6.5–9.5	5	105–4,700	−2.1843	0.5711	85
Bluegill	6.5–8.5	3	22–640	−3.7051	0.7319	97
Trichlorfon						
Pteronarcella badia	6.5–8.5	3	5.3–100	6.3414	−0.6379	90
Cutthroat trout	6.5–8.5	3	375–4,750	7.5694	−0.5513	99
Rainbow trout[c]	6.5–9.0	12	210–11,400	6.6397	−0.4337	49
Atlantic salmon	6.5–8.5	4	300–4,400	7.5368	−0.5832	81
Brook trout	6.5–9.0	7	240–9,200	8.6374	−0.6532	85
Bluegill[c]	6.5–9.0	12	1,720–50,000	8.0379	−0.4904	56

[a]Log 96-h LC50 in μg/L = $a+b$(pH).
[b]Significantly different from zero ($P < 0.05$).
[c]Combined pH effects at three temperatures (7, 12, and 17°C).

1980). Many of the hydroxylated products of carbamates have pronounced AChE inhibitory activity (O'Brien 1967; Mauck et al. 1977; Schoettger and Mauck 1978; Woodward and Mauck 1980).

Covariance analysis was conducted on 49 chemicals and 100 pH tests (pH range, 6.0–9.5) with fishes and a few aquatic invertebrates. Slopes significantly different from zero were observed in 23 of the tests on 10 chemicals (Table 8). Although pH significantly affected the toxicity of only 20% of the chemicals, it caused greater average changes in 96-h LC50's than did any other factor examined. The ratio of the highest to lowest LC50 within a pH test averaged 16 (range, 4.2 to 45) among the 10 chemicals affected. The slopes were relatively consistent among species within a chemical, indicating chemical rather than biological differences. No significant effects of pH on acute toxicity were found for the nine industrials, two organochlorine insecticides, or five pyrethrin and pyrethroid insecticides tested, although five of the industrials were organophosphate compounds.

BEST AVAILABLE COPY

ED_005172C_00000035-00027

The toxicities of only 2 of the 11 organophosphate insecticides—phosmet and trichlorfon—were significantly affected by pH. The toxicity of phosmet decreased and that of trichlorfon increased within a change of 2 to 3 pH units. O'Brien (1967) reported that phosphates hydrolyze more readily than their corresponding phosphorothionates under alkaline conditions; our data agree, in that eight of the nine organophosphates not affected by pH were thionates of one form or another. The only exception was the phosphate dichlorvos, which is structurally similar to trichlorfon. The 96-h LC50's of dichlorvos for cutthroat trout were the same (170 μg/L) from pH 6.0 to 9.0, and Eisler (1970) found that dichlorvos aged for 4 days was slightly less toxic than fresh material to mummichogs (*Fundulus heteroclitus*). Either dichlorvos is hydrolyzed more slowly than trichlorfon, or the toxicity of the dichlorvos hydrolysis products differs little from that of the parent compound. Trichlorfon rapidly hydrolyzes to dichlorvos under alkaline conditions (O'Brien 1967), and dichlorvos was 2.6 to 350× more toxic than trichlorfon to aquatic organisms. Phosmet, the only thionate significantly affected by pH, differs from the other thionates in being a phthalimide. It is rapidly metabolized by animals to phthalamic acid by amidase cleavage and subsequent hydrolysis (O'Brien 1967).

Carbamate toxicity increased with pH, indicating the formation of toxic hydrolysis products under alkaline conditions (Table 8). Effects of pH were significant with three methyl carbamate insecticides, but not with methomyl (thioacetimidate). However, the toxicity of benomyl, a carbamate fungicide with a chemical structure (benzimidazole) totally different from that of the insecticides, decreased significantly with increasing pH.

The effects of pH on carbamate and organophosphate insecticides corresponded to the age of the test solutions. Toxicity decreased or increased with solution age (Tables 5 and 6) and with increased pH. Thus the impact of pH on toxicity of these chemicals appears to be primarily a hydrolytic phenomenon, although aging does not necessarily involve only hydrolysis.

Many of the herbicides tested were dissociable organic chemicals, but only 4 of 17 were significantly affected by pH (Table 8). Of these four, diquat is cationic and 2,4-D, dinoseb, and glyphosate are dissociable. The toxicity of dinoseb decreased, as would be expected, but the toxicities of dodecyl/tetradodecyl amine salt of 2,4-D and glyphosate increased. When dissolved in water, organic amines convert partly to ions (O'Brien 1967; Sills and Allen 1971), and the increased accumulation of the un-ionized form of dodecyl/tetradodecyl amine at higher pH's could oppose the expected effects of pH on 2,4-D. The glyphosate tested was formulated, and the increased toxicity was due to the surfactant (MON 0818) in the formulation (Folmar et al. 1979). The acidic technical glyphosate was more toxic at pH 6.5 than at pH 9.5. The toxicity of most of the acidic herbicides decreased with increasing pH, as would be expected—though the decreases were not statistically significant.

Predicting the effects of pH on acute toxicity of organic chemicals is difficult because the impact of the effects is varied. We agree with Saarikoski and Viluksela (1981), who reported that pH effects have been poorly analyzed for predicting the toxicity and accumulation of ionizable organic chemicals or the toxicity of those that hydrolyze. Before the effects of pH on bioavailability and toxicity can be adequately predicted, more research relating pH to structure–activity interactions is needed, in which properly designed replications of toxicity tests and appropriate statistical and mathematical analyses are used.

Hardness

Water hardness is known to affect chemical toxicity. Inglis and Davis (1972) and Sprague (1970) reviewed the effects of hardness on chemical toxicity to aquatic organisms. Although the results varied, the toxicity of most inorganic and some organic chemicals was affected by water hardness. We agree with the intimations of Inglis and Davis (1972) and Pickering and Henderson (1964), who found that hardness had little effect on toxicity of organic chemicals, and that any changes were probably due to differences in pH.

Analysis of covariance on 39 tests with 25 organic chemicals (herbicides, insecticides, and solvents) and fishes indicated that the slopes of only two chemicals, dimethylamine and dodecyl/tetradodecyl amine salts of 2,4-D, differed significantly from zero. The increase in hardness (from 12 to 250–300 mg/L as $CaCO_3$) caused a 2.8× increase in the LC50 of the dimethylamine salt (96-h LC50 range, 285 to 800 mg/L) and a 2.1× increase in that of the dodecyl/tetradodecyl amine salt (96-h LC50 range, 1.9 to 4.0 mg/L). The increase in LC50 values was probably due to the increase in pH (1.5 units) between the very soft and very hard test waters. Because 2,4-D is a weak acid (pK_a = 2.80; Weber

ED_005172C_00000035-00028

Table 9. *Distribution of toxicity change of technical pesticides of various formulations under acute static conditions.*

Formulation	Pesticides (n)	Tests (n)	Toxicity change distribution (%)[a] None	Decrease	Increase
All Formulations	48	161	57	11	32
Emulsifiable concentrate	23	62	65	11	24
Granular	4	10	50	10	40
Liquid	10	35	66	3	31
Wettable powder	16	39	56	13	51
Other	3	15	13	20	67
Oil disperson	1	3	0	100	0
Oil soluble concentrate	1	11	9	0	91
Spray concentrate	1	1	100	0	0

[a]Change in toxicity due to formulation was determined by the ratio of technical to formulated material, and the ratios were categorized by change: none, 0.5–1.5; decrease, <0.5, increase, >1.5.

1972), its toxicity should decrease in alkaline water. However, toxicity of the dodecyl/tetradodecyl amine salt was the only one of five 2,4-D compounds that increased (rather than decreased) with increasing pH (Table 8), and was the only one of four 2,4-D compounds that increased in toxicity with aging (Table 5). The dodecyl/tetradodecyl amine group may be the main factor in toxicity differences, and may mask the effect of 2,4-D.

Hardness-related toxicity changes of carbamate and organophosphate insecticides have been indicated by others (Schoettger and Mauck 1978; Woodward and Mauck 1980). In these types of chemicals, toxicity changes could be caused by the hydrolysis products, which are formed more rapidly under alkaline pH, as previously discussed.

Formulation

The effects of formulation were determined for 48 pesticides (Table 9). Ratios of 48-h EC50's or 96-h LC50's of technical grade pesticides to those of formulated materials were compared ($n = 161$). A ratio greater than 1.0 indicated increased toxicity of the formulated material and a ratio less than 1.0 indicated decreased toxicity. However, ratios between 0.5 and 1.5 are probably within the range of normal experimental variation, since this range frequently occurred among groups of species (invertebrates or fishes) within a chemical. Overall, toxicity was not affected 57% of the time, decreased 11% of the time, and increased 32% of the time. The average ratios (ranges in parentheses) were as follows: emulsifiable concentrate, 2.5 (0.10 to 24); granular, 1.7 (0.34 to 5.0); liquid, 3.6 (0.08 to 33); oil dispersion, 0.30 (0.14 to 0.47); oil soluble concentrate, 69 (1.2 to 318); spray concentrate, 0.59 (one test); and wettable powder, 1.4 (0.06 to 5.1).

Among the formulations in which toxicity increased, toxicity doubled 61% of the time and increased within a factor of five 80% of the time. The ratios exceeded one order of magnitude in 10% of the tests involving aminocarb, cyprazine, glyphosate, and temephos; liquid formulated aminocarb increased in toxicity as much as 318×. The toxicities of wettable powder and oil dispersion formulations were frequently less than the toxicity of technical material—the wettable powder probably due to chemical adsorption to talc in the formulation, and the oil dispersion formulation due to an altered octanol–water partition coefficient in the presence of petroleum hydrocarbons (Stecher 1983). Hashimoto and Nishiuchi (1981) reported that, in general, emulsifiable concentrates were the most toxic, followed by technical materials, wettable powders, dust, and granules. Pickering et al. (1962) also found emulsifiable concentrates to be slightly more toxic than technical materials. In our study, formulations ranked in order of most toxic to least toxic were oil soluble concentrate, liquid, emulsifiable concentrate, granular, wettable powder, spray concentrate, and oil dispersion.

Materials added to technical chemicals in the preparation of formulations should not be regarded, as they often are, as "inert" ingredients with little or no biological activity. Although the effects of "inerts" on acute toxicity of technical chemicals are

BEST AVAILABLE COPY

ED_005172C_00000035-00029

Table 10. *Diet effects on the acute static toxicity of organic chemicals to rainbow trout.*[a]

Chemical and diet	Weight[b] (g)	96-h LC50 (µg/L)	
		LC50	95% Confidence limits
Antimycin A			
Glencoe	1.10	0.032	(0.027–0.038)
Ewos	0.80	0.016	(0.012–0.021)
High protein (45%)	1.10	0.052	(0.044–0.062)
Fish protein concentrate	0.75	0.024	(0.020–0.029)
Carbaryl			
Glencoe	1.10	1,900	(1,590–2,260)
Ewos	0.80	2,080	(1,580–2,730)
High protein (45%)	1.10	2,300	(1,500–3,500)
Fish protein concentrate	0.80	1,360	(900–1,860)
Chlordane			
Oregon Moist	1.00	8.2	(6.1–11)
Glencoe	1.10	9.1	(4.8–17)
Silver Cup	1.00	20	(14–28)
Ewos	1.10	31	(22–43)
Low protein (23%)	1.00	28	(20–41)
High protein (45%)	1.50	47	(38–58)

[a]Handling and feeding (3% body weight/day) were consistent with all three chemicals and the fish source was the same. The chlordane study was conducted at a different time than that with Antimycin A and carbaryl, and initial fish weights differed (0.4 and 0.2 g, respectively). Conditions of the toxicity tests were as follows: chlordane—temperature 12°C, pH 7.1, hardness 44 mg/L; Antimycin A and carbaryl—temperature 10°C, pH 7.4, hardness 40–44 mg/L.
[b]Weight at time of testing.

generally less than 0.5 order of magnitude, they may be as high as 2.5 orders of magnitude. In view of the widespread use of various formulations for pesticides and some industrial chemicals and the alteration in toxicity of the technical chemicals involved, it is evident that further investigation of the toxicology of those formulations is needed.

Diet

Diet affects body composition, physiological and biochemical functions, and nutritional status of the subject, and thereby influences toxicity (Doull 1980). The 96-h LC50 values for rainbow trout exposed to Antimycin A, carbaryl, and chlordane were affected by the type of diet fed 42 days before exposure (Table 10). Proximate analyses of the various diets were reported by Mehrle et al. (1974). The LC50 for trout fed a high protein diet before exposure to Antimycin A was significantly higher than that for trout fed other diets. The LC50's for all dietary groups differed significantly from each other. No significant effects were observed with carbaryl, although it was the least toxic in the group fed the high protein diet.

Significant differences occurred in chlordane toxicity among rainbow trout fed commercial or synthetic diets (Mehrle et al. 1974; Table 10). Differences existed between the group fed the Ewos diet and the other three commercial diets, and between the group fed Silver Cup and the groups fed Glenco and Oregon Moist diets. The difference in LC50's between the fish fed the Oregon Moist and Glenco diets was not statistically significant. The LC50 for the group receiving the high protein diet differed from the LC50's of all other groups, but the groups fed the low protein and Ewos diets were not significantly different. Differences in size of fish fed the various diets may have contributed to the differences in LC50's.

Feeding of protein-deficient diets has resulted in decreased tolerances to organochlorine, organophosphate, and carbamate pesticides in rats (Boyd and Boulanger 1968; Boyd and DeCastro 1968; Boyd and Tanikella 1969). This response was observed for acute toxicities of Antimycin A, carbaryl, and chlordane to rainbow trout, where a decrease in protein in the diet increased pesticide toxicity. However, protein may be only one of the many dietary factors affecting toxicity. Fish fed fish protein concentrate ranged from most susceptible to intermediate when tested against Antimycin A or carbaryl; fish fed

BEST AVAILABLE COPY

ED_005172C_00000035-00030

Table 11. *Acute toxicity (LC50, µg/L) of pesticides to rainbow trout from various sources and steelhead under static conditions.*[a]

Chemical and duration of test (hours)	Source of rainbow trout							
	Donaldson	Iowa	Missouri	Mt. Whitney	New Hampshire	Soap Lake	Wytheville	Steelhead
Antimycin A								
24	>0.087	0.049	0.10	>0.087	0.085	>0.075	0.10	
96	0.013	<0.008	0.019	0.031	0.009	0.007	0.008	
Carbaryl								
24	6,200	2,800	3,900		4,600	4,600	6,100	
96	<750	<320	1,000		<420	1,500	<320	
Carbofuran								
24							680	1,000
96							380	600
Chlordane								
24			44		78	62		13
96			27		20	13		2.9
d-trans Allethrin								
24							14	19
96							14	18
Malathion								
24			160			39		
96			94			4.1		
Trichlorfon								
24							>12,000	>7,000
96							1,800	1,400

[a]Temperature, 12°C; pH, 7.1–7.5; hardness, 40–40 mg/L.

Ewos (58% protein) were the most sensitive to Antimycin A; and the LC50's among the groups of fish fed the commercial diets were not always directly related to dietary protein concentrations. The primary sources of protein in fish food were fish meal, soybean meal, cottonseed meal, and wheat germ meal in the Glenco, Oregon Moist, and Silver Cup diets; fish protein concentrate in the Ewos diet; and casein in the low and high protein diets. Miranda and Webb (1972) reported that the oral toxicity of heptachlor was less for rats fed a low quality protein (gluten) than for rats fed casein; thus both qualitative and quantitative alterations in dietary proteins may affect acute toxicity.

Dietary constituents other than protein may also affect chronic toxicity of chemicals to fish. Mehrle et al. (1977) reported that a diet high in the amino acid methionine increased the toxicity of DDT to rainbow trout, and Mayer et al. (1978) found that increased levels of dietary vitamin C decreased toxaphene toxicity to channel catfish.

The reason for differences in susceptibility to chemicals due to diet is not completely understood. Differences in commercial diets alone altered the 96-h LC50 by an average of 2.3× (range, 1.1 to 3.8×) or 3.6× (range, 1.7 to 5.7×), if synthetic diets were included. However, the type of diet fed before chemical exposure is important, and the standardization of diets fed to aquatic organisms used in toxicological research would help reduce the variation in toxicity data.

Source of Fish

Rainbow trout from seven sources (several strains) and steelhead were tested with seven chemicals in 25 static tests. The ratios of the highest to lowest LC50's (24 h and 96 h) within a chemical ranged from 1.3 to 23 and averaged 4.8 (Table 11). The highest LC50 was 1.8 to 4.7× that of the lowest LC50 77% of the time, but exceptions occurred with chlordane and malathion. When the effects of steelhead were excluded, the highest LC50 was 1.8 to 4.7× the lowest 88% of the time. The sensitivity of steelhead (anadromous rainbow trout) to chlordane far exceeded that of any other trout. Because the weight of steelhead fry was only 12 to 23% of that

BEST AVAILABLE COPY

ED_005172C_00000035-00031

Table 12. *Acute static 96-h LC50 values (mg/L) of organic chemicals for various life stages of fishes.*

Chemical and species	Life stage: Eyed egg	Yolk-sac fry	Swim-up fry	Advanced fry[a]
Aminocarb				
Rainbow trout	>32	0.036	0.048	0.14
Benomyl				
Rainbow trout		0.28	0.16	0.23
Channel catfish		0.006	0.012	0.029
2,4-D dimethylamine salt				
Fathead minnow	1,450	630	425	320
2,4-D dodecyl/tetra-dodecyl amine salt				
Rainbow trout	47	7.6	1.4	3.2
Dowell M38N				
Cutthroat trout	1.6	0.28		0.46
Fenitrothion				
Rainbow trout	16	4.3	2.4	2.4
Channel catfish		3.8	1.4	4.8
Fire-Trol 100				
Coho salmon		90	920	>1,500
Rainbow trout		150	780	>1,000
Fire-Trol 931				
Coho salmon		580	930	1,000
Rainbow trout		700	790	940
MBC				
Rainbow trout		0.14	0.32	0.65
Channel catfish		0.007	0.012	0.016
Methomyl				
Rainbow trout		3.2	1.3	2.0
Channel catfish		1.8	<0.56	0.76
Methoxychlor				
Brook trout	>50	0.017	0.015	0.019
Mexacarbate				
Brown trout		16		20
Phos-Chek 202				
Coho salmon		145	200	320
Rainbow trout		105	110	230
Phos-Chek 259				
Coho salmon		145	170	250
Rainbow trout		115	94	160
Phosmet				
Rainbow trout	>10	>10	0.28	0.48
Picloram				
Rainbow trout		8.0	8.0	11
Channel catfish		5.8	6.8	16
Pydraul 50E				
Lake trout		2.8	2.9	1.5
Toxaphene				
Channel catfish		0.008	0.001	0.004
Tretolite JW-8226				
Cutthroat trout	0.048	2.9		2.2
Trifluralin				
Rainbow trout		1.6	0.083	0.086
Channel catfish		0.66	0.33	2.2

[a]Weight = 0.2 to 1.2 g.

ED_005172C_00000035-00032

of the other strains tested, size may have accounted for some of the difference in sensitivity, but most probably represented physiological differences. Due to incompleteness of data and the small size of the data base, no rainbow trout from a particular source or of a particular strain could be identified as being the most sensitive. However, differences in LC50's among rainbow trout from different sources within the chemicals tested were similar to the mean interlaboratory differences reported by Lemke (1981) in a round-robin study where different laboratories obtained fish from different sources.

Life Stage and Size

Many investigators have reported the effects of life stage and size of invertebrates and fishes on susceptibility to toxicants. Susceptibility of invertebrates generally decreases with maturity (Sanders and Cope 1968; Sanders 1972). The sensitivity of tadpoles of Fowler's toad to DDT increases as the animals mature (Sanders 1970), whereas that of tadpoles of the southern leopard frog (*Rana sphenocephala*) to toxaphene decreases with age (Hall and Swinford 1980). Similarly, decreases in sensitivity have been observed with increased size in fishes (Anderson and Weber 1975; King 1962; Marking 1966; Mount 1962; Pickering et al. 1962; Surber 1948), although responses varied considerably among the early life stages (Akiyama 1970; Bills and Marking 1976; Dean et al. 1977; Iatomi et al. 1958; Klaverkamp et al. 1977; Mauck et al. 1977; Olson and Marking 1973; Piavis and Howell 1975; Pickering and Vigor 1965; Spotila and Paladino 1979).

Life stage. Analysis of 96-h LC50's for yolk-sac and swim-up fry exposed to 20 chemicals indicated that each was the most sensitive 50% of the time (Table 12). Advanced fry were generally less sensitive than yolk-sac and swim-up fry, but yolk-sac and swim-up fry were less sensitive than advanced fry 3% and 7% of the time, respectively. The ratios of the highest to lowest 96-h LC50 among yolk-sac fry, swim-up fry, and advanced fry averaged 4.5 (range, 1.2 to 36), and were 5.0 or less 83% of the time.

The eyed-egg stage was the least sensitive stage tested, except that it was the most sensitive stage when exposure was to Tretolite JW-8226. Although eggs are generally the least susceptible stage, several exceptions are known. For example, Akiyama (1970) found eggs to be more resistant than fry to two mercurials, but not to a third mercurial. And Mauck et al. (1977) reported that brown trout eggs were significantly more sensitive than yolk-sac fry and fingerlings to mexacarbate. In contrast, TFM was most toxic to green eggs of coho salmon and rainbow trout, followed by swim-up fry and yolk-sac fry; eyed-eggs were the least sensitive (Bills and Marking 1976; Olson and Marking 1973).

Toxicity decreased with increasing maturity in other species that were exposed to different chemicals (shown in parentheses): apple snail, *Pomacea paludosa* (Cutrine-plus); crayfish, *Orconectes nais* (DDT, endrin, malathion, and parathion); and stoneflies, *Pteronarcys californica* (lindane). The 96-h LC50's increased by 1.3× for apple snails (from 1–2 days to 2–4 weeks), 56 to 375× for crayfish (3–5 weeks to maturity), and 4.5× for stonefly naiads (first and second year classes). Although toxicity decreased with age, the change may be small within the stages normally tested. The 96-h LC50's for crayfish exposed to DDT were consistent up to 5 weeks of age (LC50's, μg/L: 1 day, 0.30; 1 week, 0.18; 2 weeks, 0.20; 3 weeks, 0.24; and 5 weeks, 0.90), but increased markedly thereafter (8 weeks, 28; mature, 100).

Fish size. Analysis of covariance was conducted on 27 chemicals used in 51 tests for fish of different weights (Table 13). Toxicity decreased as weight increased, but in only 10 of the equations (six chemicals) were the slopes significantly different from zero. The six regressions with a large range in fish size (antimycin and chlordane) varied in slopes among species, the average being 0.0370 (95% CL, ±0.0213). Ratios of the highest to the lowest LC50 averaged 13 (range, 5.3 to 54) within differences in weight of one to two orders of magnitude.

Most acute toxicity tests are conducted with fish weighing 0.2 to 1.5 g, but of 38 regressions within that size range, only 4 slopes (glyphosate, merphos, toxaphene, and trichlorfon) differed significantly from zero. Ratios of the highest to the lowest LC50 value ranged from 4.9 to 6.2 and averaged 5.6. The slopes for these four chemicals were significantly greater than those for chlordane. However, the four tests with glyphosate, merphos, toxaphene, and trichlorfon represented only 10% of the 38 tests (24 chemicals) in which fish weighing 0.2 to 1.5 g were analyzed; thus size effects would not usually be of great concern in standardized acute toxicity testing, as judged by the range of weights represented here.

BEST AVAILABLE COPY

ED_005172C_00000035-00033

Table 13. *Effect of weight on acute static toxicity (96-h LC50) of organic chemicals to fishes.*[a]

Chemical and species	Weight (g)	LC50's (n)	LC50 range (μg/L)	Intercept (a)	Slope[b] (b)	r^2 (%)
Antimycin A						
Coho salmon	0.9–19	5	0.009–0.06	−1.8883	0.0273	51
Rainbow trout	0.7–107	16	0.009–0.12	−1.7712	0.0167	38
Bluegill	1.2–20	4	0.038–0.20	−1.3390	0.0369	77
Chlordane						
Coho salmon	0.6–19	5	14–80	1.1177	0.0461	87
Rainbow trout	1.2–46	5	10–135	1.3601	0.0187	65
Channel catfish	1.9–18	4	6.7–230	1.1896	0.0764	61
Glyphosate						
Rainbow trout	0.4–1.0	4	1,300–8,300	2.6078	1.4158	77
Merphos						
Bluegill	0.5–0.8	3	1,300–8,000	2.0045	2.3758	75
Toxaphene						
Channel catfish	0.02–0.3	4	0.82–4.2	−0.0429	2.1283	75
Trichlorfon						
Atlantic salmon	0.2–0.6	4	610–2,970	2.4416	1.6326	93

[a]Log 96-h LC50 in μg/L = $a+b$(weight in grams).
[b]Significantly different from zero ($P < 0.05$).

Conclusions

1. No one species, family, or class was the most sensitive to all chemicals all of the time.
2. By rank-order comparison, stoneflies (*Claassenia sabulosa*, *Pteronarcys californica*, *Pteronarcella badia*) were the most sensitive aquatic animals tested, followed by glass shrimp (*Palaemonetes kadiakensis*), amphipods (*Gammarus fasciatus*), daphnids (*Daphnia pulex*, *D. magna*, *Simocephalus serrulatus*), brown trout, rainbow trout, seed shrimp (*Cypridopsis vidua*), largemouth bass, cutthroat trout, bluegills, sow bugs (*Asellus brevicaudus*), coho salmon, yellow perch, channel catfish, common carp, black bullheads, green sunfish, fathead minnows, goldfish, western chorus frog, and Fowler's toad. The ranking was slightly altered by standardizing test media characteristics and organism size.
3. Of the four most commonly tested forms compared together, daphnids (*Daphnia magna* or *D. pulex*) were the most sensitive 58% of the time followed by rainbow trout (35%), bluegills (5%), and fathead minnows (2%). However, the sensitivity of daphnids or rainbow trout either equaled or exceeded that of the other two species, as determined by analysis of variance and multiple mean comparisons.
4. When daphnids and rainbow trout were compared with 4 to 40 other species tested with the same chemicals, (a) the lowest of the two LC50's for daphnids and rainbow trout never exceeded 15× the LC50 of the most sensitive species tested 95% of the time, or never exceeded 25× the LC50 of the most sensitive species 100% of the time; and (b) testing of three forms (*Daphnia*, *Gammarus*, and rainbow trout) provided the lowest toxicity value 88% of the time, and could not be improved more than 2.5% by adding any other single species.
5. Interspecies correlation models could be used to predict acute toxicity values, but the confidence limits were smallest when extrapolation was within families.
6. Of the seven major families tested, the Pteronarcidae were the most sensitive, followed by the Daphnidae, Gammaridae, Salmonidae, Centrarchidae, Ictaluridae, and Cyprinidae. The sensitivities of Daphnidae and Gammaridae may be more realistic here than in the species comparisons, since all three daphnid and amphipod species are represented in the family comparisons. Invertebrates were the most sensitive 95% of the time.
7. The class Insecta was the most sensitive 50% of the time, followed by Crustacea (31%), Osteichthyes (19%), and Amphibia (0%).
8. Frequency distributions of chemical toxicity (EC50's or LC50's) were generally bimodal; insecticides were mainly in the <100 μg/L mode, and herbicides, fungicides, industrials, and other chemicals in the >1,000 μg/L mode.

ED_005172C_00000035-00034

9. Flow-through toxicity values may be estimated by multiplying static values by 0.51, but the error can be large: ratios of static to flow-through tests varied by 1.8 orders of magnitude.

10. Toxicity of aged test solutions increased by a factor of 2 or more only 11% of the time and never increased by a factor of more than 4. Toxicity decreased 22% of the time, and no change occurred in 69% of the tests.

11. Temperature affected the toxicity of 40% of the chemicals tested, generally increasing it. However, the toxicity of a few chemicals (DDT, dimethrin, and methoxychlor) decreased with increasing temperature.

12. The effect of temperature on toxicity conforms to the Q_{10} concept and can be predicted. Temperature increased the toxicity of most chemicals by a factor of 3.1 per 10°C rise in temperature; the factor for organophosphate insecticides was higher (5.1)—possibly due to the simultaneous increase in acetylcholinesterase (AChE) activity and rate of AChE inhibition.

13. Although pH affected the toxicity of only 20% of the chemicals tested, it caused a greater average change in toxicity (16×; range, 4.2 to 45) than did any other factor examined.

14. Regression slopes were consistent among species within a chemical for both pH and temperature, indicating chemical rather than biological differences in toxicity.

15. Hardness had little if any effect on the toxicity of organic chemicals; the negligible effects we observed were probably due to pH differences in solutions of differing hardness.

16. Formulations of pesticides increased the toxicity of the technical material used in preparing them 32% of the time, and decreased it 11% of the time. The effect of formulation on toxicity was within a factor of 0.2 to 5× 85% of the time, but was as high as 318×. Formulations ranked in order of most toxic to least toxic were as follows: oil soluble concentrate, liquid, emulsifiable concentrate, granular, wettable powder, spray concentrate, and oil dispersion. Tests on both formulated and technical pesticides are strongly indicated.

17. Differences in the diet fed to fish before they were tested altered toxicity by as much as 5.7×, but factors of 1.1 to 3.8× would be anticipated within available commercial diets. Standardized diets would help reduce variation in acute toxicity data.

18. Source of rainbow trout contributed to variation in toxicity data; the differences ranged from 1.8 to 4.7× about 88% of the time.

19. Sensitivity of invertebrates and fish decreased with development and increased size. Differences in toxicity among the life stages of fish were 5× or less 83% of the time. The change in toxicity was usually small within the sizes normally tested, and thus size should be of no great concern in standardized acute toxicity testing. However, life stage could be of considerable importance in the natural environment.

20. Of the factors affecting toxicity (except for pesticide formulation), the ratios of the highest to lowest LC50's for a chemical were 5.0 or less 80% or more of the time; this generalization includes pH and temperature, when evaluated on the basis of changes of 1.0 pH unit and 10°C.

21. Generalizations and predictions are accurate for comparative toxicity and factors affecting toxicity 80% of the time. The utility of predictive techniques should be realized, but no single approach correctly predicts acute toxicity under all situations. Factors affecting acute toxicity of a chemical (except pesticide formulations) appear to vary within 5× under standard laboratory conditions. Also, these factors often alter bioavailability of the chemical in the field. Studies involving a range of field conditions, therefore, are strongly indicated to determine the interactions of such factors on both bioavailability and acute toxicity. Species sensitivity varied more than factors affecting toxicity, since toxicity differences between the least to the most sensitive species tested within a chemical averaged two orders of magnitude and were as high as five orders of magnitude. Due to inherent variation and exceptions in acute toxicity data, it is best to test the chemical and species of concern under environmental conditions of interest only when exact information is required.

The acute toxicity test provides a rapid, cost-efficient way of measuring relative toxicity to organisms representing different trophic levels, toxicity in different types of water, and toxicity of different forms and formulations of a chemical. Its main value lies in its use as a relative starting point for determining effects in hazard assessment, along with the expected or measured environmental concentration. Toxicity studies are too often conducted to develop data, without subsequent comprehensive interpretation. We hope that this manual will serve as a guide for future research and bring into focus the pertinent aspects of laboratory studies on acute toxicity, thereby providing a foundation for predictive correlations and hazard assessments.

BEST AVAILABLE COPY

ED_005172C_00000035-00035

References

Akiyama, A. 1970. Acute toxicity of two organic mercury compounds to the teleost, *Oryzias latipes*, in different stages of development. Bull. Jpn. Soc. Sci. Fish. 36:563–570.

Aly, O. M., and M. A. El-Dib. 1971. Hydrolysis of Sevin, Baygon, Pyrolan, and Dimetilan in waters. Water Res. 5:1191–1205.

American Society for Testing and Materials. 1980. Standard practice for conducting acute toxicity tests with fishes, macroinvertebrates, and amphibians. *In* Annual Book of ASTM Standards, E729-80. American Society for Testing and Materials, Philadelphia, Pa. 25 pp.

Anderson, P. D., and L. J. Weber. 1975. Toxic response as a quantitative function of body size. Toxicol. Appl. Pharmacol. 33:471–483.

Bills, T. D., and L. L. Marking. 1976. Toxicity of 3-trifluromethyl-4-nitrophenyl (TFM), 2',5-dichloro-4'-nitrosalicylanilide (Bayer 73), and a 98:2 mixture to fingerlings of seven fish species and to eggs and fry of coho salmon. U.S. Fish Wildl. Serv., Invest. Fish Control 69. 9 pp.

Birge, W. J., and J. A. Black. 1982. Statement on surrogate species clusters concept. Pages A6-5–A6-7 *in* Surrogate Species Workshop Report TR-507-36B. U.S. Environmental Protection Agency, Washington, D.C.

Birge, W. J., J. A. Black, and R. A. Kuehne. 1980. Effects of organic compounds on amphibian reproduction. U.S. Fish Wildl. Serv., Res. Rep. 121. 39 pp.

Black, J. A., W. J. Birge, W. E. McDonnell, A. G. Westerman, B. A. Ramey, and D. M. Bruser. 1982. The aquatic toxicity of organic compounds to embryo-larval stages of fish and amphibians. U.S. Fish. Wildl. Serv., Res. Rep. 133. 61 pp.

Blanck, H. 1982. Aquatic organisms differ in their sensitivity to chemicals. Implications for test strategies. OIKOS Conf. Ecotoxicol. 3:37.

Blanck, H., G. Wallin, and S.-A. Wangberg. 1984. Species-dependent variation in algal sensitivity to chemical compounds. Ecotoxicol. Environ. Saf. 8:339–351.

Boyd, E. M., and M. A. Boulanger. 1968. Augmented susceptibility to carbaryl toxicity in albino rats fed purified casein diets. J. Agric. Food Chem. 16:834–838.

Boyd, E. M., and E. S. DeCastro. 1968. Protein-deficient diet and DDT toxicity. Bull. WHO 38:141–150.

Boyd, E. M., and T. K. Tanikella. 1969. The acute oral toxicity of malathion in relation to dietary protein. Arch. Toxikol. 24:292–303.

Brungs, W. A., and D. I. Mount. 1978. Introduction to a discussion of the use of aquatic toxicity tests for evaluation of the effects of toxic substances. Pages 15–26 *in* J. Cairns, Jr., K. L. Dickson, and A. W. Maki, eds. Estimating the Hazard of Chemical Substances to Aquatic Life. Am. Soc. Test. Mater. STP 657.

Buikema, A. L., Jr., B. R. Niederlehner, and J. Cairns, Jr. 1982. Biological monitoring. Part IV—Toxicity testing. Water Res. 16:239–262.

Cairns, J., Jr., A. G. Heath, and B. C. Parker. 1975a. The effects of temperature upon the toxicity of chemicals to aquatic organisms. Hydrobiolgia 47:135–171.

Cairns, J., Jr., A. G. Heath, and B. C. Parker. 1975b. Temperature influence on chemical toxicity to aquatic organisms. J. Water Pollut. Control Fed. 47:267–280.

Committee on Methods for Toxicity Tests with Aquatic Organisms. 1975. Methods for acute toxicity tests with fish, macroinvertebrates, and amphibians. U.S. Environ. Prot. Agency, Ecol. Res. Ser. EPA-660/3-75-009. 61 pp.

Crandall, A. C., and C. J. Goodnight. 1959. The effect of various factors on the toxicity of sodium pentachlorophenate to fish. Limnol. Oceanogr. 4:53–56.

Dean, H. J., J. R. Colquhoun, and H. A. Simonin. 1977. Toxicity of methoxychlor and naled to several life stages of landlocked Atlantic salmon. N.Y. Fish Game J. 24:144–153.

DeGraeve, G. M., D. L. Geiger, J. S. Meyer, and H. L. Bergman. 1980. Acute and embryo-larval toxicity of phenolic compounds to aquatic biota. Arch. Environ. Contam. Toxicol. 9:557–568.

Doherty, F. G. 1983. Interspecies correlations of acute aquatic median lethal concentration for four standard testing species. Environ. Sci. Technol. 17:661–665.

Doull, J. 1980. Factors influencing toxicology. Pages 70–83 *in* J. Doull, C. D. Klaassen, and M. O. Amdur, eds. Casarett and Doull's Toxicology: The Basic Science of Poisons. Macmillan Publishing Co., New York.

Eisler, R. 1970. Factors affecting pesticide-induced toxicity [illegible] an estuarine fish. U.S. Fish Wildl. Serv., Tech. Pap. [illegible]. 20 pp.

Folmar, L. C., H. O. Sanders, and A. M. Julin. 1979. Toxicity of the herbicide glyphosate and several of its formulations to fish and aquatic invertebrates. Arch. Environ. Contam. Toxicol. 8:269–278.

Gammon, D. W., M. A. Brown, and J. E. Casida. 1981. Two classes of pyrethroid action in the cockroach. Pestic. Biochem. Physiol. 15:181–191.

Giese, A. C. 1968. Cell physiology. W. B. Saunders Co., Philadelphia, Pa. 671 pp.

Goodnight, C. J. 1942. Toxicity of sodium pentachlorophenate and pentachlorophenol to fish. Ind. Eng. Chem. 34:868–872.

Hall, R. J., and D. Swineford. 1980. Toxic effects of endrin and toxaphene on the southern leopard frog (*Rana sphenocephala*). Environ. Pollut. (Series A) 23:53–65.

Hashimoto, Y., and Y. Nishiuchi. 1981. Establishment of bioassay methods for the evaluation of acute toxicity of pesticides to aquatic organisms. Nippon Noyaku Gakkai 6:257–264.

Hogan, J. W. 1970. Water temperature as a source of variation in specific activity of brain acetylcholinesterase of bluegills. Bull. Environ. Contam. Toxicol. 5:347–353.

Holcombe, G. A., J. T. Fiandt, and G. L. Phipps. 1980. Effects of pH increases and sodium chloride additions on the acute toxicity of 2,4-dichlorophenol to the fathead minnow. Water Res. 14:1073–1077.

Iatomi, K., T. Tamura, Y. Itazawa, I. Hanyu, and S. Sugiura. 1958. Toxicity of endrin to fish. Prog. Fish-Cult. 20:155–162.

Inglis, A., and E. L. Davis. 1972. Effects of water hardness on the toxicity of several organic and inorganic herbicides to fish. U.S. Fish. Wildl. Serv., Tech. Pap. 67. 22 pp.

Johnson, D. W. 1968. Pesticides and fishes—a review of selected literature. Trans. Am. Fish. Soc. 97:398–424.

ED_005172C_00000035-00036

Johnson, W. W., and M. T. Finley. 1980. Handbook of acute toxicity of chemicals to fish and aquatic invertebrates. U.S. Fish Wildl. Serv., Resour. Publ. 137. 98 pp.

Kaila, K., and J. Saarikoski. 1977. Toxicity of pentachlorophenol and 2,3,6-trichlorophenol to the crayfish (*Astacus fluviatilis* L.). Environ. Pollut. 12:119–123.

Kenaga, E. E. 1978. Test organisms and methods useful for early assessment of acute toxicity of chemicals. Environ. Sci. Technol. 12:1322–1329.

Kimerle, R. A., A. F. Werner, and W. J. Adams. 1983. Aquatic hazard evaluation principles applied to the development of water quality criteria. Seventh ASTM Symposium on Aquatic Toxicology, Milwaukee, Wis. 15 pp.

King, S. F. 1962. Some effects of DDT on the guppy and the brown trout. U.S. Fish Wildl. Serv., Spec. Sci. Rep.—Fish. 399. 21 pp.

Klaverkamp, J. F., M. Duangsawasdi, W. A. Macdonald, and H. S. Majewski. 1977. An evaluation of fenitrothion toxicity in four life stages of rainbow trout, *Salmo gairdneri*. Pages 231–240 *in* F. L. Mayer and J. L. Hamelink, eds. Aquatic toxicology and hazard evaluation. Am. Soc. Test. Mater. STP 634.

Kobayashi, K., and T. Kishino. 1980. Effect of pH on the toxicity and accumulation of pentachlorophenol in goldfish. Bull. Jpn. Soc. Sci. Fish. 46:167–170.

Konemann, H., and A. Musch. 1981. Quantitative structure–activity relationships in fish toxicity studies. Part II: The influence of pH on the QSAR of chlorophenols. Toxicology 19:223–228.

LeBlanc, G. A. 1984. Interspecies relationships in acute toxicity of chemicals to aquatic organisms. Environ. Toxicol. Chem. 3:47–60.

Lemke, A. E. 1981. Interlaboratory comparison: acute testing set. U.S. Environmental Protection Agency, Duluth, Minn. 29 pp.

Levy, G., and S. P. Gucinski. 1964. Studies on biologic membrane permeation kinetics and acute toxicity of drugs by means of goldfish. J. Pharmacol. Exp. Ther. 146:80–86.

Macek, K., W. Birge, F. L. Mayer, A. L. Buikema, Jr., and A. W. Maki. 1978. Discussion session synopsis—use of aquatic toxicity tests for evaluation of the effects of toxic substances. Pages 27–32 *in* J. Cairns, Jr., K. L. Dickson, and A. W. Maki, eds. Estimating the hazard of chemical substances to aquatic life. Am. Soc. Test. Mater. STP 657.

Macek, K. J., C. Hutchinson, and O. B. Cope. 1969. The effects of temperature on the susceptibility of bluegills and rainbow trout to selected pesticides. Bull. Environ. Contam. Toxicol. 4:174–183.

Macek, K. J., and W. A. McAllister. 1970. Insecticide susceptibility of some common fish family representatives. Trans. Am. Fish. Soc. 99:20–27.

Maki, A. W., K. L. Dickson, and J. Cairns, Jr. 1979. Introduction. Pages 1–6 *in* K. L. Dickson, A. W. Maki, and J. Cairns, Jr., eds. Analyzing the hazard evaluation process. American Fisheries Society, Washington, D.C.

Maren, T. H., R. Embry, and L. E. Broder. 1968. The excretion of drugs across the gill of the dogfish, *Squalus acanthias*. Comp. Biochem. Physiol. 26:853–864.

Marking, L. L. 1966. Evaluation of *p,p'*-DDT as a reference toxicant in bioassays. Invest. Fish Control 10:1–10.

Marking, L. L. 1972. Methods of estimating the half-life of biological activity of toxic chemicals in water. Invest. Fish Control 46:1–9.

Marking, L. L. 1975. Effects of pH on toxicity of antimycin to fish. J. Fish. Res. Board Can. 32:769–773.

Marking, L. L., and V. K. Dawson. 1972. The half-life of biological activity of antimycin determined by fish bioassay. Trans. Am. Fish. Soc. 101:100–105.

Marking, L. L., and L. E. Olson. 1975. Toxicity of the lampricide 3-trifluoromethyl-4-nitrophenol (TFM) to nontarget fish in static tests. Invest. Fish Control 60:1–27.

Mauck, W. L., L. E. Olson, and J. W. Hogan. 1977. Effects of water quality on deactivation and toxicity of mexacarbate (Zectran®) to fish. Arch. Environ. Contam. Toxicol. 6:385–393.

Mauck, W. L., L. E. Olson, and L. L. Marking. 1976. Toxicity of natural pyrethrins and five pyrethroids to fish. Arch. Environ. Contam. Toxicol. 4:18–29.

Mayer, F. L., P. M. Mehrle, and P. L. Crutcher. 1978. Interactions of toxaphene and vitamin C in channel catfish. Trans. Am. Fish. Soc. 107:326–333.

McLeay, D. J., C. C. Walden, and J. R. Munro. 1979. Influence of dilution water on the toxicity of kraft pulp and paper mill effluent, including mechanisms of effect. Water Res. 13:151–158.

Mehrle, P. M., W. W. Johnson, and F. L. Mayer. 1974. Nutritional effects on chlordane toxicity in rainbow trout. Bull. Environ. Contam. Toxicol. 12:513–517.

Mehrle, P. M., F. L. Mayer, and W. W. Johnson. 1977. Diet quality in fish toxicology: effects on acute and chronic toxicity. Pages 269–280 *in* F. L. Mayer and J. L. Hamelink, eds. Aquatic toxicology and hazard evaluation. Am. Soc. Test. Mater. STP 634.

Miranda, C. L., and R. E. Webb. 1972. Effect of quality of dietary protein on heptachlor toxicity. Fed. Proc. Am. Soc. Exp. Biol. 31:726.

Mount, D. I. 1962. Chronic effects of endrin on bluntnose minnows and guppies. U.S. Fish Wildl. Serv., Res. Rep. 58. 38 pp.

Mount, D. 1982. Aquatic surrogates. Pages A6-2–A6-4 *in* Surrogate Species Workshop Report TR-507-36B. U.S. Environmental Protection Agency, Washington, D.C.

O'Brien, R. D. 1967. Insecticides. Action and metabolism. Academic Press, New York. 332 pp.

Olson, L. E., and L. L. Marking. 1973. Toxicity of TFM (lampricide) to six early life stages of rainbow trout (*Salmo gairdneri*). J. Fish. Res. Board Can. 30:1047–1052.

Piavis, G. W., and J. W. Howell. 1975. Effects of 3-trifluoromethyl-4-nitrophenol (TFM) on developmental stages of the sea lamprey. U.S. Fish Wildl. Serv., Invest. Fish Control 64:1–8.

Pickering, Q. H., and C. Henderson. 1964. Acute toxicity of some important petrochemicals to fish. J. Water Pollut. Control Fed. 35:1419–1429.

Pickering, Q. H., C. Henderson, and A. F. Lemke. 1962. The toxicity of organic phosphorus insecticides to different species of warm water fishes. Trans. Am. Fish. Soc. 91:175–184.

Pickering, Q. H., and W. N. Vigor. 1965. The acute toxicity of zinc to eggs and fry of the fathead minnow. Prog. Fish-Cult. 27:153–157.

BEST AVAILABLE COPY

ED_005172C_00000035-00037

Saarikoski, J., and M. Viluksela. 1981. Influence of pH on the toxicity of substituted phenols to fish. Arch. Environ. Contam. Toxicol. 10:747–753.

Sanders, H. O. 1969. Toxicity of pesticides to the crustacean *Gammarus lacustris*. U.S. Fish Wildl. Serv., Tech. Pap. 25. 18 pp.

Sanders, H. O. 1970. Pesticide toxicities to tadpoles of the western chorus frog *Pseudacris triseriata* and Fowler's toad *Bufo woodhousii fowleri*. Copeia 1970: 246–251.

Sanders, H.O. 1970. Toxicities of some herbicides to six species of freshwater crustaceans. J. Water Pollut. Control Fed. 42(Part I):1544–1550.

Sanders, H. O. 1972. Toxicity of some insecticides to four species of malacostracon crustaceans. U.S. Fish Wildl. Serv., Tech. Pap. 66. 19 pp.

Sanders, H. O., and O. B. Cope. 1968. The relative toxicities of several pesticides to naiads of three species of stoneflies. Limnol. Oceanogr. 13:112–117.

Sanders, H. O., M. T. Finley, and J. B. Hunn. 1983. Acute toxicity of six forest insecticides to three aquatic invertebrates and four fishes. U.S. Fish. Wildl. Serv., Tech. Pap. 110. 5 pp.

Schoettger, R. A. 1970. Toxicology of thiodan in several fish and aquatic invertebrates. Invest. Fish Control 35:1–31.

Schoettger, R. A., and A. M. Julin. 1969. Efficacy of quinaldine as an anesthetic for seven species of fish. Invest. Fish Control 22:1–10.

Schoettger, R. A., and W. L. Mauck. 1978. Toxicity of experimental forest insecticides to fish and aquatic invertebrates. Pages 11–27 *in* W. R. Swain and N. K. Ivanikiw, eds. Vol. 2. Proceedings of the Second USA-USSR Symposia on the effects of pollutants upon aquatic ecosystems. U.S. Environ. Prot. Agency, Spec. Publ. EPA-600/3-78-076.

Sills, J. B., and J. L. Allen. 1971. The influence of pH on the efficacy and residues of quinaldine. Trans. Am. Fish. Soc. 100:544–545.

Smith, M. J., and A. G. Heath. 1979. Acute toxicity of copper, chromate, zinc, and cyanide to freshwater fish: effect of different tempertures. Bull. Environ. Contam. Toxicol. 22:113–119.

Snedecor, G. W., and W. G. Cochran. 1967. Statistical methods. Iowa State University Press, Ames. 593 pp.

Snedecor, G. W., and W. G. Cochran. 1980. Statistical methods. Iowa State University Press, Ames. 507 pp.

Spotila, J. R., and F. V. Paladino. 1979. Toxicity of arsenic to developing muskellunge fry (*Esox masquinongy*). Comp. Biochem. Physiol. 62C:67–69.

Sprague, J. B. 1979. Measurement of pollutant toxicity to fish. II. Utilizing and applying bioassay results. Water Res. 4:3–32.

Stecher, K. J. 1983. Susceptibility of rainbow trout (*Salmo gairdneri*) to enteric redmouth disease following exposure to technical PCBs and waste transformer oil containing PCBs. M.S. Thesis. University of Missouri, Columbia. 74 pp.

Sullivan, J. K. 1977. Effects of salinity and temperature on the acute toxicity of cadmium to the estuarine crab *Paragrapsus gaimardii* (Milne Edwards). Aust. J. Mar. Freshwater Res. 28:739–743.

Surber, E. W. 1948. Chemical control agents and their effects on fish. Prog. Fish-Cult. 10:125–131.

Suter, G. W., II, and D. S. Vaughn. 1985. Extrapolation of ecotoxicity data: choosing tests to suite the assessment. Pages 387–399 *in* K. E. Cowser, ed. Synthetic fossil fuel technologies—results of health and environmental studies. Butterworth Publ., Boston, Mass.

Tucker, R. K., and J. S. Leitzke. 1979. Comparative toxicology of insecticides for vertebrate wildlife and fish. J. Pharmacol. Ther. 6:167–220.

U.S. Environmental Protection Agency. 1979. Water quality criteria. Part V: Request for comments. Fed. Regist. 44:15926–15981.

Weber, J. B. 1972. Interaction of organic pesticides with particulate matter in aquatic and soil systems. Pages 55–120 *in* R. F. Gould, ed. Fate of organic pesticides in the aquatic environment. Am. Chem. Soc. Adv. Chem. Ser. 111.

Weiss, C. M., and J. L. Botts. 1957. Factors affecting the responses of fish to toxic materials. Sewage Ind. Wastes 29:810–818.

Woodward, D. F. 1976. Toxicity of the herbicides dinoseb and picloram to cutthroat (*Salmo clarki*) and lake trout (*Salvelinus namaycush*). J. Fish. Res. Board Can 33:1671–1676.

Woodward, D. F., and W. L. Mauck. 1980. Toxicity o five forest insecticides to cutthroat trout and two specie of aquatic invertebrates. Bull. Environ. Contam. Toxico 25:846–853.

Wuhrmann, K. 1952. Concerning some principles of th toxicology of fish. Bull. Cent. Belge Etud. Doc. Eaux 1 49 pp.

Zanella, E. 1983. Effect of pH on acute toxicity of deh droabietic acid and chlorinated dehydroabietic acid fish and *Daphnia*. Bull. Environ. Contam. Toxic 30:133–140.

BEST AVAILABLE COPY

ED_005172C_00000035-00038

Appendix

		Page
Table A1.	Scientific and common names of invertebrates used for acute toxicity testing	34
Table A2.	Scientific and common names of vertebrates used for acute toxicity testing	35
Table A3.	Species list with chemicals tested	36
Table A4.	Common and alternate names of chemicals used for aquatic toxicity testing	55
Table A5.	Acute toxicity data for 410 chemicals and 66 species of freshwater animals from the Columbia National Fisheries Research Laboratory, U.S. Fish and Wildlife Service, 1965–1984	64

BEST AVAILABLE COPY

ED_005172C_00000035-00039

Table A1. *Scientific and common names of invertebrates used for acute toxicity testing.*

Class, order, family, genus, and species	Common name
Crustacea	
Anostraca	Fairy shrimps
Streptocephalidae	
Streptocephalus seali	
Cladocera	Waterfleas, daphnids
Daphnidae	
Daphnia magna	
Daphnia pulex	
Simocephalus serrulatus	
Podocopa	Seed shrimps, ostracods
Cypridae	
Cypridopsis vidua	
Isopoda	Aquatic sow bugs
Asellidae	
Asellus brevicaudus	
Amphipoda	Scuds, sideswimmers
Gammaridae	
Gammarus fasciatus	
Gammarus lacustris	
Gammarus pseudolimnaeus	
Decapoda	Crayfishes, shrimps
Astacidae	Crayfishes
Orconectes nais	
Procambarus sp.	
Palaemonidae	Shrimps
Palaemonetes kadiakensis	
Insecta	
Plecoptera	Stoneflies
Perlidae	
Acroneuria sp.	
Claassenia sabulosa	
Perlodidae	
Isogenus sp.	
Isoperla sp.	
Skwala sp.	
Pteronarcidae	
Pteronarcella badia	
Pteronarcys californica	
Ephemeroptera	Mayflies
Baetidae	
Baetis sp.	
Ephemerella sp.	
Ephemeridae	
Hexagenia bilineata	
Odonata	Dragonflies, damselflie
Coenagrionidae	Damselflies
Ischnura verticalis	
Lestes congener	
Gomphidae	Dragonflies
Phiogomphus sp.	
Libellulidae	Dragonflies
Macromia sp.	
Trichoptera	Caddisflies
Hydropsychidae	
Hydropsyche sp.	
Limnephilidae	
Limnephilus sp.	
Diptera	True flies
Culicidae	Phantom midges
Chaoborus sp.	
Rhagionidae	Snipe flies
Atherix variegata	
Tendipedidae	Midges
Chironomous plumosus	
Pentaneura sp.	
Tipulidae	True crane flies
Tipula sp.	
Gastropoda	
Ctenobranchiata	
Ampullariidae	
Pomacea paludosa	Apple snail

BEST AVAILABLE COPY

ED_005172C_00000035-00040

Table A2. *Scientific and common names of vertebrates used for acute toxicity testing.*

Class, order, family, genus, and species	Common name
Osteichthyes	
Acipenseriformes	
Polyodontidae	
Polyodon spathula	Paddlefish
Salmoniformes	
Salmonidae	
Oncorhynchus kisutch	Coho salmon
Oncorhynchus tshawytscha	Chinook salmon
Salmo clarki	Cutthroat trout
Salmo gairdneri	Rainbow trout
Salmo salar	Atlantic salmon
Salmo trutta	Brown trout
Salvelinus fontinalis	Brook trout
Salvelinus namaycush	Lake trout
Esocidae	
Esox lucius	Northern pike
Cypriniformes	
Cyprinidae	
Carassius auratus	Goldfish
Cyprinus carpio	Common carp
Notropis blennius	River shiner
Pimephales promelas	Fathead minnow
Catostomidae	
Catostomus catostomus	Longnose sucker
Catostomus commersoni	White sucker
Siluriformes	
Ictaluridae	
Ictalurus melas	Black bullhead
Ictalurus punctatus	Channel catfish
Clariidae	
Clarias batrachus	Walking catfish
Atheriniformes	
Poeciliidae	
Gambusia affinis	Mosquitofish
Perciformes	
Centrarchidae	
Lepomis cyanellus	Green sunfish
Lepomis macrochirus	Bluegill
Lepomis microlophus	Redear sunfish
Micropterus dolomieui	Smallmouth bass
Micropterus salmoides	Largemouth bass
Pomoxis annularis	White crappie
Pomoxis nigromaculatus	Black crappie
Percidae	
Perca flavescens	Yellow perch
Stizostedion vitreum vitreum	Walleye
[illegible]idae	
[illegible]*pia mossambica*	Tilapia
[illegible]ia	
Anura	
Bufonidae	
Bufo woodhousei fowleri	Fowler's toad
Hylidae	
Pseudacris triseriata	Western chorus frog

BEST AVAILABLE COPY

ED_005172C_00000035-00041

Table A3. *Species list with chemicals tested.*

STREPTOCEPHALUS SEALI — 1 TESTS

1	PHOSMET

DAPHNIA MAGNA — 110 TESTS

2	ALACHLOR	2	METHOMYL
1	AMORO	2	METHYL PARATHION
3	AMINOCARB	2	METOLACHLOR
1	ANTIMYCIN A	1	METRIBUZIN
1	AZINPHOS-ETHYL	1	MIREX
1	BARBAN	1	MON 0818
1	BENOMYL	1	NITRALIN
1	BENZOTHIAZOLE 2-METHYL MERCAPTO	2	OXAMYL
1	CARBARYL	1	PARATHION
1	CCA TYPE III	2	PENTACHLOROPHENOL
1	CHLORDECONE	1	PENTACHLOROPHENOL (DOWICIDE EC-7)
1	CHLORFENETHOL	2	PENTACHLOROPHENOL COPPER SALT
1	CLONITRALIDE	1	PERMETHRIN
1	CYHEXATIN	1	PHOSFLEX 31P
1	CYTROL AMITROLE-T	2	PHOSMET
4	DCPA	1	PICLORAM
2	DDD	2	PROFENOFOS
1	DDT	2	PROPACHLOR
1	DEF	1	PYDRAUL 50E
1	DICAMBA	1	SANTICIZER 148
1	DICHLOROPROPENE	1	SANTICIZER 154
2	DIFLUBENZURON	2	SIMAZINE
1	DIOXATHION	2	SODIUM SELENITE
1	DIPHENAMIDE	1	SRCII SYNFUEL
3	ENDRIN	1	TERBUFOS
1	ETHION	1	THANITE
3	FENITROTHION	1	TOXAPHENE
1	FENVALERATE	2	TRIALLATE
2	FLUCHLORALIN	1	TRIFLURALIN
2	FLUOMETURON	1	WATER GLYCOLS 894-44A
1	FLUORENE	1	WATER GLYCOLS 894-44B
4	FLURIDONE	1	WATER GLYCOLS 894-44C
1	FOSAMINE AMMONIUM	1	XYLENOL DIMETHYLAMINO
1	FYRQUEL GT	1	2,3,4,6-TETRACHLOROPHENOL
1	GLYPHOSATE	3	2,4-D BUTOXYETHANOL ESTER
1	HOUGHTO-SAFE 520	2	2,4-D DIMETHYLAMINE SALT
1	KRONITEX 200	1	2,4-D PROPYLENE GLYCOL BUTYL ETHER ES
2	LINURON	1	6-CHLORO-2-PICOLINIC ACID
1	MALATHION		

DAPHNIA PULEX — 45 TESTS

1	ALDRIN	1	ETHION
1	ALLETHRIN	1	FENAC
1	ARAMITE	1	FENTHION
1	AZIDE POTASSIUM	1	HEPTACHLOR
1	AZIDE SODIUM	1	LIME SULFUR
1	AZINPHOS-ETHYL	1	LINDANE
1	BENZENE HEXACHLORIDE	1	MALATHION
1	CARBARYL	1	METHOXYCHLOR
1	CHLORDANE	1	MEVINPHOS
1	CHLOROBENZILATE	1	MEXACARBATE
1	CRYOLITE	1	MIREX
1	DALAPON	1	NALED
1	DDD	1	PARAQUAT
1	DDT	1	PARATHION
1	DEAD-X	1	PHOSPHAMIDON
1	DEMETON	1	PROPHAM
1	DIAZINON	1	ROTENONE
1	DICHLOBENIL	1	SILVEX PROPYLENE GLYCOL BUTYL ETHER E
1	DICHLORVOS	1	SODIUM ARSENITE
1	DIELDRIN	1	TOXAPHENE
1	DIURON	1	TRICHLORFON
1	DNOC	1	TRIFLURALIN
1	ENDRIN		

BEST AVAILABLE COPY

ED_005172C_00000035-00042

Table A3. *Continued.*

SIMOCEPHALUS SERRULATUS — 67 TESTS

2	ALDRIN	1	FENAC
1	ALLETHRIN	1	FENTHION
1	ARAMITE	2	HEPTACHLOR
1	AZIDE POTASSIUM	1	LIME SULFUR
1	AZIDE SODIUM	2	LINDANE
1	AZINPHOS-ETHYL	3	MALATHION
3	CARBARYL	2	METHOXYCHLOR
2	CHLORDANE	1	METHYL PARATHION
1	CHLOROBENZILATE	2	MEVINPHOS
1	COUMAPHOS	1	HEXACARBATE
1	CRYOLITE	1	MIREX
1	DALAPON	2	NALED
2	DDD	1	PARAQUAT
2	DDT	2	PARATHION
1	DEAD-X	2	PHOSPHAMIDON
2	DIAZINON	1	PROPHAM
1	DICHLOBENIL	1	ROTENONE
2	DICHLORVOS	1	SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER
1	DICROTOPHOS	1	SODIUM ARSENITE
2	DIELDRIN	2	TOXAPHENE
1	DIURON	2	TRICHLORFON
2	ENDRIN	1	TRIFLURALIN
1	ETHION	1	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER

CYPRIDOPSIS VIDUA — 22 TESTS

1	ALDRIN	1	METAM-SODIUM
1	CARBARYL	1	METHOXYCHLOR
1	CLONITRALIDE	1	MOLINATE
1	DDD	1	SILVEX BUTOXYETHANOL ESTER
1	DDT	2	SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER
1	DICHLONE	1	SIMAZINE
1	DIPHENAMIDE	1	VERNOLATE
1	ENDRIN	1	2,4-D BUTOXYETHANOL ESTER
1	FENTHION	1	2,4-D DIMETHYLAMINE SALT
1	LINDANE	1	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER
1	MALATHION		

ASELLUS BREVICAUDUS — 39 TESTS

1	ANTIMYCIN A	1	METHOXYCHLOR
1	AZINPHOS-METHYL	1	MEVINPHOS
1	BINAPACRYL	1	HEXACARBATE
1	CARBARYL	1	NALED
1	CYTROL AMITROLE-T	1	ORYZALIN
1	DDD	1	OXYDEMETON-METHYL
1	DDT	2	PARATHION
1	DICAMBA	2	PHOSMET
1	DICHLOBENIL	1	PIPERONYL BUTOXIDE
1	DICHLONE	1	SILVEX
1	DIELDRIN	2	SILVEX BUTOXYETHANOL ESTER
1	DIPHENAMIDE	1	SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER
1	DIURON	1	TRIFLURALIN
1	ENDRIN	1	VERNOLATE
1	EPTC	1	2,4-D BUTOXYETHANOL ESTER
1	FENTHION	1	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER
1	LINDANE	1	2,4,5-T BUTOXYETHANOL ESTER
1	MALATHION	1	6-CHLORO-2-PICOLINIC ACID

GAMMARUS FASCIATUS — 149 TESTS

2	ALDRIN	1	FENAMINOSULF
1	ALLETHRIN	1	FENITROTHION
1	AMINOCARB	1	FENSULFOTHION
2	AMITROLE	1	FENTHION
1	ANILAZINE	1	FOLPET
1	ANTIMYCIN A	1	GLYODIN
1	ARAMITE	2	HEPTACHLOR

BEST AVAILABLE COPY

ED_005172C_00000035-00043

Table A3. *Continued.*

(CONT.) GAMMARUS FASCIATUS — 149 TESTS

1	AZIDE POTASSIUM	2	LETHANE 384
1	AZIDE SODIUM	2	LINDANE
2	AZINPHOS-METHYL	3	MALATHION
1	BENEFIN	1	METHOXYCHLOR
1	BENSULIDE	2	METHYL PARATHION
1	BENZYL BENZOATE	1	METHYL TRITHION
2	BUFENCARB	3	MEVINPHOS
1	BUTOXY POLYPROPYLENE GLYCOL	1	MEXACARBATE
2	BUTYLATE	1	MOLINATE
2	CACODYLIC ACID	1	MONOCROTOPHOS
1	CARBARYL	1	MSMA
1	CARBOPHENOTHION	1	NALED
1	CHLORDANE	1	NITRALIN
1	CHLORFENVINPHOS	1	NITROFEN
1	CLONITRALIDE	1	NOREA
2	CRUFOMATE	1	ORYZALIN
1	CYANAZINE	2	OXYDEMETON-METHYL
1	CYHEXATIN	1	PARAQUAT
2	DDD	3	PARATHION
3	DDT	1	PCB AROCLOR 1248
1	DEAD-X	1	PCB AROCLOR 1254
1	DEET	1	PEBULATE
1	DEF	2	PHORATE
2	DEMETON	2	PHOSMET
1	DIAZINON	3	PHOSPHAMIDON
2	DICAMBA	1	PICLORAM
2	DICHLOFENTHION	1	PROPANIL
1	DICHLONE	1	PROPHAM
1	DICROTOPHOS	2	PYRETHRUM
2	DIELDRIN	2	RONNEL
1	DINOCAP	1	ROTENONE
2	DIOXATHION	1	SILVEX
1	DIPHENAMIDE	1	SIMAZINE
1	DIQUAT	1	TEPP
2	DISULFOTON	1	TETRADIFON
1	DIURON	1	TETRAMINE
1	DNOC	2	THANITE
1	DODINE	2	TOXAPHENE
1	ENDOSULFAN	1	TRIFLURALIN
1	ENDOTHALL HYDROTHOL 47	1	TRIPHENYLTIN HYDROXIDE
1	ENDOTHALL POTASSIUM SALT	2	VERNOLATE
2	ENDRIN	2	2,4-D BUTOXYETHANOL ESTER
1	EPN	1	2,4-D DIMETHYLAMINE SALT
1	EPTC	1	2,4-D ISOOCTYL ESTER
1	ETHION	3	2,4-D PROPYLENE GLYCOL BUTYL ETHER ES
1	ETHYLENE DICHLORIDE	2	2,4,5-T BUTOXYETHANOL ESTER
1	FENAC	1	6-CHLORO-2-PICOLINIC ACID

GAMMARUS LACUSTRIS — 28 TESTS

1	BENZENE HEXACHLORIDE	1	DIMETHOATE
1	CAPTAFOL	1	ENDOSULFAN
1	CARBARYL	1	ENDOTHALL HYDROTHOL 191
1	CARBOPHENOTHION	1	ENDRIN
1	CHLORONITROPROPANE	1	FENTHION
1	CHLORPYRIFOS	1	HEPTACHLOR
1	COUMAPHOS	1	LINDANE
1	CROTOXYPHOS	1	METHOXYCHLOR
1	DDT	1	METHYL DEMETON
1	DICHLOBENIL	1	MEXACARBATE
1	DICHLOFENTHION	1	PARATHION
1	DICHLORVOS	1	PROPOXUR
1	DICROTOPHOS	1	TEMEPHOS
1	DILAN	1	TRICHLORFON

GAMMARUS PSEUDOLIMNAEUS — 90 TESTS

2	ACEPHATE	1	MIREX
9	AMINOCARB	1	PCB AROCLOR 1242

ED_005172C_00000035-00044

Table A3. *Continued.*

(CONT.) GAMMARUS PSEUDOLIMNAEUS — 90 TESTS

1	BENOMYL	1	PCB 2,2',4,4',6,6'-HEXACHLOROBIPHENYL
1	BENZOTHIAZOLE 2-METHYL MERCAPTO	1	PCB 2,2',4,5,5'-PENTACHLOROBIPHENYL
4	CARBARYL	1	PCB 2,3,4'-TRICHLOROBIPHENYL
1	CCA TYPE III	1	PCB 2,4-DICHLOROBIPHENYL
1	CHLORDECONE	1	PCB 4,4'-DICHLOROBIPHENYL
1	CLONITRALIDE	1	PERMETHRIN
2	DCPA	1	PHOS-CHEK 202
1	DEF	1	PHOS-CHEK 259
2	DIFLUBENZURON	1	PHOSFLEX 31P
4	FENITROTHION	1	PHTHALATE DI-2-ETHYLHEXYL
1	FENVALERATE	1	PHTHALATE DIBUTYL
1	FIRE-TROL 100	1	PICLORAM
1	FIRE-TROL 931	2	PROFENOFOS
1	FLUCHLORALIN	2	PYDRAUL 50E
4	FLURIDONE	1	SANTICIZER 148
1	FOSAMINE AMMONIUM	1	SANTICIZER 154
1	FYRQUEL GT	1	TERBUFOS
1	GLYPHOSATE	2	TFM
1	HOUGHTO-SAFE 1120	9	TRICHLORFON
1	HOUGHTO-SAFE 520	1	WATER GLYCOLS 894-44A
1	KRONITEX 200	1	WATER GLYCOLS 894-44B
7	METHOMYL	1	WATER GLYCOLS 894-44C
5	METHOXYCHLOR		

ORCONECTES NAIS — 30 TESTS

1	BENOMYL	1	HEPTACHLOR
1	CHLORDANE	2	MALATHION
1	CLONITRALIDE	1	METHOXYCHLOR
7	DDT	1	METHYL PARATHION
1	DEF	2	PARATHION
1	DICROTOPHOS	1	PCB AROCLOR 1242
1	DIELDRIN	1	PCB AROCLOR 1254
2	ENDRIN	1	PHTHALATE DIBUTYL
2	FENTHION	1	PYDRAUL 50E
1	GLYPHOSATE	1	TFM

PROCAMBARUS SP. — 13 TESTS

1	AZINPHOS-METHYL	1	PARATHION
1	BENOMYL	1	PCB AROCLOR 1254
1	CARBARYL	1	PHOXIM
1	LEPTOPHOS	2	TRICHLORFON
4	MEXACARBATE		

PALAEMONETES KADIAKENSIS — 44 TESTS

1	ALDRIN	1	FENTHION
1	ANTIMYCIN A	1	HEPTACHLOR
1	AZINPHOS-METHYL	3	MALATHION
1	BUTENCARB	1	METHOXYCHLOR
1	CACODYLIC ACID	1	MEVINPHOS
1	CARBARYL	2	MEXACARBATE
1	CARBOPHENOTHION	1	MOLINATE
1	CLONITRALIDE	1	NALED
2	DDD	2	PARATHION
2	DDT	1	PCB AROCLOR 1254
1	DICAMBA	1	SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER
1	DIPHENAMIDE	1	SIMAZINE
1	DISULFOTON	1	TRIFLURALIN
1	ENDOTHALL HYDROTHOL 191	1	VERNOLATE
2	ENDRIN	2	2,4-D BUTOXYETHANOL ESTER
1	EPN	1	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER
1	ETHANOL	1	2,4,5-T BUTOXYETHANOL ESTER
1	ETHION	1	6-CHLORO-2-PICOLINIC ACID

ACRONEURIA SP. — 1 TESTS

1	ENDRIN

BEST AVAILABLE COPY

ED_005172C_00000035-00045

Table A3. *Continued.*

	CLAASSENIA SABULOSA		10 TESTS
1	CARBARYL	1	HEPTACHLOR
1	CHLORPYRIFOS	1	MALATHION
1	DDT	1	PARATHION
1	DIELDRIN	1	TOXAPHENE
1	ENDRIN	1	TRICHLORFON
	ISOGENUS SP.		14 TESTS
2	ACEPHATE	2	METHOMYL
6	CARBARYL	2	TRICHLORFON
2	DIFLUBENZURON		
	ISOPERLA SP.		2 TESTS
1	DDT	1	MALATHION
	SKWALA SP.		4 TESTS
2	AMINOCARB	2	FENITROTHION
	PTERONARCELLA BADIA		41 TESTS
3	ACEPHATE	2	METHOMYL
6	AMINOCARB	2	METHOXYCHLOR
4	CARBARYL	1	PARATHION
1	DDT	1	PCB AROCLOR 1016
1	DIELDRIN	1	TEMEPHOS
1	ENDRIN	1	TOXAPHENE
3	FENITROTHION	1	TRETOLITE J-146
1	FENTHION	4	TRICHLORFON
2	GREEN CRUDE OIL	1	2,4-D BUTYL ESTER
1	HEPTACHLOR	1	2,4-D PROPYLENE GLYCOL BUTYL ETHER E
3	MALATHION		
	PTERONARCYS CALIFORNICA		77 TESTS
1	ALDRIN	1	ETHION
1	ALLETHRIN	1	ETHYLENE DICHLORIDE
1	ARAMITE	2	FENAC
1	AZIDE POTASSIUM	1	FENAMINOSULF
1	AZIDE SODIUM	1	FENITROTHION
1	AZINPHOS-ETHYL	1	FENTHION
1	AZINPHOS-METHYL	1	HEPTACHLOR
1	BACILLUS THURINGIENSIS	2	LINDANE
1	BENZENE HEXACHLORIDE	1	MALATHION
1	CAPTAFOL	1	METHIOCARB
1	CARBARYL	1	METHOXYCHLOR
1	CHLORDANE	1	METHYL DEMETON
1	CHLORFENVINPHOS	1	METHYL TRITHION
1	CHLORPYRIFOS	1	MEVINPHOS
1	CLONITRALIDE	1	HEXACARBATE
1	CROTOXYPHOS	3	MOLINATE
1	DALAPON	1	NALED
1	DDD	1	OVEX
1	DDT	1	PARAQUAT
1	DEAD-X	1	PARATHION
1	DEF	1	PHORATE
1	DIAZINON	1	PHOSPHAMIDON
1	DICHLOBENIL	1	PICLORAM
1	DICHLOFENTHION	1	PROPOXUR
1	DICHLORVOS	1	SILVEX PROPYLENE GLYCOL BUTYL ETHER
1	DICOFOL	1	SIMAZINE
1	DICROTOPHOS	1	SODIUM ARSENITE
1	DIELDRIN	1	TEMEPHOS
1	DIMETHOATE	1	TERPENE POLYCHLORINATES
1	DISULFOTON	1	TOXAPHENE
1	DIURON	1	TRICHLORFON

ED_005172C_00000035-00046

Table A3. *Continued.*

(CONT.) PTERONARCYS CALIFORNICA — 77 TESTS

1	DNOC	1	TRICHLORONATE
1	ENDOSULFAN	1	TRIFLURALIN
1	ENDOTHALL HYDROTHOL 191	1	2,4-D BUTYL ESTER
1	ENDRIN	2	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER
1	EPN	1	2,4-DB

BAETIS SP. — 1 TESTS

1	ENDRIN

EPHEMERELLA SP. — 1 TESTS

1	DDT

HEXAGENIA BILINEATA — 3 TESTS

1	DDT	1	PARATHION
1	ENDRIN		

ISCHNURA VERTICALIS — 9 TESTS

1	DDD	1	PARATHION
1	DIELDRIN	1	PCB AROCLOR 1242
2	ENDRIN	1	PCB AROCLOR 1254
1	METHYL PARATHION	1	TFM

LESTES CONGENER — 1 TESTS

1	MALATHION

OPHIOGOMPHUS SP. — 1 TESTS

1	DDT

MACROMIA SP. — 2 TESTS

1	PCB AROCLOR 1242	1	PCB AROCLOR 1254

HYDROPSYCHE SP. — 1 TESTS

1	MALATHION

LIMNEPHILUS SP. — 1 TESTS

1	MALATHION

CHAOBORUS SP. — 1 TESTS

1	DDT

ATHERIX VARIEGATA — 4 TESTS

1	DDT	1	MALATHION
1	ENDRIN	1	TOXAPHENE

CHIRONOMUS PLUMOSUS — 95 TESTS

6	ACEPHATE	2	OXAMYL
2	ALACHLOR	2	PENTACHLOROPHENOL COPPER SALT
1	AMDRO	1	PERMETHRIN
3	AMINOCARB	1	PHOSFLEX 31P
1	BARBAN	8	PHOSMET
1	BENOMYL	1	PHTHALATE DI-2-ETHYLHEXYL
1	BENZOTHIAZOLE 2-METHYL MERCAPTO	2	PHTHALATE DIBUTYL
1	CHLORDECONE	1	PHTHALATE 2-ETHYLHEXYL
1	CLONITRALIDE	1	PHTHALIC ACID
2	DCPA	2	PROFENDFOS

BEST AVAILABLE COPY

ED_005172C_00000035-00047

Table A3. *Continued.*

(CONT.) CHIRONOMUS PLUMOSUS — 95 TESTS

1	DEF	2	PROPACHLOR	7
1	DIFLUBENZURON	1	PYDRAUL 50E	9
3	FENITROTHION	1	SANTICIZER 148	7
1	FENVALERATE	1	SANTICIZER 154	2
2	FLUCHLORALIN	2	SODIUM SELENITE	5
2	FLUOMETURON	1	SRCII SYNFUEL	1
1	FLUORENE	1	TERBUFOS	10
4	FLURIDONE	6	TOXAPHENE	2
1	FOSAMINE AMMONIUM	1	TRIALLATE	4
1	FYRQUEL GT	1	WATER GLYCOLS 894-44A	1
3	GLYPHOSATE	1	WATER GLYCOLS 894-44B	1
1	HOUGHTO-SAFE 520	1	WATER GLYCOLS 894-44C	1
1	KRONITEX 200	1	XYLENOL DIMETHYLAMINO	2
2	LINURON	1	2-ETHYLHEXANOL	1
2	METHOMYL	2	2,4-D BUTOXYETHANOL ESTER	2
2	METOLACHLOR	1	2,4-D DIMETHYLAMINE SALT	2
1	MIREX	1	2,4-D DODECYL/TETRADODECYL AMINE SALT	5
1	MON 0818			2
				1

PENTANEURA SP. — 1 TESTS

1	DDT		7
			14
			1
			15

TIPULA SP. — 3 TESTS

1	DDT	1	TOXAPHENE		1
1	ENDRIN				1
					6

APPLE SNAIL — 6 TESTS

1	AQUA-VATOR	1	DIQUAT		1
3	CUTRINE-PLUS	1	KOMEEN		1
					2

PADDLEFISH — 1 TESTS

1	ANTIMYCIN A				15
					2
					9
					5

COHO SALMON — 105 TESTS

10	ANTIMYCIN A	2	LINDANE		1
3	AZINPHOS-METHYL	2	MALATHION		8
2	CAPTAN	1	METHYL PARATHION		
5	CARBARYL	15	MEXACARBATE	9	
1	CARBOFURAN	4	PHOS-CHEK 202	1	
5	CHLORDANE	4	PHOS-CHEK 259	1	
2	D-TRANS ALLETHRIN	9	PHOXIM	2	
2	DDT	1	PHTHALATE DI-2-ETHYLHEXYL	2	
3	DINITRAMINE	1	POTASSIUM NAPHTHALENEACETATE	6	
1	DIURON	1	PYDRAUL 50E	1	
1	ENDOTHALL AQUATHOL K	2	PYRETHRUM	1	
1	ENDRIN	2	RESMETHRIN	1	
1	FENAMINOSULF	2	RU-11679	26	
1	FENITROTHION	1	SD 16898	1	
1	FENTHION	1	SD 17250	20	
4	FIRE-TROL 100	1	SILVEX	1	
4	FIRE-TROL 931	4	TEMEPHOS	1	
1	FOLPET	1	TOXAPHENE	1	
1	HEXACHLOROBENZENE	1	TRICHLORFON	1	
1	HOUGHTO-SAFE 1120			1	
				1	

CHINOOK SALMON — 35 TESTS

1	ALDRIN	2	PHOSMET	1	
1	CAPTAN	1	PURIFLOC C-3	5	
1	CARBARYL	1	PYDRAUL 50E	1	
1	HEXACHLOROBENZENE	1	PYRETHRUM	14	
8	MEXACARBATE	3	UREABOR	1	
3	PENTACHLOROPHENOL	3	2,4-D DIMETHYLAMINE SALT	2	
6	PENTACHLOROPHENOL SODIUM SALT	3	2,4-D DODECYL/TETRADODECYL AMINE SALT	1	

BEST AVAILABLE COPY

ED_005172C_00000035-00048

Table A3. *Continued.*

CUTTHROAT TROUT — 305 TESTS

7	ACEPHATE
9	AMINOCARB
7	ANTIMYCIN A
2	BENZENE HEXACHLORIDE
5	BLACK CRUDE OIL
1	CAPTAN
10	CARBARYL
2	CHLORDANE
4	CHLORPYRIFOS
1	COUMAPHOS
1	CROTOXYPHOS
1	D-D SOIL FUMIGANT
2	DDT
1	DEMETON
2	DIAZINON
2	DICHLOFENTHION
5	DICHLORVOS
2	DICOFOL
1	DIELDRIN
7	DIFLUBENZURON
14	DINOSEB
1	DIOXATHION
15	DIURON
1	DOWELL A170
1	DOWELL F75A
1	DOWELL F75N
1	DOWELL L47
6	DOWELL M38N
1	DOWELL W35
1	DOWICIL 75
1	ENDOTHALL HYDROTHOL 191
1	ENDRIN
2	EPN
15	EPTC
2	ETHION
9	FENITROTHION
5	FENTHION
1	FLIT MLO
8	GREEN CRUDE OIL
3	LEPTOPHOS
2	LETHANE 384
5	MALATHION
1	METHOMYL
5	METHOXYCHLOR
2	METHYL PARATHION
1	METHYL TRITHION
3	MEXACARBATE
1	NALED
2	PARATHION
1	PCB AROCLOR 1221
1	PCB AROCLOR 1232
1	PCB AROCLOR 1242
1	PCB AROCLOR 1248
1	PCB AROCLOR 1254
1	PCB AROCLOR 1260
1	PCB AROCLOR 1262
1	PCB AROCLOR 1268
1	PCB AROCLOR 4465
1	PCB AROCLOR 5442
1	PCB AROCLOR 5460
1	PENTACHLOROPHENOL
1	PENTACHLOROPHENOL (DOWICIDE EC-7)
1	PENTACHLOROPHENOL SODIUM SALT
2	PHORATE
1	PHOTO-DIELDRIN
22	PICLORAM
2	RONNEL
2	TEMEPHOS
2	THANITE
1	TRETOLITE J-146
1	TRETOLITE JN-9045
7	TRETOLITE JW-8226
17	TRICHLORFON
6	ZINC SULFATE
12	2,4-D
17	2,4-D BUTYL ESTER
16	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER

RAINBOW TROUT — 917 TESTS

9	ACEPHATE
1	ACETONE
1	AKTON
2	ALACHLOR
2	ALDICARB
6	ALDRIN
1	ALLETHRIN
1	AMDRO
1	AMETRYN
26	AMINOCARB
1	ANILAZINE
20	ANTIMYCIN A
1	APHOLATE
1	AQUA COP
1	AQUA-VATOR
1	ARAMITE
1	ATRAZINE 4L
1	AZIDE POTASSIUM
1	AZIDE SODIUM
1	AZINPHOS-ETHYL
5	AZINPHOS-METHYL
1	BACILLUS THURINGIENSIS
14	BENOMYL
1	BENSULIDE
2	BENTHIOCARB
1	BENZALKONIUM CHLORIDE
1	BENZENE
4	CHLORPYRIFOS
2	CHLORPYRIFOS-METHYL
1	CLARIFITE
1	CLONITRALIDE
1	COPPER SULFATE
1	COPPER-COUNT-N
1	COPPER-COUNT-NS
2	CORREX
1	COUMAPHOS
1	CROTOXYPHOS
1	CRYOLITE
1	CUMYLPHENYL DIPHENYL PHOSPHATE
1	CYANAZINE
1	CYANO(METHYLMERCURI)GUANIDINE
2	D-D SOIL FUMIGANT
1	D-TRANS ALLETHRIN
2	DDD
1	DDE
7	DDT
1	DEAD-X
11	DEF
2	DEMETON
1	DIAZINON
1	DICAMBA
2	DICHLOBENIL
1	DICHLOFENTHION
1	DICHLONE

BEST AVAILABLE COPY

ED_005172C_00000035-00049

Table A3. *Continued.*

(CONT.) RAINBOW TROUT 917 TESTS

1	BENZENE HEXACHLORIDE
1	BENZOTHIAZOLE 2-MERCAPTO
1	BENZOTHIAZOLE 2-METHYL MERCAPTO
3	BENZOYLPROP ETHYL
1	BINAPACRYL
2	BOMYL
1	BUTYLATE
2	CAPTAFOL
1	CAPTAN
14	CARBARYL
1	CARBOFURAN
1	CCA TYPE III
1	CHLORAMBEN
1	CHLORBROMURON
19	CHLORDANE
3	CHLORDANE-HCS-3260
2	CHLORDECONE
1	CHLORDIMEFORM
3	CHLORENDATE DIBUTYL
3	CHLORENDATE DIMETHYL
1	CHLORFENVINPHOS
1	CHLORINATED ALPHA OLEFIN 22/44
1	CHLORMEQUAT CHLORIDE
1	CHLOROBENZILATE
1	CHLORONITROPROPANE
1	CHLOROWAX LV
1	CHLOROWAX 40
1	CHLOROWAX 50
6	CHLOROWAX 500C
1	CHLOROWAX 70
1	CHLOROXURON
1	ETHANOL
1	ETHION
2	ETHOFUMESATE
1	ETHYLAN
1	ETHYLBENZENE
1	ETHYLENE DICHLORIDE
2	ETHYLENE GLYCOL
2	FENAC
1	FENAMINOSULF
1	FENAZAFLOR
2	FENBUTATIN-OXIDE
21	FENITROTHION
1	FENSON
2	FENTHION
11	FENVALERATE
4	FIRE-TROL 100
4	FIRE-TROL 931
2	FLAMPROP-METHYL
2	FLUCHLORALIN
3	FLUOMETURON
1	FLUORODIFEN
10	FLURIDONE
2	FOLPET
1	FONOFOS
1	FOSAMINE AMMONIUM
2	FOSPIRATE
9	FYROQUEL GT
1	GERANIOL
1	GLENBAR
1	GLYCEROL
10	GLYPHOSATE
6	HEPTACHLOR
1	HEPTACHLOR EPOXIDE
1	HEXACHLOROBUTADIENE
2	HEXAZINONE
3	HOUGHTO-SAFE 1120
4	HOUGHTO-SAFE 520
1	JODFENPHOS
9	KRONITEX 200
1	DICHLOROPHENYL METHYLSULFONATE
1	DICHLORVOS
1	DICLOFOP METHYL
1	DICOFOL
1	DICROTOPHOS
6	DIELDRIN
8	DIFLUBENZURON
2	DILAN
2	DIMETHOATE
1	DIMETHYL FORMAMIDE
1	DIMETHYLSULFOXIDE
2	DINITRAMINE
1	DINOBUTON
2	DINOCAP
1	DIOXATHION
1	DIPHENAMIDE
1	DIQUAT
1	DISULFOTON
11	DIURON
1	DNOC
2	EMCOL AD-410
4	ENDOSULFAN
1	ENDOTHALL
2	ENDOTHALL AQUATHOL K
1	ENDOTHALL COPPER SALT
1	ENDOTHALL DES-I-CATE
1	ENDOTHALL HERBICIDE 282
1	ENDOTHALL HYDROTHOL 191
1	ENDOTHALL POTASSIUM SALT
6	ENDRIN
1	EPN
2	MIREX
1	MOLINATE
3	MON 0818
1	MONOCROTOPHOS
2	MONOETHANOLAMINE
1	MSMA
5	NALED
2	NEODOL 25-9
1	NITRALIN
4	NITRAPYRIN
1	NONYLPHENYL DIPHENYL PHOSPHATE
2	ORTHO 11775
3	OVEX
2	OXAMYL
2	OXYDEMETON-METHYL
1	OXYTHIOQUINOX
1	PARA-DICHLOROBENZENE
1	PARAQUAT
4	PARATHION
1	PARATHION DITHIOATE ANALOGUE
2	PAROIL 1032
1	PAROIL 1048
1	PAROIL 160
3	PCB AROCLOR 1016
1	PCB AROCLOR 1242
1	PCB AROCLOR 1248
1	PCB AROCLOR 1254
1	PCB AROCLOR 1260
1	PCB 2-CHLOROBIPHENYL
1	PCB 3-CHLOROBIPHENYL
1	PCB 4-CHLOROBIPHENYL
4	PENTACHLOROPHENOL
6	PENTACHLOROPHENOL SODIUM SALT
10	PERMETHRIN
2	PHORATE
1	PHORAZETIM
4	PHOS-CHEK 202
4	PHOS-CHEK 259
1	PHOSALONE

BEST AVAILABLE COPY

ED_005172C_00000035-00050

Table A3. *Continued.*

(CONT.) RAINBOW TROUT 917 TESTS

1	LANDRIN	9	PHOSFLEX 31P
11	LEPTOPHOS	18	PHOSMET
1	LIME SULFUR	1	PHOSPHAMIDON
5	LINDANE	1	PHOXIM
5	MALATHION	1	PHTHALATE DI-2-ETHYLHEXYL
4	MBC	4	PHTHALATE DIBUTYL
1	MCPB	14	PICLORAM
5	MERPHOS	3	PIPERONYL BUTOXIDE
1	METHANOL	2	PROFENOFOS
1	METHIDATHION	1	PROPANIL
1	METHIOCARB	1	PROPHAM
23	METHOMYL	1	PROPOXUR
2	METHOPRENE	1	PROPYLENE GLYCOL
10	METHOXYCHLOR	2	PURIFLOC C-3
2	METHYL PARATHION	2	PYDRAUL 115E
1	METHYL TRITHION	3	PYDRAUL 50E
1	METRIBUZIN	1	PYRETHRUM
1	MEVINPHOS	2	RESMETHRIN
10	MEXACARBATE	2	RONNEL
1	ROTENONE	1	TRANID
1	ROWMATE	1	TREFMID
1	RU-11679	2	TRIALLATE
2	RYANIA	83	TRICHLORFON
1	RYLEX D (2-HYDROXY-4-DODECOXYBENZOPHENONE)	1	TRICHLORONATE
1	RYLEX H (2,4-DIHYDROBENZOPHENONE)	1	TRICLOPYR
1	RYLEX NBC (NICKEL DIBUTYLDITHIO CARBAMATE)	2	TRICRESYL PHOSPHATE
6	SANTICIZER 148	25	TRIFLURALIN
9	SANTICIZER 154	1	TRIPHENYL PHOSPHATE
2	SD 16898	1	TRIPHENYLTIN HYDROXIDE
1	SD 17250	1	UC 10854
1	SD 7438	4	UREABOR
1	SD 8339	1	VERNOLATE
2	SILVEX	4	WATER GLYCOLS 894-44A
1	SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER	4	WATER GLYCOLS 894-44B
1	SIMAZINE	4	WATER GLYCOLS 894-44C
1	SODIUM ARSENITE	4	XYLENE
9	TEMEPHOS	1	2,3,4,6-TETRACHLOROPHENOL
1	TEPA	1	2,4-D
1	TEPP	4	2,4-D DIMETHYLAMINE SALT
10	TERBUFOS	8	2,4-D DODECYL/TETRADODECYL AMINE SALT
1	TERBUTRYN	2	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER
1	TERPENE POLYCHLORINATES	1	2,4-D/2,4,5-T 24%/28%
1	TETRACHLORVINPHOS	1	2,4-D/2,4,5-T 30%/28%
2	TETRADIFON	1	2,4-D/2,4,5-T 34%/17%
1	TETRAMINE	2	2,4-DB
1	TETRASUL	1	2,4,5-T TRIETHYLAMINE SALT
1	TH 285-N	1	2,6-DICHLOROBENZOIC ACID
1	TH 336-N	1	2,6-DIOCTADECYL-P-CRESOL
1	TOLUENE	1	6-CHLORO-2-PICOLINIC ACID
7	TOXAPHENE		

RAINBOW TROUT DONALDSON 2 TESTS

1	ANTIMYCIN A	1	CARBARYL

RAINBOW TROUT IOWA 2 TESTS

1	ANTIMYCIN A	1	CARBARYL

RAINBOW TROUT MISSOURI 9 TESTS

6	ANTIMYCIN A	1	CHLORDANE
1	CARBARYL	1	MALATHION

RAINBOW TROUT MT. WHITNEY 3 TESTS

3	ANTIMYCIN A		

RAINBOW TROUT NEW HAMPSHIRE 3 TESTS

1	ANTIMYCIN A	1	CHLORDANE
1	CARBARYL		

BEST AVAILABLE COPY

ED_005172C_00000035-00051

Table A3. *Continued.*

RAINBOW TROUT SOAP LAKE — 4 TESTS

	ANTIMYCIN A	1	CHLORDANE
	CARBARYL	1	MALATHION

RAINBOW TROUT STEELHEAD — 13 TESTS

	CARBOFURAN	3	PYRETHRUM
	CHLORDANE	2	RESMETHRIN
	D-TRANS ALLETHRIN	2	RU-11679
	PHOXIM	1	TRICHLORFON

RAINBOW TROUT WYTHEVILLE — 2 TESTS

	ANTIMYCIN A	1	CARBARYL

ATLANTIC SALMON — 131 TESTS

	ACEPHATE	5	METHOXYCHLOR
2	AMINOCARB	3	MEXACARBATE
17	AZINPHOS-METHYL	1	PCB AROCLOR 1016
15	CARBARYL	3	PHOXIM
1	DDE	1	PYRETHRUM
1	DDT	2	RU-11679
1	DIFLUBENZURON	21	TEMEPHOS
19	FENITROTHION	11	TRICHLORFON
10	METHOMYL		

BROWN TROUT — 77 TESTS

13	AMINOCARB	3	LINDANE
6	AZINPHOS-METHYL	1	MALATHION
2	CAPTAN	1	METHYL PARATHION
2	CARBARYL	6	MEXACARBATE
2	CARBOFURAN	1	PCB AROCLOR 1016
2	CHLORDANE	1	PCB AROCLOR 1260
1	DDT	3	PHOXIM
2	DINITRAMINE	3	PYDRAUL 50E
15	FENITROTHION	2	PYRETHRUM
1	FENTHION	1	RESMETHRIN
2	FOLPET	1	RU-11679
1	GERANIOL	1	TOXAPHENE
2	HEXACHLOROBENZENE	1	TRICHLORFON
1	LEPTOPHOS		

BROOK TROUT — 201 TESTS

11	ACEPHATE	88	METHOXYCHLOR
16	AMINOCARB	1	MEXACARBATE
1	AZINPHOS-METHYL	1	PCB AROCLOR 1016
9	CARBARYL	3	PERMETHRIN
2	CHLORPYRIFOS-METHYL	3	PHOXIM
7	DIFLUBENZURON	1	PYDRAUL 50E
23	FENITROTHION	6	TEMEPHOS
1	HEXACHLOROBENZENE	1	TFM
2	HEXAZINONE	3	TOXAPHENE
1	HOUGHTO-SAFE 1120	18	TRICHLORFON
3	METHOMYL		

LAKE TROUT — 146 TESTS

1	ANTIMYCIN A	2	MALATHION
3	CAPTAN	2	METHOXYCHLOR
5	CARBARYL	2	METHYL PARATHION
1	CARBOFURAN	1	MEXACARBATE
6	CHLORPYRIFOS	1	NALED
1	COUMAPHOS	2	PARATHION
2	D-TRANS ALLETHRIN	2	PCB AROCLOR 1016
1	DIAZINON	13	PICLORAM
2	DICHLORVOS	3	PYDRAUL 50E

Table A3. C

1	DICOFOL
1	DINITRAM
10	DINOSEB
15	DIURON
8	EPTC
3	FENTHION
1	FLIT MLO
2	FOLPET
1	LEPTOPHO
2	LETHANE
2	LINDANE

1	AZINPHOS
1	D-TRANS
1	DDT
1	DINITRAM
1	EPN
1	FENTHION
1	HEPTACHL

3	ANTIMYCI
1	AZINPHOS
1	BENEFIN
1	BENZENE
1	BUFENCAI
2	CAPTAFOI
2	CARBARYI
1	COPPER :
2	DDT
2	DICHLOB
1	DICHLOR
2	DIELDRI
1	DINOCAP
1	DIPHENA
1	DIQUAT
1	DITHIAN

1	AZINPHO
1	CARBARY
1	DDT
1	DINITRA
1	ENDRIN
1	FENITRO

1	DDT

2	ACEPHA
1	AKTON
1	ALDRIN
1	ALUMIN
1	ALUMIN
1	AMDRO
2	AMINOC
1	AMITRO
9	ANTIMY
3	AZINPH
1	BENEFI
4	BENOMI
1	BENZEI

BEST AVAILABLE COPY

ED_005172C_00000035-00052

Table A3. *Continued.*

(CONT.) LAKE TROUT — 146 TESTS

1	DICOFOL	2	PYRETHRUM
1	DINITRAMINE	2	RESMETHRIN
10	DINOSEB	2	RONNEL
15	DIURON	2	RU-11679
8	EPTC	4	TEMEPHOS
3	FENTHION	2	THANITE
1	FLIT MLO	2	TRICHLORFON
2	FOLPET	6	2,4-D
1	LEPTOPHOS	14	2,4-D BUTYL ESTER
2	LETHANE 384	14	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER
2	LINDANE		

NORTHERN PIKE — 15 TESTS

1	AZINPHOS-METHYL	1	METHOXYCHLOR
1	D-TRANS ALLETHRIN	1	METHYL PARATHION
1	DDT	1	PHORATE
1	DINITRAMINE	2	PHOXIM
1	EPN	1	PYRETHRUM
1	FENTHION	1	RESMETHRIN
1	HEPTACHLOR	1	RU-11679

GOLDFISH — 45 TESTS

3	ANTIMYCIN A	1	ENDRIN
1	AZINPHOS-METHYL	1	FENITROTHION
1	BENEFIN	2	FENTHION
1	BENZENE HEXACHLORIDE	2	LINDANE
1	BUFENCARB	1	MALATHION
2	CAPTAFOL	2	METHOXYCHLOR
2	CARBARYL	1	METHYL PARATHION
1	COPPER SULFATE	2	MEXACARBATE
2	DDT	1	MSMA
2	DICHLOBENIL	1	NOREA
1	DICHLOROPROPENE	2	PARATHION
2	DIELDRIN	1	PIPERALIN
1	DINOCAP	1	SILVEX BUTOXYETHANOL ESTER
1	DIPHENAMIDE	2	TOXAPHENE
1	DIQUAT	1	TRIFLURALIN
1	DITHIANON	1	TRIPHENYLTIN HYDROXIDE

CARP — 12 TESTS

1	AZINPHOS-METHYL	1	FENTHION
1	CARBARYL	1	LINDANE
1	DDT	1	MALATHION
1	DINITRAMINE	1	METHYL PARATHION
1	ENDRIN	1	MEXACARBATE
1	FENITROTHION	1	TOXAPHENE

RIVER SHINER — 1 TESTS

1	DDT

FATHEAD MINNOW — 280 TESTS

2	ACEPHATE	1	FENBUTATIN-OXIDE
1	AKTON	5	FENITROTHION
1	ALDRIN	3	FENTHION
1	ALUMINUM NITRATE	2	FENVALERATE
1	ALUMINUM SULFATE	1	FIRE-TROL 100
1	AMDRO	1	FIRE-TROL 931
2	AMINOCARB	2	FLURIDONE
1	AMITROLE	2	FOLPET
9	ANTIMYCIN A	1	FOSAMINE AMMONIUM
3	AZINPHOS-METHYL	1	FOSPIRATE
1	BENEFIN	1	FYROQUEL GT
6	BENOMYL	2	GERANIOL
1	BENZENE HEXACHLORIDE	5	GLYPHOSATE

BEST AVAILABLE COPY

ED_005172C_00000035-00053

Table A3. *Continued.*

(CONT.) FATHEAD MINNOW 280 TESTS

3	BENZOYLPROP ETHYL	1	HALOWAX 1099	1
2	CALCIUM HYPOCHLORITE	1	HEPTACHLOR	2
2	CAPTAN	1	HEXACHLOROBENZENE	3
3	CARBARYL	2	HOUGHTO-SAFE 1120	1
3	CARBOFURAN	1	HOUGHTO-SAFE 520	2
2	CHLORDANE	1	LEPTOPHOS	1
1	CHLORDANE-HCS-3260	1	LIME SULFUR	2
2	CHLORDECONE	4	LINDANE	1
1	CHLORENDATE DIBUTYL	2	MALATHION	
2	CHLORENDATE DIMETHYL	1	MBC	
2	CHLORFENETHOL	1	MCPB	
1	CHLORMEQUAT CHLORIDE	3	METHOMYL	2
1	CHLOROWAX 500C	2	METHOPRENE	1
1	CHLORPYRIFOS-METHYL	1	METHOXYCHLOR	1
1	COPPER SULFATE	3	METHYL PARATHION	1
1	COPPER-COUNT-N	2	METOLACHLOR	1
1	COPPER-COUNT-NS	2	MEXACARBATE	1
1	CORREX	1	MIREX	5
1	CROTOXYPHOS	1	MITIN FF	2
1	CUMYLPHENYL DIPHENYL PHOSPHATE	1	MON 0818	1
4	CYANAZINE	1	MONOCROTOPHOS	9
2	CYPRAZINE	1	MSMA	1
2	D-TRANS ALLETHRIN	1	NALED	1
1	DDD	2	NEODOL 25-12	1
3	DDT	1	NEODOL 25-9	6
2	DICHLOBENIL	2	NITRAPYRIN	2
1	DICHLOROPHENYL METHYLSULFONATE	1	NONYLPHENYL DIPHENYL PHOSPHATE	1
1	DICHLOROPROPENE	1	NOREA	1
1	DICHLORVOS	2	ORTHO 11775	1
1	DIELDRIN	2	PARATHION	1
1	DIFLUBENZURON	1	PENTACHLOROPHENOL	1
1	DIMETHRIN	1	PENTACHLOROPHENOL SODIUM SALT	1
2	DINITRAMINE	2	PERMETHRIN	1
1	DIPHENAMIDE	1	PHOS-CHEK 202	3
1	DISULFOTON	1	PHOS-CHEK 259	1
1	DITHIANON	1	PHOSALONE	1
1	ENDOSULFAN	1	PHOSFLEX 31P	1
1	ENDOTHALL HYDROTHOL 191	2	PHOSMET	1
2	ENDRIN	1	PHOSPHAMIDON	5
1	ETHION	1	PHOXIM	1
1	PHTHALATE DI-2-ETHYLHEXYL	1	TRICHLORFON	1
2	PHTHALATE DIBUTYL	3	TRIFLURALIN	1
1	PROPOXUR	1	TRIPHENYL PHOSPHATE	1
2	PURIFLOC C-3	1	TRIPHENYLTIN HYDROXIDE	2
1	PYDRAUL 115E	1	UREABOR	1
3	PYDRAUL 50E	1	WATER GLYCOLS 894-44A	1
2	PYRETHRUM	1	WATER GLYCOLS 894-44B	2
3	RESMETHRIN	1	WATER GLYCOLS 894-44C	1
2	RU-11679	1	2-HYDROXY-2',4,4'-TRICHLORODIPHENYL ETHER	1
2	S-BIOALLETHRIN	1	2-HYDROXY-4,4'-DICHLORODIPHENYL ETHER	1
1	SANTICIZER 154	1	2,3,4,6-TETRACHLOROPHENOL	1
2	SILVEX	1	2,3,5-TRIMETHYL NAPHTHALENE	1
1	SILVEX BUTOXYETHANOL ESTER	1	2,3,6-TRICHLOROBENZOIC ACID	1
1	SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER	1	2,3,6-TRIMETHYL NAPHTHALENE	1
3	SIMAZINE	1	2,4-D	1
1	SODIUM SELENITE	1	2,4-D BUTOXYETHANOL ESTER	1
1	TEMEPHOS	24	2,4-D DIMETHYLAMINE SALT	2
1	TEPP	18	2,4-D DODECYL/TETRADODECYL AMINE SALT	1
2	TERBUFOS	1	2,4-DB	1
10	TOXAPHENE	1	2,4,5-T TRIETHYLAMINE SALT	1
1	TRI-N-BUTYL PHOSPHATE			

LONGNOSE SUCKER 1 TESTS

1	PCB AROCLOR 1016

WHITE SUCKER 8 TESTS

2	CHLORDANE	2	RESMETHRIN
1	D-TRANS ALLETHRIN	2	RU-11679
1	PCB AROCLOR 1016		

BEST AVAILABLE COPY

ED_005172C_00000035-00054

Table A3. *Continued.*

BLACK BULLHEAD — 25 TESTS

1	ALDRIN
2	ANTIMYCIN A
3	AZINPHOS-METHYL
1	CARBARYL
2	DDT
1	DIELDRIN
2	DIQUAT
1	ENDRIN
2	FENTHION
1	HEPTACHLOR
1	LINDANE
2	MALATHION
1	METHYL PARATHION
2	MEXACARBATE
2	TOXAPHENE
1	TRICHLORFON

CHANNEL CATFISH — 341 TESTS

2	ACEPHATE
1	AERO XANTHATE 343
1	AEROFROTH 71
1	AKTON
1	ALDRIN
1	AMDRO
5	AMINOCARB
2	AMITROLE
1	ANILAZINE
9	ANTIMYCIN A
1	AQUA COP
*	AQUA-VATOR
1	AZINPHOS-METHYL
6	BENOMYL
2	BENTHIOCARB
1	BENZENE
1	BENZENE HEXACHLORIDE
1	BENZOTHIAZOLE 2-MERCAPTO
1	BENZOTHIAZOLE 2-METHYL MERCAPTO
1	BINAPACRYL
1	CAPTAFOL
1	CAPTAN
3	CARBARYL
1	CARBOFURAN
1	CARBOPHENOTHION
1	CCA TYPE III
1	CHLORBROMURON
5	CHLORDANE
1	CHLORDANE-HCS-3260
1	CHLORDECONE
1	CHLORDIMEFORM
1	CHLORENDATE DIBUTYL
2	CHLORENDATE DIMETHYL
1	CHLORFENETHOL
1	CHLORMEQUAT CHLORIDE
2	CHLOROWAX 500C
1	CHLOROXURON
1	CHLORPYRIFOS
2	CORREX
1	COUMAPHOS
1	CROTOXYPHOS
1	CUMYLPHENYL DIPHENYL PHOSPHATE
3	CYANAZINE
2	CYTROL AMITROLE-T
1	D-D SOIL FUMIGANT
2	D-TRANS ALLETHRIN
1	DDD
6	DDT
2	DEF
1	DEMETON
1	DICHLOFENTHION
1	DICOFOL
1	DICROTOPHOS
2	DIELDRIN
2	DIFLUBENZURON
2	DIMETHRIN
1	DINITRAMINE
1	DISULFOTON
1	DITHIANON
1	ENDOSULFAN
1	ENDOTHALL
2	ENDOTHALL AQUATHOL K
1	ENDOTHALL HYDROTHOL 191
2	ENDRIN
1	EPN
1	ETHION
1	ETHYLBENZENE
2	FENBUTATIN-OXIDE
8	FENITROTHION
2	FENTHION
2	FENVALERATE
3	FLAMPROP-METHYL
2	FLUCHLORALIN
3	FLUOMETURON
1	FLUORODIFEN
10	FLURIDONE
1	FOLPET
1	FOSAMINE AMMONIUM
1	FYRQUEL GT
6	GLYPHOSATE
1	HALOWAX 1099
1	HEPTACHLOR
3	HEXACHLOROBENZENE
1	HEXACHLOROBUTADIENE
3	HOUGHTO-SAFE 1120
1	JODFENPHOS
1	LEAD ARSENATE
1	LEPTOPHOS
2	LINDANE
2	LINURON
2	MALATHION
5	MBC
2	MERPHOS
6	METHOMYL
1	METHOPRENE
1	METHOXYCHLOR
1	METHYL PARATHION
1	METHYL TRITHION
2	METRIBUZIN
1	MEVINPHOS
2	MEXACARBATE
1	MON 0818
1	MONOCROTOPHOS
1	MSMA
1	NALED
1	NEODOL 25-12
1	NEODOL 25-9
1	NITRAPYRIN
1	ORTHO 11775
2	OXAMYL
1	OXYDEMETON-METHYL
1	PARAQUAT
1	PARATHION
1	PARATHION DITHIOATE ANALOGUE
1	PAROIL 1032
1	PAROIL 1048

BEST AVAILABLE COPY

ED_005172C_00000035-00055

Table A3. *Continued.*

(CONT.) CHANNEL CATFISH 341 TESTS

1	PAROIL 160	2	RU-11679
3	PCB AROCLOR 1016	1	RYANIA
1	PCB AROCLOR 1242	1	S-BIOALLETHRIN
1	PCB AROCLOR 1248	2	SILVEX
1	PCB AROCLOR 1254	1	SODIUM SELENITE
1	PCB AROCLOR 1260	5	TEMEPHOS
2	PENTACHLOROPHENOL	1	TERBUFOS
2	PENTACHLOROPHENOL COPPER SALT	1	TETRACHLORVINPHOS
2	PENTACHLOROPHENOL SODIUM SALT	1	TETRADIFON
1	PERMETHRIN	1	TOLUENE
1	PHORATE	26	TOXAPHENE
1	PHOSFLEX 31P	2	TRIALLATE
2	PHOSMET	3	TRICHLORFON
1	PHOSPHAMIDON	1	TRICHLOROMANDELIC ACID
1	PHOTO-DIELDRIN	1	TRICRESYL PHOSPHATE
1	PHOXIM	6	TRIFLURALIN
2	PHTHALATE DI-2-ETHYLHEXYL	1	TRIPHENYL PHOSPHATE
2	PHTHALATE DIBUTYL	1	UREABOR
7	PICLORAM	1	WARFARIN
2	PROFENOFOS	2	XYLENOL DIMETHYLAMINO
2	PROPACHLOR	1	Z-200
1	PROPYL ISOME	1	2,3,4,6-TETRACHLOROPHENOL
1	PURIFLOC C-3	2	2,4-D BUTOXYETHANOL ESTER
3	PYDRAUL 115E	6	2,4-D DIMETHYLAMINE SALT
4	PYDRAUL 50E	2	2,4-D DODECYL/TETRADODECYL AMINE SALT
4	PYRETHRUM	1	2,4-D OLEYLPROPYLENEDIAMINE SALT
3	RESMETHRIN	1	2,4-D/2,4,5-T 24%/28%
1	RONNEL	1	2,4,5-T TRIETHYLAMINE SALT
2	ROTENONE	1	6-CHLORO-2-PICOLINIC ACID

WALKING CATFISH 1 TESTS

1	ANTIMYCIN A

MOSQUITOFISH 2 TESTS

1	DICHLORVOS	1	ENDRIN

MOSQUITOFISH NON-RESISTANT 2 TESTS

1	ANTIMYCIN A	1	PARATHION

MOSQUITOFISH RESISTANT 2 TESTS

1	ANTIMYCIN A	1	PARATHION

GREEN SUNFISH 32 TESTS

1	ANTIMYCIN A	1	FENITROTHION
1	AZINPHOS-METHYL	3	FENTHION
1	BINAPACRYL	2	LINDANE
2	CARBARYL	3	MALATHION
1	CARBOPHENOTHION	2	METHYL PARATHION
2	COPPER SULFATE	2	MEXACARBATE
3	DDT	2	PARATHION
2	DICHLOBENIL	2	TOXAPHENE
2	ENDOTHALL COPPER SALT		

BLUEGILL 959 TESTS

7	ACEPHATE	1	CHLOROWAX 70
1	AERO XANTHATE 343	5	CHLORPYRIFOS
1	AEROFROTH 71	1	COPPER OXYCHLORIDE
1	AKTON	2	COPPER SULFATE
2	ALACHLOR	1	COPPER-COUNT-N
2	ALDICARB	1	COPPER-COUNT-NS
6	ALDRIN	2	CORREX
1	ALLETHRIN	1	COUMAPHOS
1	AMDRO	1	CROTOXYPHOS

BEST AVAILABLE COPY

ED_005172C_00000035-00056

Table A3. *Continued.*

(CONT.) BLUEGILL 959 TESTS

1	AMETRYN
21	AMINOCARB
1	ANILAZINE
17	ANTIMYCIN A
1	APHOLATE
1	AQUA COP
1	AQUA-VATOR
1	ARAMITE
1	ATRAZINE 4L
1	AZIDE POTASSIUM
1	AZIDE SODIUM
1	AZINPHOS-ETHYL
7	AZINPHOS-METHYL
1	BACILLUS THURINGIENSIS
12	BENOMYL
1	BENSULIDE
2	BENTHIOCARB
1	BENZALKONIUM CHLORIDE
9	BENZENE
1	BENZENE HEXACHLORIDE
4	BENZOTHIAZOLE 2-MERCAPTO
4	BENZOTHIAZOLE 2-METHYL MERCAPTO
3	BENZOYLPROP ETHYL
1	BINAPACRYL
1	BOMYL
3	BUTYLATE
1	CACODYLIC ACID
2	CAPTAFOL
1	CAPTAN
14	CARBARYL
2	CARBOFURAN
1	CARBOPHENOTHION
1	CCA TYPE III
1	CHLORAMBEN
3	CHLORDANE
1	CHLORDANE CIS
1	CHLORDANE TRANS
3	CHLORDANE-HCS-3260
1	CHLORDECONE
2	CHLORENDATE DIBUTYL
2	CHLORENDATE DIMETHYL
2	CHLORFENVINPHOS
1	CHLORINATED ALPHA OLEFIN 22/44
1	CHLORMEQUAT CHLORIDE
1	CHLORONITROPROPANE
1	CHLOROWAX LV
1	CHLOROWAX 40
1	CHLOROWAX 50
4	CHLOROWAX 500C
7	ENDRIN
1	EPN
1	ETHION
2	ETHOFUMESATE
1	ETHYLAN
10	ETHYLBENZENE
1	ETHYLENE GLYCOL
2	FENAC
1	FENAMINOSULF
1	FENAZAFLOR
1	FENBUTATIN-OXIDE
24	FENITROTHION
1	FENSON
1	FENSULFOTHION
2	FENTHION
11	FENVALERATE
1	FIRE-TROL 100
1	FIRE-TROL 931
2	FLAMPROP-METHYL
2	FLUCHLORALIN
2	FLUOMETURON
2	CRUFOMATE
1	CRYOLITE
2	CUTRINE-PLUS
2	CYANAZINE
4	CYHEXATIN
2	CYPRAZINE
3	CYTROL AMITROLE-T
1	D-D SOIL FUMIGANT
13	D-TRANS ALLETHRIN
1	DALAPON
1	DCPA
1	DDD
1	DDE
7	DDT
1	DEAD-X
11	DEF
1	DEMETON
1	DIAZINON
1	DICAMBA
2	DICHLOBENIL
1	DICHLOFENTHION
1	DICHLONE
1	DICHLORVOS
1	DICLOFOP METHYL
1	DICOFOL
1	DICROTOPHOS
8	DIELDRIN
8	DIFLUBENZURON
1	DILAN
1	DIMETHOATE
14	DIMETHRIN
1	DIMETHYLSULFOXIDE
1	DINITRAMINE
1	DINOBUTON
2	DINOCAP
1	DIOXATHION
1	DIPHENAMIDE
10	DIQUAT
1	DISULFOTON
12	DIURON
1	DNOC
1	EMCOL AD-410
1	ENDOSULFAN
1	ENDOTHALL
9	ENDOTHALL AQUATHOL K
1	ENDOTHALL COPPER SALT
1	ENDOTHALL HERBICIDE 282
1	ENDOTHALL HYDROTHOL 191
1	ENDOTHALL POTASSIUM SALT
2	MIREX
1	MITIN FF
1	MNFA
1	MOLINATE
4	MON 0818
1	MONOCROTOPHOS
1	MONOETHANOLAMINE
3	MSMA
1	N-BUTANOL
1	N,N-DIMETHYL-2,4-DICHLOROPHENOXYACETAMIDE
1	NALED
1	NITRALIN
1	NOREA
2	ORTHO 11775
1	OVEX
2	OXAMYL
1	OXYDEMETON-METHYL
1	OXYTHIOQUINOX
1	PARAQUAT
7	PARATHION
1	PARATHION DITHIOATE ANALOGUE

BEST AVAILABLE COPY

ED_005172C_00000035-00057

Table A3. *Continued.*

(CONT.) BLUEGILL 959 TESTS

1	FLUORENE
2	FLURIDONE
1	FOLPET
1	FONOFOS
1	FOSAMINE AMMONIUM
9	FYRQUEL GT
16	GLYPHOSATE
3	HEPTACHLOR
1	HEPTACHLOR EPOXIDE
1	HERCULES 7175
2	HEXACHLOROBENZENE
1	HEXACHLOROBUTADIENE
1	HEXAZINONE
3	HOUGHTO-SAFE 1120
1	HOUGHTO-SAFE 520
1	HYDROXY-S-TRIAZINYL ALANINE
1	KOMEEN
9	KRONITEX 200
1	LANDRIN
2	LEPTOPHOS
1	LIME SULFUR
6	LINDANE
8	MALATHION
1	MCPA DIMETHYLAMINE SALT
1	MCPB
7	MERPHOS
1	METHIDATHION
1	METHIOCARB
21	METHOMYL
2	METHOPRENE
1	METHOPROTRYNE
5	METHOXYCHLOR
3	METHYL PARATHION
2	METHYL TRITHION
1	METRIBUZIN
3	MEVINPHOS
16	MEXACARBATE
11	S-BIOALLETHRIN
6	SANTICIZER 148
9	SANTICIZER 154
2	SD 16898
2	SD 17250
1	SD 7438
1	SD 8339
3	SILVEX
2	SILVEX BUTOXYETHANOL ESTER
2	SILVEX POTASSIUM SALT
2	SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER
1	SIMAZINE
1	SODIUM ARSENITE
4	TEMEPHOS
3	TEPP
10	TERBUFOS
1	TERBUTRYN
2	TERPENE POLYCHLORINATES
2	TETRACHLORVINPHOS
1	TETRADIFON
1	TETRAMINE
1	TH 285-N
1	TH 336-N
9	TOLUENE
12	TOXAPHENE
1	TRANID
2	TREFMID
65	TRICHLORFON
1	TRICHLORONATE
1	PAROIL 1032
1	PAROIL 1048
1	PAROIL 160
2	PCB AROCLOR 1016
1	PCB AROCLOR 1242
2	PCB AROCLOR 1248
2	PCB AROCLOR 1254
1	PCB AROCLOR 1260
2	PENTACHLOROPHENOL
2	PENTACHLOROPHENOL SODIUM SALT
10	PERMETHRIN
6	PHORATE
1	PHOS-CHEK 202
1	PHOS-CHEK 259
1	PHOSALONE
9	PHOSFLEX 31P
33	PHOSMET
1	PHOSPHAMIDON
1	PHOTO-DIELDRIN
14	PHOXIM
2	PHTHALATE DI-2-ETHYLHEXYL
5	PHTHALATE DIBUTYL
10	PICLORAM
2	PIPERONYL BUTOXIDE
2	PROFENOFOS
1	PROPANIL
1	PROPHAM
1	PROPOXUR
2	PURITIOC C-3
1	PYDRAUL 115E
3	PYDRAUL 50E
15	PYRETHRUM
3	RESMETHRIN
3	RONNEL
1	ROTENONE
2	RU-11679
1	RYANIA
1	TRICLOPYR
2	TRICRESYL PHOSPHATE
16	TRIFLURALIN
1	TRIPHENYLTIN HYDROXIDE
1	TRITON B-1956
1	UC 10854
4	UREABOR
2	VERNOLATE
1	WATER GLYCOLS 894-44A
1	WATER GLYCOLS 894-44B
1	WATER GLYCOLS 894-44C
12	XYLENE
2	XYLENOL AMINO
2	XYLENOL DIMETHYLAMINO
1	Z-200
1	2-ETHYLHEXANOL
1	2,3,4,6-TETRACHLOROPHENOL
1	2,4-D
3	2,4-D BUTOXYETHANOL ESTER
8	2,4-D DIMETHYLAMINE SALT
5	2,4-D DODECYL/TETRADODECYL AMINE SALT
1	2,4-D OLEYLPROPYLENEDIAMINE SALT
2	2,4-D PROPYLENE GLYCOL BUTYL ETHER EST
2	2,4-D/2,4,5-T 24%/28%
1	2,4-D/2,4,5-T 30%/28%
1	2,4-D/2,4,5-T 34%/17%
2	2,4-DB
1	2,4,5-T TRIETHYLAMINE SALT
1	2,6-DICHLOROBENZOIC ACID

BEST AVAILABLE COPY

ED_005172C_00000035-00058

Table A3. *Continued.*

REDEAR SUNFISH — 18 TESTS

1	AKTON	5	HEPTACHLOR
1	ANILAZINE	1	MALATHION
5	CHLORDECONE	1	SILVEX BUTOXYETHANOL ESTER
1	DDT	2	2,4,5-T BUTOXYETHANOL ESTER
1	FENAC		

SMALLMOUTH BASS — 9 TESTS

1	D-TRANS ALLETHRIN	1	RU-11679
1	FOLPET	1	2,4-D DIMETHYLAMINE SALT
1	PHOSMET	2	2,4-D DODECYL/TETRADODECYL AMINE SALT
2	PYRETHRUM		

LARGEMOUTH BASS — 74 TESTS

1	ALDRIN	1	FIRE-TROL 931
1	AMINOCARB	1	HEPTACHLOR
5	ANTIMYCIN A	1	HEXACHLOROBENZENE
1	AZINPHOS-METHYL	1	LEPTOPHOS
1	BENZENE HEXACHLORIDE	1	LINDANE
1	CARBARYL	2	MALATHION
5	CHLORDANE	2	METHOMYL
1	COUMAPHOS	2	METHOXYCHLOR
1	CROTOXYPHOS	1	METHYL PARATHION
2	CYHEXATIN	1	METHYL TRITHION
1	D-D SOIL FUMIGANT	1	MEVINPHOS
1	D-TRANS ALLETHRIN	1	MEXACARBATE
1	DDD	1	MIREX
2	DDT	1	NALED
1	DEMETON	1	OXYDEMETON-METHYL
1	DICHLOFENTHION	1	PARATHION
1	DICHLOROPROPENE	1	PHORATE
1	DICOFOL	1	PHOS-CHEK 202
1	DIELDRIN	1	PHOS-CHEK 259
2	DIOXATHION	1	PHOSMET
1	DISULFOTON	1	PYRETHRUM
1	ENDRIN	1	RESMETHRIN
1	EPN	1	RU-11679
1	ETHION	4	TEMEPHOS
1	FENAZAFLOR	2	TOXAPHENE
3	FENTHION	1	TRICHLORFON
1	FIRE-TROL 100	1	TRIFLURALIN

WHITE CRAPPIE — 1 TESTS

1	ANTIMYCIN A

BLACK CRAPPIE — 3 TESTS

1	AZINPHOS-METHYL	1	DDT
1	CARBARYL		

YELLOW PERCH — 133 TESTS

7	ACEPHATE	1	METHYL PARATHION
14	AMINOCARB	3	MEXACARBATE
1	ANTIMYCIN A	1	MIREX
17	AZINPHOS-METHYL	1	PAROIL 1032
1	CAPTAN	1	PAROIL 1048
14	CARBARYL	1	PAROIL 160
3	CARBOFURAN	1	PCB AROCLOR 1016
1	CHLORDANE	1	PCB AROCLOR 1242
1	D-TRANS ALLETHRIN	1	PCB AROCLOR 1248
1	DDT	1	PCB AROCLOR 1254

BEST AVAILABLE COPY

ED_005172C_00000035-00059

Table A3. *Continued.*

(CONT.) YELLOW PERCH — 133 TESTS

2	DIFLUBENZURON	1	PCB AROCLOR 1260
1	DIMETHRIN	3	PHOXIM
2	DINITRAMINE	1	PHTHALATE DIBUTYL
2	DIQUAT	1	PYDRAUL 50E
1	ENDRIN	1	PYRETHRUM
13	FENITROTHION	1	RESMETHRIN
1	FENTHION	1	RU-11679
1	FOLPET	1	S-BIOALLETHRIN
1	HOUGHTO-SAFE 1120	1	TOXAPHENE
15	LEPTOPHOS	1	TRICHLORFON
2	LINDANE	1	TRICRESYL PHOSPHATE
1	MALATHION	2	XYLENOL DIMETHYLAMINO
4	METHOXYCHLOR		

WALLEYE — 17 TESTS

1	AMINOCARB	1	EPN
1	COUMAPHOS	1	MALATHION
1	D-D SOIL FUMIGANT	1	MIREX
1	DDD	1	OXYDEMETON-METHYL
2	DDT	2	PHORATE
1	DEMETON	1	PYRETHRUM
1	DICHLOFENTHION	1	TRIFLURALIN
1	DICHLOROPROPENE		

TILAPIA — 10 TESTS

3	DDT	2	MALATHION
2	DIELDRIN	1	METHOXYCHLOR
1	ENDRIN	1	METHYL TRITHION

FOWLERS TOAD — 25 TESTS

1	ALDRIN	1	LINDANE
1	AZINPHOS-METHYL	1	MALATHION
1	BENZENE HEXACHLORIDE	1	METHOXYCHLOR
1	DDD	1	MOLINATE
5	DDT	1	PARAQUAT
1	DCF	1	PARATHION
1	DICLORIN	1	SILVEX PROPYLENE GLYCOL BUTYL ETHER ES
1	ENDOTHALL HYDROTHOL 191	1	TOXAPHENE
1	ENDRIN	2	TRIFLURALIN
1	HEPTACHLOR	1	2,4-D BUTOXYETHANOL ESTER

WESTERN CHORUS FROG — 13 TESTS

1	AZINPHOS-METHYL	1	METHYL PARATHION
1	DDT	1	MEVINPHOS
1	DIELDRIN	1	PARAQUAT
1	ENDRIN	1	PARATHION
1	LINDANE	1	PIPERONYL BUTOXIDE
1	MALATHION	1	TOXAPHENE
1	METHOXYCHLOR		

BEST AVAILABLE COPY

ED_005172C_00000035-00060

Table A4. *Common and alternate names of chemicals used for aquatic toxicity testing.*

Common name	Alternate name
ACEPHATE	Orthene; Ortho 12420; Ortran
ACETONE	dimethylformaldehyde; dimethylketol; dimethyl ketone; ketone propane; methy ketone; propanone; pyroacetic acid; pyroacetic ether
AKTON	Axiom
ALACHLOR	Alanex; Pillarzo
ALDICARB	Temik; OMS 771; UC21149
ALDRIN	Aldrex 30; Aldrine; Aldrite; Aldrosol; Altox; Drinox; HHDN; Octalene
ALLETHRIN	Pallethrine; Pyna:nin
ALUMINUM NITRATE	Aluminum trinitrate; nitric acid aluminum salt; nitric acid aluminum (3+) salt
ALUMINUM SULFATE	Alum; aluminum sulfate (2:3); aluminum trisulfate; cake alum; dialuminum sulfate; dialuminum trisulfate; luminum alum; sulfuric acid aluminum salt (3:2)
AMDRO	AC 217,300; CL 217,300
AMETRYN	Ametrex; Crisatrine; Evik; G-34162; Gesapax
AMINOCARB	A 363; aminocarbe; Bay 44646; Matacil
AMITROLE	Amerol; Amino-triazole; Amino Triazole Weedkiller 90; Amitrol T; Amizol; AT-90; ATA; AT Liquid; Azolan; Azole; Cytrol; Cytrol Amitrole-T; Diurol; Herbizole; Simazol; Weedazol; Weedazol TL, Amitrol
ANILAZINE	Direz; Dyrene; Kemate; Triasyn
ANTIMYCIN A	Antipiricullin; Virosin
APHOLATE	APN; ENT-26,316; NSC-26812; OLIN MO. 2174; pholate; SQ 8388
ARAMITE	Aracide; Niagaramite; 88-R
AROCLOR 1221	Arochlor 1221; chlorodiphenyl (21% Cl); polychlorinated biphenyl (Aroclor 1221)
AROCLOR 1232	Arochlor 1232; chlorodiphenyl (32% Cl); polychlorinated biphenyl (Aroclor 1232)
AROCLOR 1242	Arochlor 1242; chlorodiphenyl (42% Cl); polychlorinated biphenyl (Aroclor 1242)
AROCLOR 1248	Arochlor 1248; chlorodiphenyl (48% Cl); polychlorinated biphenyl (Aroclor 1248)
AROCLOR 1254	Arochlor 1254; chlorodiphenyl (54% Cl); NCI-CO2664; polychlorinated biphenyl (Aroclor 1254)
AROCLOR 1260	Arochlor 1260; chlorodiphenyl (60% Cl); Clophen A60; Phenoclor DP6; polychlorinated biphenyl (Aroclor 1260)
AROCLOR 1262	Arochlor 1262; chlorodiphenyl (62% Cl); polychlorinated biphenyl (Aroclor 1262)
AROCLOR 1268	Arochlor 1268; chlorodiphenyl (68% Cl); polychlorinated biphenyl (Aroclor 1268)
AROCLOR 4465	Arochlor 4465; polychlorinated biphenyl (Aroclor 4465)
AROCLOR 5442	Arochlor 5442; polychlorinated triphenyl (Aroclor 5442)
AROCLOR 5460	Arochlor 5460; polychlorinated triphenyl (Aroclor 5460)
ATRAZINE 4L	AAtrex; Atranex; Atred; Crisatrina; Crisazine; Farmco Atrazine; Griffex; Shell Atrazine Herbicide; Vectal SC
AZIDE (SODIUM)	
AZINPHOS ETHYL	Azinos; Bay 16259; Cotnion-Ethyl; Crysthion; Gusathion A; Triazotion
AZINPHOS METHYL	Bay 17147; Carfene; Cotnion-Methyl; Gusathion M; Guthion; metiltriazotion
BARBAN	barbamate; barbane; Carbyne; chlorinat
BENEFIN	Balan; Balfin; benfluralin; Benefex; Bethrodine; Quilan
BENOMYL	Beniate; Tersan 1991
BENSULIDE	Betamec; Betasan; Exporsan; Prefar; Pre-San
BENTHIOCARB	Bolero; Saturn; Saturno; Siacarb; Tamariz; Thiobencarb
BENZALKONIUM CHLORIDE	Barquat MB-50; Barquat MB-80; BTC; Roccal; Zephiran
BENZENE	Benzol
BENZENE HEXACHLORIDE	Benzahex; Benzex; BHC; Dol; Dolmix; Gammexane; Gexane; HCCH; HCH; Hexablanc; hexachlor; hexachloran; Hexafor; Hexamul; Hexapoudre; Hexyclan; Hilbeech; Kotol; Soprocide; Submar; 666

BEST AVAILABLE COPY

ED_005172C_00000035-00061

Table A4. *Continued.*

Common name	Alternate name
BENZOTHIAZOLE 2-METHYL MERCAPTO	USAF EK-4008
BENZOYLPROP ETHYL	SD 30053; Suffix; Suffix 25; WL 17731
BENZYL BENZOATE	Ascabin; Ascabiol; Benzylets; Colebenz; Novoscabin; Peruscabin; Vanzoate; Venzonate
BINAPACRYL	Acricid; Ambox; Dapacryl; dinoseb; Endosan; FMC 9044; Hoe 2784; methacrylate; Morocide; Morrocid; NIA 9044
BOMYL	Fly Bait Grits; GC-3707; Swat
BUFENCARB	Bux; Ortho 5353
BUTOXYL POLYPROPYLENE GLYCOL	Stabilene Fly Repellent
BUTYLATE	R-1910; Sutan
CACODYLIC ACID	Dilic; Rad-E-Cate 25
CALCIUM HYPOCHLORITE	bleaching powder; chloride of lime
CAPTAFOL	Difolatan; Haipen; Merpafol; Pillartan; Sanspor
CAPTAN	Captane; Merpan; Orthocide; Pillarcap; Vondcaptan
CARBARYL	Carbamine; Cekubaryl; Denapon; Devicarb; Dicarbam; Hexavin; Karbaspray; Nac; Ravyon; Septene; Sevin; Tercyl; Tricarnam; UC 7744
CARBOFURAN	Bay 70143; Curaterr; D 1221; ENT 27164; FMC 10242; Furadan; NIA 10242; Yaltox
CARBOPHENOTHION	Dagadip; Garrathion; R 1303; Trithion
CHLORAMBEN	Amiben; Amiben DS; chlorambene; Ornamental Weeder
CHLORBROMURON	C6313; chlorobromuron; Maloran
CHLORDANE	Belt; Chlordan; Chlor Kil; Chlortox; Corodane; Kypchlor; Niran; Niran 5% granular bait; Octachlor; Octa-Klor; Ortho-Klor; Snyklor; Topiclor 20; Velsicol 1068
CHLORDANE CIS	*alpha*-Chlordane
CHLORDANE TRANS	*gamma*-Chlordane
CHLORDECONE	GC 1189; Kepone
CHLORDIMEFORM	Bermat; C 8514; ENT 27567; EP-333; Fundal; Galecron; SN 36268
CHLORFENETHOL	BCPE; DCPC; dichlorodiphenylethanol; Dimite; DMC; Qikron
CHLORFENVINPHOS	Apachlor; Birlane; Birlane 24; C 8949; CFV; CGA 26351; Compound 4072; Sapecron; SD 7859; Steladone; Supona; Vinylphate
CHLORMEQUAT CHLORIDE	CeCeCe; Chlormequat-chloride + cholin chloride; Cycocel; Cycocel-Extra; Cyogan; Hico CCC; Hormocel-2CCC; Increcel; Lihocin
CHLOROBENZILATE	Acaraben; Akar; Benzilan; Folbex smoke-strips; Kop-Mite
CHLORONITROPROPANE	Korax; Lanstan
CHLOROWAX 40	NCI-C53543
CHLOROWAX 500C	NCI-C53587
CHLOROXURON	C-1983; chloroxifenidim; Tenoran
CHLORPYRIFOS	Brodan; Dursban; Eradex; Lorsban; Pyrinex
CHLORPYRIFOS-METHYL	DOWCO 214; Ent 27520; Reldan
CLONITRALIDE	Bay 6067; Bayluscid; Niclosamide; SR73
COPPER-COUNT-N	
COPPER OXYCHLORIDE	BASF-Grunkupfer; basic copper chloride; Blitox; Chempar; Cobox; Coprantol; Cop Tox; Cupramar; Cupravit; Cuprokylt; Cuprosana; Cuprovinol; Cuprox; Devicopper; Fytolan; Kauritil; Kilex; Recop; Rhodiacuivre; Viricuivre; Vitigran
COPPER SULFATE	Bluestone; blue copperas; blue vitriol; Triangle
CORREX	
COUMAPHOS	Asuntol; Bay 21/199; Baymix; Co-Ral; Diolice; Meldane; Muscatox; Resistox
CROTOXYPHOS	Ciodrin; Ciovap; Cypona E.C.; Decrotox; Duo-Kill; Duravos; SD 4294
CRUFOMATE	Dowco 132; Ruelene
CRYOLITE	Kryocide

ED_005172C_00000035-00062

Table A4. *Continued.*

Common name	Alternate name
CUTRINE-PLUS	
CYANAZINE	Bladex; Fortral; SD 15418; WL 19805
CYANO (METHYLMERCURI) GUANIDINE	Morsodren; Panodrin A-13; Panogen Turf Fungicide
CYHEXATIN	Dowco 213; Plictran
CYPRAZINE	Outfox; S-9115
CYTROL AMITROLE-T	Amitril T.L.; Amitrol T
2,4-D ACID	Agrotect; Amoxone; Aqua-Kleen; BH 2,4-D; Chipco Turf Herbicide "D"; Chloroxone; Crop Rider; D50; Dacamine; Debroussaillant 600; Ded-Weed; Desormone; Dinoxol; DMA 4; Dormone, Emulsamine BK; Emulsamine E-3; Envert 171; Envert DT; Esteron Brush Killer; Esteron 99 Concentrate; Estone; Farmco; Fernesta; Fernimine; Fernoxone; Ferxone; Formula 40; Hedonal; Herbidal; Lawn-Keep; Macrondray; Miracle; Netagrone 600; Pennamine D; Planotox; Plantgard; Rhodia; Salvo; Spritz-Hormin/2,4-D; Spritz-Hormit/2,4-D; Super D Weedone; Superormone Concentre; Transamine; Tributon; U 46; U 46 D-Ester; U 46 D-Fluid; Visko-Rhap; Weedar; Weedatul; Weed-B-Gon; Weedone; Weed-Rhap; Weed Tox; Weedtrol
2,4 DB	Butoxon; Butoxone; Butoxone amine; Butoxone ester; Butyrac; Butyrac 200; Butyrac ester; Embutox; Embutox E
2,4-D BUTOXYETHANOL ESTER	Bladex-B; Brush Killer 64; 2,4-D-Bee; 2,4-D butoxyethyl ester; Planotox; Weedone LV4
2,4-D BUTYL ESTER	Butyl 2,4-D; butyl dichloro-phenoxyacetate; Esso Herbicide 10; Fernesta; Lironox; Shell 40
2,4-D DIMETHYLAMINE SALT	2,4-D amine; 2,4-D amine salt; Bladex G; Formula 40; Hormin; Phordene
2,4-D ISOOCTYL ESTER	
DALAPON	Basfapon/Basfapon N; BH Dalapon; Crisapon; Dalapon 85; Dalapon-Na; Ded-Weed; Devipon; Dowpon; Dowpon M; DPA; Gramevin; Revenge; Unipon
DCPA	chlorthal dimethyl; Dacthal
D-D SOIL FUMIGANT	Nemafene
DDD	Rhothane; TDE
DDE	
DDT	Anofex; Arkotine; Chlorophenothane; DDT technical; DDT 75% WDP; Dedelo; Didimac; Digmar; Genitox; Gyron; Hildit; Ixodex; Kopsol; Micro DDT 75; Neocid; Pentachlorin; R50; Rukseam; Zeidane; pp'Zeidane; Zerdane
DEET	Detamide; Metadelphene; MGK Diethyltoluamide; OFF
DEF	De-Green; E-Z-Off D; Fos-Fall "A"; Ortho Phosphate Defoliant
DEMETON	Bay 10756; demeton-O + demeton-S; E 1059; mercaptofos; Systemox; Systox
DIAZINON	Basudin; Dazzel; Diagran; Dianon; Diaterr-Fos; Diazajet; Diazatol; Diazide; Diazol; Dizinon; Dyzol; G-24480; Gardentox; Kayazinon; Kayazol; Neocidol; Nipsan; Sarolex; Spectracide
DICAMBA	Banex; Banvel 45; Banvel CST; Banvel D; Banvel Herbicide; Banvel II Herbicide; Banvel 4WS; CST; dianat
DICHLOBENIL	Casoron; Decabane
DICHLOFENTHION	dDichlofention; dichlorofenthion; Mobilawn; Tri-VC13; VC-13 Nemacide
DICHLONE	Phygon; Quintar
DICHLOROPROPENE	Telone II Soil Fumigant
DICHLORVOS	Apavap; Benfos; Cekusan; Cypona; DDVP; Dedevap; Devikol; Divipan; Duo-Kill; Duravos; Fly-Die; Fly Fighter; Herkol; Mafu; Marvex; Nogos; No-Pest; Nuvan; Oko; Phosvit; Tetravos, UDVF; Vapona; Vaponite; Vapora II; Verdican; Verdipor; Verdisol
DICLOFOP METHYL	Hoe-23408; Hoe-Grass; Hoelon; Hoelon 3 EC; Illoxan; Iloxan

BEST AVAILABLE COPY

ED_005172C_00000035-00063

Table A4. *Continued.*

Common name	Alternate name
DICOFOL	Acarin; Cekudifol; FW-293; Hifol; Kelthane; Mitigan
DICROTOPHOS	Bidrin; C 709; Carbicron; Diapadrin; Ektafos; SD 3562
DIELDRIN	Dieldrex; Dieldrine; Dieldrite; Octalox; Panoram D-31
DIFLUBENZURON	Dimilin; DU 112307; ENT-29054; OMS 1804; PDD 6040-I; PH 60-40; TH 6040
DILAN	
DIMETHOATE	AC-12880; Bi 58 EC; Cekuthoate; Cygon; Daphene; De-Fend; Demos-L40; Devigon; Dimate 267; Dimethoate technisch 95%; Dimethogen; Fosfamid; Fostion MM; Perfekthion; Rebelate; Rogodial; Rogor; Roxion; Trimetion
DIMETHRIN	Dimethrine
DIMETHYL FORMAMIDE	DMF; NCI-C60913; NSC 5356; U-4224
DIMETHYLSULFOXIDE	Demasorb; Demavet; Demeso; Dermasorb; dimethyl sulfoxide; dimethyl sulphoxide; DMS-70; DMS-90; DMSO; Doligur; Dromisol; Gamasol 90; Hyadur; Infiltrina; methyl sulfoxide; Somipront; SQ 9453; Syntexan
DINITRAMINE	Cobex; Cobexo; Dinitroamine; USB-3584
DINOBUTON	Acrex; Dessin; Dinofen; Drawinol; UC 19786
DINOCAP	Arathane; Crotothane; Iscothane; Karathane; Mildex
DINOSEB	Basanite; Caldon; Chemox General; Chemox PE; Chemsect DNBP; dinitro; Dinitro-3; Dinitro General; dinosebe; DNBP; Dynamyte; Elgetol 318; Gebutox; Hel-Fire; Kiloseb; Nitropone C; Premerge 3; Sinox General; Subitex; Unicrop DNBP; Vertac Dinitro Weed Killer 5; Vertac General Weed Killer; Vertac Selective Weed Killer
DIOXATHION	Delnav; Detic; Hercules AC528
DIPHENAMIDE	DIF 4; Dimid; diphenamid; Dymid; Enide; Enide 50; Fenam; L-34314; Lilly 34,314; U 4513; 80W
DIQUAT	Aquacide; Deiquat; Dextrone; reglon; Reglone; Reglox; Weedtrine-D
DISULFOTON	Bay 19639; Bay S275; Di-Syston; Disyston; dithiodemeton; dithiosystox; Ethylthiodemeton; Frumin AL; M-74; Solvirex; thiodemeton
DITHIANON	Delan; Thynon
DIURON	Cekiuron; Dailon; Diater; dichlorfenidim; Di-on; Direx 4L; Diurex; Diurol; Drexel Diuron 4L; Dynex; Farmco Diuron; Karmex; Unidron; Urox "D"; Vonduron
DNOC	Chemsect DNOC; DNC; Elgetol 30; Nitrador; Selinon; Sinox; Trifocide; Trifrina
DODINE	AC 5223; Apadodine; Carpene; Curitan; Cyprex; Doquadine; Melprex; Syllit; tsitrex; Venturol; Vondodine
DOWELL A170	corrosion inhibitor A170
DOWELL F75A	Ezeflo F75A surfactant
DOWELL F75N	Ezeflo F75N surfactant
DOWELL L47	Gypban L47 scale inhibitor
DOWELL M38N	silicate control additive M38W
DOWELL W35	emulsion and sludge preventor W35
DOWICIDE EC-7 (PENTACHLOROPHENOL)	88% pentachlorophenol; 12% other chlorophenols
*d*TRANS ALLETHRIN	Bioallethrin
ENDOSULFAN	Benzoepin thiodan; Beosit; Chlorthiepin; Crisulfan; Cyclodan; Devisulphan; Endocel; Endosol; EnSure; FMC 5462; Hildan; Hoe 2671; Insectophene; Kop Tiodan; Malix; N1A 5462; Thifor; Thimul; Thiodan; Thiofor; Thionex; Thiosulfan; Tiovel
ENDOTHALL	Accelerate; Aquathol; Aquathol K; Des-i-cate; Endothal; Endothal Turf Herbicide; Endothal Weed Killer; Herbicide 273; Hydout; Hydrothol; Hydrothol 47; Hydrothol 191
ENDRIN	Endrex; Hexadrin; Nendrin
EPN	
EPTC	Eptam
ETHANOL	Alcohol; ethyl alcohol

03

Table A4. *Continued.*

Common name	Alternate name
ETHION	*Diethion; Ethanox; Ethiol; Hylemox;* Rhodiacide; Rhodocide; Vegfru Fosmite
ETHOFUMESATE	NC 8438; Nortranese; Nortron; Tramat
ETHYLAN	Perthane
ETHYLBENZENE	EB; ethylbenzol; NCI-C56393; phenylethane
ETHYLENE DICHLORIDE	EDC
ETHYLENE GLYCOL	ethylene alcohol; ethylene dihydrate; glycol; glycol alcohol; Lutrol-9; Macrogol 400 BPC; M.E.G.; monoethylene glycol; NCI-C00920; Tescol
2-ETHYLHEXANOL	
FENAC	chlorfenac; Tri-Fen; Trifene
FENAMINOSULF	Bay 22555; Bayer 5072; Lesan
FENAZAFLOR	Fenozaflor
FENBUTATIN-OXIDE	SD 14114; Torque; Vendex
FENITROTHION	Accothion; Agrothion; Bay 41831; Bay S 5660; Cekutrothion; Cytel; Dybar; Fenitox; *Folithion; MEP; Novathion; Nuvanol; S 5660; Sumithion*
FENSON	CPBS; Fenizon; Murvesco; PCPBS
FENSULFOTHION	Bay 25141; Dasanit; S 767; Terracur P
FENTHION	Bay 29493; Baycid; Baytex; Entex; Lebaycid; mercaptophos; S 1752; Tiguvon
FENVALERATE	Belmark; Ectrin; Pydrin; S-5602; Sanmarton; SD 43775; Sumicidin; Sumifly; Sumipower
FLAMPROP-METHYL	Lancer; WL 29761
FLIT MLO	
FLUCHLORALIN	Basalin
FLUMETURON	C-2059; Cotoran; Cottonex; Lanex
FLUORODIFEN	C 6989; Preforan
FLURIDONE	Sonar
FOLPET	Folpan; Phaltan; Thiophal
FONOFOS	Dyfonate; N-2790
FOSAMINE AMMONIUM	Krenite brush control agent
FOSPIRATE	Dowco 217; ENT 27521; Torelle
GERANIOL	geraniol alcohol; geraniol extra; geranyl alcohol; Guaniol; Lemonol
GLENBAR	OCS-21944
GLYCEROL	glycerin; glycerine; glyceritol; glycyl alcohol; synthetic glycerin; 90 technical glycerine; trihydroxypropane
GLYODIN	Crag Fruit Fungicide 341
GLYPHOSATE	glyphosate isopropylamine salt; Roundup
HEPTACHLOR	Drinox H-34; Heptachlore; Heptamul; Heptox
HEPTACHLOR EPOXIDE	ENT 25,584; epoxyheptachlor; HCE; heptachlor epoxide; Velsicol 53-CS-17
HERCULES 7175	
HEXACHLOROBENZENE	Anticarie; Ceku C.B.; HCB; No Bunt
HEXACHLOROBUTADIENE	C-46; Dolen-Pur; GP-40-66:120; HCBD; hexachlorbutadiene; perchlorobutadiene
HEXAZINONE	Velpar weed killers
JODFENPHOS	C-9491; *iodofenphos;* Nuvanol N
KOMEEN	
LANDRIN	
LEAD ARSENATE	Gypsine; Soprabel; Talbot
LEPTOPHOS	MBCP; Phosvel
LETHANE 384	
LIME SULFUR	Security Lime Sulphur
LINDANE	Gamma BHC; Exagama; Forlin; Gallogama; Gamaphex; Gamma-col; Gammalin; Gammex; Gammexane; gamma HCH; Inexit; Isotox; Lindafor; Lindagam; Lindagrain; Lindagranox; Lindalo; Lindamul; Lindapoudre; Lindaterra; Lintox; Novigam; Silvanol

BEST AVAILABLE COPY

ED_005172C_00000035-00065

Table A4. *Continued.*

Common name	Alternate name
LINURON	Afalon; Hoe 2810; Linex 4L; Linorox; Linurex; Lorox; Sarclex
MALATHION	Calmathion; carbofos; Celthion; Cythion; Detmol MA 96%; Emmatos; Emmatos Extra; For-Mal; Fyfanon; Hilthion; Karbofos; Kop-Thion; Kypfos; Malamar; Malaphele; Malaspray; Malathion ULV Concentrate; Malatol; maldison; Malmed; Maltox; Mercaptothion; mercaptotion; MLT; Sumitox; Vegfru Malatox; Zithiol
MBC	
MCPB	Can-Trol; 2,4-MCPB; 2M-4Kh-M; PDQ; Thistrol; Tropotox
MERPHOS	Deleaf Defoliant; Easy Off-D; Folex
METAM-SODIUM	A7 Vapam; carbam; Karbation; Maposol; Metam-Fluid BASF; SMDC; Solasan 500; Sometam; Trimaton; Vapam; VPM
METHANOL	Alcohol
METHIDATHION	GS-13005; Somonil; Supracide; Ultracide
METHIOCARB	Bay 37344; Draza; H 321; mercaptodimethur; Mesurol; metmercapturon
METHOMYL	Lannate; Nu-Bait II; Nudrin; SD14999
METHOPRENE	ZR-515
METHOPROTRYNE	G 33393; Gesaran; Methoprotryn
METHOXYCHLOR	Chemform; Flo Pro Mc Seed Protectant; Marlate
METHYL DEMETON	Bay 15203; demeton methyl
METHYL PARATHION	Cekumethion; Devithion; Dimethyl Parathion; Drexel Methyl Parathion 4E; E601; Folidol M; Fosferno M50; Gearphos; Metacide; metafos; Metaphos; Nitrox 80; Parataf; Parathion-methyl; Paratox; Partron M; Penncap-M; Tekwaisa; Vertac Methyl Parathion technisch 80%; Wofatox
METHYL TRITHION	
METOLACHLOR	Bicep; CGA-24705; Codal; Cotoran Multi; Dual; Milocep; Ontrack 8E; Primagram; Primextra
METRIBUZIN	Bay 94337; Bay DIC 1468; Lexone; Sencor; Sencoral; Sencorex
MEVINPHOS	Apavinphos; Duraphos; Gesfid, Menite; OS-2046; Phosdrin; Phosfene
MEXACARBATE	Zectran
MIREX	Dechlorane; GC-1283
MITIN FF	
MNFA	FAM; Nissol
MOLINATE	Hydram; Ordram
MONOCROTOPHOS	Apadrin; Azodrin; Bilobran; C 1414; Crisodrin; Hazodrin; Monocil 40; Monocron; Nuvacron; Villardrin; Plantdrin; SD 9129; Susvin
MONOETHANOLAMINE	Colamine; ethanolamine; ethylolamine; glycinol; MEA; olamine; Thiofaco M-50; USAF EK-1597
MSMA	Ansar 170 H.C.; Ansar 529 H.C.; Arsonate Liquid; Bueno 6; Daconate 6; Dal-E-Rad; Herb-All; Merge 823; Mesamate; Monate; Target MSMA; Trans-Vert; Weed-E-Rad; Weed-Hoe
NALED	Bromchlophos; Bromex; Dibrom; RE 4355
N-BUTANOL	1-butanol; butan-1-ol; butyl hydroxide; butyric or normal primary butyl alcohol; CCS 203; 1-hydroxybutane; methylolpropane; n-butyl alcohol; propylcarbinol; propylmethanol
NEODOL 25-9	
NEODOL 25-12	
NITRALIN	Planavin
NITRAPYRIN	Dowco 163; N-Serve
NITROFEN	nitrofene; nitrophen; Nip; Tok; Tok E-25; Tok WP-50; Trizilin
NOREA	Herban; Hercules 7531; noruron
ORTHO 11775	

ED_005172C_00000035-00066

Table A4. *Continued.*

Common name	Alternate name
ORYZALIN	Dirimal; Ryzelan; Surflan
OVEX	chlorofenizon; Chlorfenson; Corotran; CPCBS; difenson; ephirsulphonate; Estonmite; Niagaratran; ovatran; Ovochlor; Ovotran; Sappiran
OXAMYL	DPX 1410; Vydate L insecticide/nematicide
OXYDEMETON-METHYL	Bay 21097; demeton-S-methyl sulfoxid; metasystemox; Metasystox-R; metilmercaptofosoksid
OXYTHIOQUINOX	Bay 36205; chinomethionat; chinomethionate; Morestan; quinomethionate; SS 2074
PARA-DICHLOROBENZENE	Paracide; PDB
PARAQUAT	Cekuquat; Crisquat; Dextrone; Dexuron; Esgram; Gramonol; Gramoxone; Gramuron; Herbaxon; Herboxone; Paracol; Pathclear; Pillarquat; Pillarxone; Sweep; Terraklene; Totacol; Toxer Total; Weedol
PARATHION	AC3422; Alkron; Alleron; Aphamite; Bladan; Corothion; Drexel parathion 8E; E-605; ENT 15108; Ethyl Parathion; Etilon; Folidol E-605; Fosferno 50; Niran; Orthophos; Panthion; Paramar; Paraphos; Parathene; Parawet; Phoskil; Rhodiatox; Soprathion; Stathion; Thiophos
PEBULATE	PEBC; R-2061; Tillam
PENTACHLOROPHENATE SODIUM	Santobrite; sodium pentachlorophenate; Weedbeads
PENTACHLOROPHENOL	Dow Pentachlorophenol; DP-2 Antimicrobial; PCP; penchlorol; penta; Pentacon; Penwar; Priltox; Santobrite; Santophen; Sinituho; Weedone
PERMETHRIN	Ambush; BW-21-Z; Ectiban; FMC-33297; Indothrin; Kafil; Permasect; Perthrine; Pounce; PP 557
PHORATE	AC 3911; Phorate-10G; Rampart; Thimet; Timet; Vegfru Foratox
PHORAZETIM	Bay 38819; Gophacide
PHOSALONE	Azofene; RP 11974; Rubitox; Zolone
PHOSMET	Appa; Imidan; Kemolate; phosmet; PMP; Prolate R-1504
PHOSPHAMIDON	Apamidon; C 570; Dimecron; Dixon
PHOTO-DIELDRIN	NCI-C00599
PHOXIM	Baythion; Bay 77488; phoxime; Volator
PICLORAM	Amdon; Grazon; Tordon
PIPERALIN	Pipron
PIPERONYL BUTOXIDE	Butacide; Piperonyl Butoxide technical; Prentox Piperonyl Butoxide technical
PROFENOFOS	CGA-15324; Curacron; Polycron; Selecron
PROPACHLOR	
PROPANIL	Bay 30130; Chem Rice; DPA; Farmco Propanil; FW-734; Herbox technical; Propanex; Prop-Job; Riselect; Stam F-34; Stam M-4; Stam Supernox; Stampede 3E; Strel; Supernox; Surcopur; Surpur; S 10165; Vertac
PROPHAM	Ban-Hoe; Beet-Kleen; Chem-Hoe; IFC; IPC; Premalox; Triherbide-IPC; Tuberite
PROPOXUR	Aprocarb; Bay 39007; Baygon; Blattenex; Propyon; Suncide; Tugon Fliegenkugel; Unden
PROPYL ISOME	N-Propyl isome
PROPYLENE GLYCOL	Methyl glycol; Methylethylene glycol; monopropylene glycol; PG 12; Sirlene; trimethyl glycol
PYRETHRUM	Pyrethrins
RESMETHRIN	Benzofuroline; Chrysron; Premgard; Pynosect; Pyretherm; Synthrin
RONNEL	Ectoral; Etrolene; fenchlorfos; fenchlorphos; Nankorl; Korlan; Trolene; Viozene
ROTENONE	Chem Fish; Derris; Nicouline; Prentox; Tubatoxin
ROWMATE	Sirmate; UC 22463
RYANIA	Bonide Ryatox; ground Ryania specisa (Vahl) stemwood (alkoloid ryanodine); Ryanexel; Ryania powder; Ryania speciosa; Ryanicide; Ryanodine

BEST AVAILABLE COPY

ED_005172C_00000035-00067

Table A4. *Continued.*

Common name	Alternate name
S-BIOALLETHRIN	Esbiol; Esbiol Concentrate 90%
SD 7438	ENT 25,739; Shell SD 7,438
SD 8339	Accel
SD 16898	
SD 17250	
SILVEX ACID	Amchem 2,4,5-TP; Aqua-Vex; Double Strength; fenoprop; Fruitone T; Kuron; Kurosal; Silvex; Silvi-Rhap; Weed-B-Gon
SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER	Kuron
SIMAZINE	Aquazine; Cekusan; Farmco Simazine; Framed; G-27692; Gesatop; Primatol S; Princep; Simadex; Simanex; Sim-trol
SODIUM ARSENITE	
SODIUM SELENITE	disodium selenite
2,4,5-T BUTOXYETHANOL ESTER	Bladex H; Hormoslyr 500T; Trinoxol
TEMEPHOS	Abate; Abathion; Difenthos; Ecopro; Nimitox
TEPA	Aphoxide; APO
TEPP	Tetron; Vapotone
TERBUFOS	Counter 15G Soil Insecticide-Nematicide
TERBUTRYN	GS 14260; Igran; Prebane; Terbutrex
TERPENE POLYCHLORINATES	Strobane
2,3,4,6-TETRACHLOROPHENOL	Dowicide 6; tetrachlorophenol
TETRACHLORVINPHOS	Appex; CVMP; Debantic; Dust M; Gardcide; Gardona; Rabon; Rabond; ROL; SD 8447; Stirofos
TETRADIFON	Nia 5488; Tedion; Tedion V-18
TETRAMINE	TEM
TETRASUL	Animert V-101
TFM	Hoe 02770; Lamprecid; Lamprecide; TFN
THANITE	
TOLUENE	Methacide; methylbenzene; methylbenzol; NC1-CO7272; phenylmethane; Toluol
TOXAPHENE	Attac 4-2; Attac 4-4; Attac 6; Attac 6-3; Attac 8; camphechlor; Camphoclor; Camphofene Huileux; Motox; Phenacide; Phenatox; Polychlorocamphene; Strobane T-90; Toxakil; Toxon 63; Vertac Toxaphene 90
TRANID	UC 2047A
TREFMID	
TRIALLATE	Avadex BW; Far-Go
TRICHLORFON	Bovinox; Briten; Cekufon; Ciclosom; clorofos; Crinex; Danex; dipterex; Dylox; Equino-Aid; Leivasom; Neguvon; Proxol; Trichlorphon; Trinex; Tugon
2,3,6-TRICHLOROBENZOIC ACID	Benzac; HC-1281; 2,3,6-TBA; TCBA; Tribac; trichlorobenzoic acid
TRICHLORONATE	Agrisil; Agritox; Bay 37289; fenophosphon; Phytosol; S4400; trichloronat
TRICLOPYR	Garlon 3A; Garlon 4
TRICRESYL PHOSPHATE	Celluflex 179C; cresyl phosphate; Disflamoll TKP; Fyrquel 150; IMOL S 140; Kronitex; Lindol; NCI-C61041; tris (tolyloxy) phosphine oxide; tritolyl phosphate
TRIFLURALIN	Crisalina; Digermin; Elancolan; Farmco Trifluralin; Ipersan; Sinflouran; Trefanocide; Treflan; Trifluraline; Triflurex
TRI-N-BUTYL PHOSPHATE	Celluphos 4; TBP; tributyl phosphate

ED_005172C_00000035-00068

Table A4. *Continued:*

Common name	Alternate name
TRIPHENYL PHOSPHATE	Celluflex TPP; TPP
TRIPHENYLTIN HYDROXIDE	Du-Ter; Duter; fentin hydroxide; Haitin; Suzu H; TPTH; TPTOH; Tubotin
UC 10854	Hercules AC 5727
UREABOR	
VERNOLATE	Surpass; Vernam
WARFARIN	Co-Rax; Coumafene; Cov-R-Tox; Kypfarin; Liqua-Tox; Mouse Pak; Rat Pak; Ratox; RAX; Rodex; Rodex Blox; Rutoxin; Tox-Hid; Warfarin Q; Zoocoumarin
XYLENE	Xylol
ZINC SULFATE	NU-Z

BEST AVAILABLE COPY

ED_005172C_00000035-00069

Table A5. *Acute toxicity data for 410 chemicals and 66 species of freshwater animals from the Columbia National Fisheries Research Laboratory, U.S. Fish and Wildlife Service, 1965–1984.*

Contents

Table page	Chemicals	No. of tests	Table page	Chemicals	No. of tests
1	ACEPHATE	66	65	CAPTAN	15
4	ACETONE	1	66	CARBARYL	136
5	AEROFROTH 71	2	72	CARBOFURAN	15
6	AERO XANTHATE 343	2	73	CARBOPHENOTHION	6
7	AKTON	5	74	CCA TYPE III	5
8	ALACHLOR	8	75	CHLORAMBEN	2
9	ALDICARB	4	76	CHLORBROMURON	2
10	ALDRIN	26	77	CHLORDANE	56
12	ALLETHRIN	6	80	CHLORDANE CIS	1
13	ALUMINUM NITRATE	1	81	CHLORDANE-HCS-3260	8
14	ALUMINUM SULFATE	1	82	CHLORDANE TRANS	1
15	AMDRO	6	83	CHLORDECONE	14
16	AMETRYN	2	84	CHLORDIMEFORM	2
17	AMINOCARB	144	85	CHLORENDATE DIBUTYL	7
23	AMITROLE	5	86	CHLORENDATE DIMETHYL	9
24	ANILAZINE	5	87	CHLORFENETHOL	4
25	ANTIMYCIN A	108	88	CHLORFENVINPHOS	5
30	APHOLATE	2	89	CHLORINATED ALPHA OLEFIN 22/44	2
31	AQUA COP	3	90	CHLORMEQUAT CHLORIDE	4
32	AQUA-VATOR	4	91	CHLOROBENZILATE	3
33	ARAMITE	6	92	CHLORONITROPROPANE	3
34	ATRAZINE 4L	2	93	6-CHLORO-2-PICOLINIC ACID	6
35	AZIDE POTASSIUM	6	94	CHLOROWAX LV	2
36	AZIDE SODIUM	6	95	CHLOROWAX 40	2
37	AZINPHOS-ETHYL	6	96	CHLOROWAX 50	2
38	AZINPHOS-METHYL	77	97	CHLOROWAX 70	2
42	BACILLUS THURINGIENSIS	3	98	CHLOROWAX 500C	13
43	BARBAN	2	99	CHLOROXURON	2
44	BENEFIN	3	100	CHLORPYRIFOS	23
45	BENOMYL	41	101	CHLORPYRIFOS-METHYL	5
47	BENSULIDE	3	102	CLARIFITE	1
48	BENTHIOCARB	6	103	CLONITRALIDE	9
49	BENZALKONIUM CHLORIDE	2	104	COPPER-COUNT-N	3
50	BENZENE	11	105	COPPER-COUNT-NS	3
51	BENZENE HEXACHLORIDE	12	106	COPPER OXYCHLORIDE	1
52	BENZOTHIAZOLE 2-MERCAPTO	6	107	COPPER SULFATE	7
53	BENZOTHIAZOLE 2-METHYL MERCAPTO	9	108	CORREX	7
54	BENZOYLPROP ETHYL	9	109	COUMAPHOS	9
55	BENZYL BENZOATE	1	110	CROTOXYPHOS	8
56	BINAPACRYL	5	111	CRUFOMATE	4
57	BLACK CRUDE OIL	5	112	CRYOLITE	4
58	BOMYL	3	113	CUMYLPHENYL DIPHENYL PHOSPHATE	3
59	BUFENCARB	4	114	CUTRINE-PLUS	5
60	BUTOXY POLYPROPYLENE GLYCOL	1	115	CYANAZINE	11
61	BUTYLATE	6	116	CYANO (METHYLMERCURI) GUANIDINE	1
62	CACODYLIC ACID	4	117	CYHEXATIN	8
63	CALCIUM HYPOCHLORITE	2	118	CYPRAZINE	4
64	CAPTAFOL	9			

BEST AVAILABLE COPY

ED_005172C_00000035-00070

Table A5. *Continued.*

Contents

Table page	Chemicals	No. of tests
119	CYTROL AMITROLE-T	6
120	2,4-D	21
121	2,4-D BUTOXYETHANOL ESTER	18
122	2,4-D BUTYL ESTER	33
124	2,4-D DIMETHYLAMINE SALT	51
127	2,4-D DODECYL/TETRADODECYL AMINE SALT	39
129	2,4-D ISOOCTYL ESTER	1
130	2,4-D OLEYLPROPYLENEDIAMINE SALT	2
131	2,4-D PROPYLENE GLYCOL BUTYL ETHER ESTER	45
133	2,4-D/2,4,5-T 24%/28%	4
134	2,4-D/2,4,5-T 30%/28%	2
135	2,4-D/2,4,5-T 34%/17%	2
136	DALAPON	4
137	2,4-DB	6
138	DCPA	9
139	D-D SOIL FUMIGANT	7
140	DDD	21
141	DDE	3
142	DDT	85
146	DEAD-X	6
147	DEET	1
148	DEF	31
150	DEMETON	10
151	DIAZINON	10
152	DICAMBA	7
153	DICHLOBENIL	15
154	DICHLOFENTHION	11
155	DICHLONE	5
156	2,6-DICHLOROBENZOIC ACID	2
157	DICHLOROPHENYL METHYLSULFONATE	2
158	DICHLOROPROPENE	5
159	DICHLORVOS	16
160	DICLOFOP METHYL	2
161	DICOFOL	8
162	DICROTOPHOS	8
163	DIELDRIN	37
165	DIFLUBENZURON	42
167	DILAN	4
168	DIMETHOATE	5
169	DIMETHRIN	18
170	DIMETHYL FORMAMIDE	1
171	DIMETHYLSULFOXIDE	2
172	DINITRAMINE	16
173	DINOBUTON	2
174	DINOCAP	6
175	DINOSEB	24
176	2,6-DIOCTADECYL-P-CRESOL	1
177	DIOXATHION	8
178	DIPHENAMIDE	9
179	DIQUAT	18
180	DISULFOTON	9
181	DITHIANON	3
182	DIURON	59
185	DNOC	5
186	DODINE	1
187	DOWELL A170	1
188	DOWELL F75A	1
189	DOWELL F75N	1
190	DOWELL L47	1
191	DOWELL M38N	6
192	DOWELL W35	1
193	DOWICIL 75	1
194	*D*-TRANS ALLETHRIN	29
196	EMCOL AD-410	3
197	ENDOSULFAN	10
198	ENDOTHALL	3
199	ENDOTHALL AQUATHOL K	14
200	ENDOTHALL COPPER SALT	4
201	ENDOTHALL DES-I-CATE	1
202	ENDOTHALL HERBICIDE 282	2
203	ENDOTHALL HYDROTHOL 47	1
204	ENDOTHALL HYDROTHOL 191	9
205	ENDOTHALL POTASSIUM SALT	3
206	ENDRIN	53
209	EPN	11
210	EPTC	25
212	ETHANOL	2
213	ETHION	13
214	ETHOFUMESATE	4
215	ETHYLAN	2
216	ETHYLBENZENE	12
217	ETHYLENE DICHLORIDE	3
218	ETHYLENE GLYCOL	3
219	2-ETHYLHEXANOL	2
220	FENAC	10
221	FENAMINOSULF	5
222	FENAZAFLOR	3
223	FENBUTATIN-OXIDE	6
224	FENITROTHION	160
231	FENSON	2
232	FENSULFOTHION	2
233	FENTHION	43
235	FENVALERATE	29
237	FIRE-TROL 100	12
238	FIRE-TROL 931	12
239	FLAMPROP-METHYL	7
240	FLIT MLO	2
241	FLUCHLORALIN	11

ED_005172C_00000035-00071

Table A5. *Continued.*

Contents

Table page	Chemicals	No. of tests	Table page	Chemicals	No. of tests
242	FLUOMETURON	12	302	METHOXYCHLOR	146
243	FLUORENE	3	309	METHYL DEMETON	2
244	FLUORODIFEN	2	310	METHYL PARATHION	31
245	FLURIDONE	36	312	METHYL TRITHION	9
247	FOLPET	14	313	METOLACHLOR	6
248	FONOFOS	2	314	METRIBUZIN	5
249	FOSAMINE AMMONIUM	7	315	MEVINPHOS	16
250	FOSPIRATE	3	316	MEXACARBATE	90
251	FYRQUEL GT	23	320	MIREX	13
252	GERANIOL	4	321	MITIN FF	2
253	GLENBAR	1	322	MNFA	1
254	GLYCEROL	1	323	MOLINATE	9
255	GLYODIN	1	324	MON 0818	11
256	GLYPHOSATE	51	325	MONOCROTOPHOS	5
259	GREEN CRUDE OIL	10	326	MONOETHANOLAMINE	3
260	HALOWAX 1099	2	327	MSMA	8
261	HEPTACHLOR	31	328	NALED	18
263	HEPTACHLOR EPOXIDE	2	329	N-BUTANOL	1
264	HERCULES 7175	1	330	NEODOL 25-9	4
265	HEXACHLOROBENZENE	12	331	NEODOL 25-12	3
266	HEXACHLOROBUTADIENE	3	332	NITRALIN	4
267	HEXAZINONE	5	333	NITRAPYRIN	7
268	HOUGHTO-SAFE 520	9	334	NITROFEN	1
269	HOUGHTO-SAFE 1120	15	335	N,N-DIMETHYL-2,4-DICHLOROPHENOXYACETAMIDE	1
270	2-HYDROXY-4,4'-DICHLORODIPHENYL ETHER	1	336	NONYLPHENYL DIPHENYL PHOSPHATE	2
271	HYDROXY-S-TRIAZINYL ALANINE	1	337	NOREA	4
272	2-HYDROXY-2',4,4'-TRICHLORODIPHENYL ETHER	1	338	ORTHO 11775	7
273	JODFENPHOS	2	339	ORYZALIN	2
274	KOMEEN	2	340	OVEX	5
275	KRONITEX 200	21	341	OXAMYL	10
276	LANDRIN	2	342	OXYDEMETON-METHYL	9
277	LEAD ARSENATE	1	343	OXYTHIOQUINOX	2
278	LEPTOPHOS	37	344	PARA-DICHLOROBENZENE	1
280	LETHANE 384	6	345	PARAQUAT	9
281	LIME SULFUR	5	346	PARATHION	47
282	LINDANE	45	348	PARATHION DITHIOATE ANALOGUE	3
284	LINURON	6	349	PAROIL 160	4
285	MALATHION	70	350	PAROIL 1032	5
288	MBC	10	351	PAROIL 1048	4
289	MCPA DIMETHYLAMINE SALT	1	352	PCB AROCLOR 1016	17
290	MCPB	3	353	PCB AROCLOR 1221	1
291	MERPHOS	14	354	PCB AROCLOR 1232	1
292	METAM-SODIUM	1	355	PCB AROCLOR 1242	9
293	METHANOL	1	356	PCB AROCLOR 1248	7
294	METHIDATHION	2	357	PCB AROCLOR 1254	12
295	METHIOCARB	3	358	PCB AROCLOR 1260	6
296	METHOMYL	84	359	PCB AROCLOR 1262	1
300	METHOPRENE	7	360	PCB AROCLOR 1268	1
301	METHOPROTRYNE	1			

ED_005172C_00000035-00072

Table A5. *Continued.*

Contents

No. of tests	Table page	Chemicals	No. of tests
146	361	PCB AROCLOR 4465	1
2	362	PCB AROCLOR 5442	1
31	363	PCB AROCLOR 5460	1
9	364	PCB 2-CHLOROBIPHENYL	1
6	365	PCB 3-CHLOROBIPHENYL	1
5	366	PCB 4-CHLOROBIPHENYL	1
16	367	PCB 2,4-DICHLOROBIPHENYL	1
90	368	PCB 4,4'-DICHLOROBIPHENYL	1
13	369	PCB 2,2',4,4',6,6'-HEXACHLOROBIPHENYL	1
2	370	PCB 2,2',4,5,5'-PENTACHLOROBIPHENYL	1
1	371	PCB 2,3,4'-TRICHLOROBIPHENYL	1
9	372	PEBULATE	1
11	373	PENTACHLOROPHENOL	15
5	374	PENTACHLOROPHENOL COPPER SALT	6
3	375	PENTACHLOROPHENOL (DOWICIDE EC-7)	2
8	376	PENTACHLOROPHENOL SODIUM SALT	18
18	377	PERMETHRIN	29
1	379	PHORATE	18
4	380	PHORAZETIM	1
3	381	PHOSALONE	3
4	382	PHOS-CHECK 202	12
7	383	PHOS-CHECK 259	12
1	384	PHOSFLEX 31P	23
	385	PHOSMET	74
1	389	PHOSPHAMIDON	11
2	390	PHOTO-DIELDRIN	3
5	391	PHOXIM	42
10	393	PHTHALATE DIBUTYL	18
9	394	PHTHALATE DI-2-ETHYLHEXYL	9
2	395	PHTHALATE 2-ETHYLHEXYL	1
1	396	PHTHALIC ACID	1
9	397	PICLORAM	70
47	400	PIPERALIN	1
3	401	PIPERONYL BUTOXIDE	7
4	402	POTASSIUM NAPHTHALENEACETATE	1
5	403	PROFENOFOS	12
4	404	PROPACHLOR	6
17	405	PROPANIL	3
1	406	PROPHAM	5
1	407	PROPOXUR	5
9	408	PROPYLENE GLYCOL	1
7	409	PROPYL ISOME	1
12	410	PURIFLOC C-3	8
6	411	PYDRAUL 50E	28
1	413	PYDRAUL 115E	7
1	414	PYRETHRUM	41
	416	RESMETHRIN	23
	417	RONNEL	12
	418	ROTENONE	7
	419	ROWMATE	1
	420	RU-11679	22
	421	RYANIA	4
	422	RYLEX D (2-HYDROXY-4-DODECOXYBENZOPHENONE)	1
	423	RYLEX H (2,4-DIHYDROBENZOPHENONE)	1
	424	RYLEX NBC (NICKEL DIBUTYLDITHIO CARBAMATE)	1
	425	SANTICIZER 148	15
	426	SANTICIZER 154	22
	427	S-BIOALLETHRIN	15
	428	SD 7438	2
	429	SD 8339	2
	430	SD 16898	5
	431	SD 17250	4
	432	SILVEX	12
	433	SILVEX BUTOXYETHANOL ESTER	8
	434	SILVEX POTASSIUM SALT	2
	435	SILVEX PROPYLENE GLYCOL BUTYL ETHER ESTER	12
	436	SIMAZINE	11
	437	SODIUM ARSENITE	5
	438	SODIUM SELENITE	6
	439	SRCII SYNFUEL	2
	440	2,4,5-T BUTOXYETHANOL ESTER	4
	441	2,4,5-T TRIETHYLAMINE SALT	4
	442	TEMEPHOS	68
	445	TEPA	3
	446	TEPP	6
	447	TERBUFOS	26
	449	TERBUTRYN	2
	450	TERPENE POLYCHLORINATES	4
	451	2,3,4,6-TETRACHLOROPHENOL	5
	452	TETRACHLORVINPHOS	4
	453	TETRADIFON	5
	454	TETRAMINE	3
	455	TETRASUL	1
	456	TFM	5
	457	TH 285-N	2
	458	TH 336-N	2
	459	THANITE	7
	460	TOLUENE	11
	461	TOXAPHENE	89
	465	TRANID	2
	466	TREFMID	3
	467	TRETOLITE J-146	2

ED_005172C_00000035-00073

Table A5. *Continued.*

Contents

Table page	*Chemicals*	*No. of tests*	*Table page*	*Chemicals*	*No. of tests*
468	TRETOLITE JN-9045	1	493	TRIPHENYLTIN HYDROXIDE	5
469	TRETOLITE JW-8226	7	494	TRITON B-1956	1
470	TRIALLATE	7	495	UC 10854	2
471	TRICHLORFON	229	496	UREABOR	13
481	2,3,6-TRICHLOROBENZOIC ACID	1	497	VERNOLATE	8
482	TRICHLOROMANDELIC ACID	1	498	WARFARIN	1
483	TRICHLORONATE	3	499	WATER GLYCOLS 894-44A	9
484	TRICLOPYR	2	500	WATER GLYCOLS 894-44B	9
485	TRICRESYL PHOSPHATE	6	501	WATER GLYCOLS 894-44C	9
486	TRIFLURALIN	62	502	XYLENE	16
489	2,3,5-TRIMETHYL NAPHTHALENE	1	503	XYLENOL AMINO	2
490	2,3,6-TRIMETHYL NAPHTHALENE	1	504	XYLENOL DIMETHYLAMINO	8
491	TRI-N-BUTYL PHOSPHATE	1	505	Z-200	2
492	TRIPHENYL PHOSPHATE	3	506	ZINC SULFATE	6

ED_005172C_00000035-00074

ACEPHATE

CAS # 30560-19-1 PRINCIPAL USE: INSECTICIDE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	94.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	7.4	40
2	94.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	7.4	320
3	94.00% TECHNICAL MATERIAL	ISOGENUS SP.	1ST YEAR CLASS	.	7.0	7.0	35
4	75.00% WETTABLE POWDER	ISOGENUS SP.	1ST YEAR CLASS	.	7.0	7.5	42
5	94.00% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	.	12.0	7.5	38
6	94.00% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	.	12.0	6.5	40
7	94.00% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	.	12.0	8.5	38
8	94.00% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	17.0	7.4	40
9	94.00% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	17.0	7.4	320
10	75.00% WETTABLE POWDER	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	20.0	7.2	40
11	75.00% WETTABLE POWDER	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	20.0	7.2	40
12	75.00% WETTABLE POWDER	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	20.0	7.2	40
13	75.00% WETTABLE POWDER	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	20.0	7.2	40
14	94.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.80	12.0	7.8	42
15	94.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.90	7.0	7.5	42
16	94.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00	12.0	6.5	42
17	94.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.70	12.0	7.5	42
18	94.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.90	12.0	8.5	42
19	94.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00	12.0	7.8	330
20	75.00% WETTABLE POWDER	CUTTHROAT TROUT		0.90	12.0	7.5	42
21	94.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	10.0	7.4	40
22	94.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	12.0	6.5	40
23	94.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	12.0	8.5	40
24	94.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.90	12.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC		.	.	>50.	.	.		.	.
2	STATIC	LC		.	.	>50.	.	.		.	.
3	STATIC	LC	65.0	33.900	12[illegible].000		.	.	11.7	8.700	15.800
4	STATIC	LC	>50	.	.		.	.	12.0	8.000	17.900
5	STATIC	LC	>14.	.	.		.	.	9.5	7.300	12.300
6	STATIC	LC	29.3	22.800	37.600		.	.	6.4	5.300	7.800
7	STATIC	LC	>40.	.	.		.	.	21.2	15.600	28.200
8	STATIC	EC		.	.	>50.	.	.		.	.
9	STATIC	EC		.	.	>50.	.	.		.	.
10	0 DAY DEGRA	EC		.	.	>1000	.	.		.	.
11	1 DAY DEGRA	EC		.	.	>1000	.	.		.	.
12	3 DAY DEGRA	EC		.	.	>1000	.	.		.	.
13	7 DAY DEGRA	EC		.	.	>1000	.	.		.	.
14	STATIC	LC	>50.	.	.		.	.	>50.	.	.
15	STATIC	LC	>100.	.	.		.	.	>100.	.	.
16	STATIC	LC	>50.	.	.		.	.	>50.	.	.
17	STATIC	LC	>100.	.	.		.	.	>100.	.	.
18	STATIC	LC	>60.	.	.		.	.	>60.	.	.
19	STATIC	LC	>50.	.	.		.	.	>50.	.	.
20	STATIC	LC	>100.	.	.		.	.	>100.	.	.
21	STATIC	LC	>1000.	.	.		.	.	1100.	775.000	1561.000
22	STATIC	LC	>50.	.	.		.	.	>50.	.	.
23	STATIC	LC	>50.	.	.		.	.	>50.	.	.
24	STATIC	LC	>50.	.	.		.	.	>50.	.	.

ED_005172C_00000035-00075

ACEPHATE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
25	94.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.90	12.0	7.4	320
26	75.00% WETTABLE POWDER	RAINBOW TROUT		1.20	10.0	7.4	40
27	97.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.20	7.0	7.4	40
28	97.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	17.0	7.4	40
29	97.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.20	17.0	7.4	40
30	97.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	7.0	7.5	40
31	97.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	17.0	7.5	40
32	97.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	7.5	40
33	97.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	7.5	12
34	97.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	6.5	40
35	97.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	7.5	40
36	97.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	8.5	40
37	97.00% TECHNICAL MATERIAL	ATLANTIC SALMON	YOLK-SAC FRY	.	7.0	7.5	40
38	94.00% TECHNICAL MATERIAL	BROOK TROUT		1.50	12.0	7.5	42
39	75.00% WETTABLE POWDER	BROOK TROUT		0.70	7.0	7.5	42
40	75.00% WETTABLE POWDER	BROOK TROUT		0.70	17.0	7.5	42
41	75.00% WETTABLE POWDER	BROOK TROUT		0.20	12.0	6.5	42
42	75.00% WETTABLE POWDER	BROOK TROUT		0.20	12.0	7.5	42
43	75.00% WETTABLE POWDER	BROOK TROUT		0.20	12.0	9.0	42
44	75.00% WETTABLE POWDER	BROOK TROUT		0.20	12.0	8.0	12
45	75.00% WETTABLE POWDER	BROOK TROUT		0.20	12.0	8.0	44
46	75.00% WETTABLE POWDER	BROOK TROUT		0.20	12.0	8.0	300
47	75.00% WETTABLE POWDER	BROOK TROUT		1.00	7.0	7.5	40
48	75.00% WETTABLE POWDER	BROOK TROUT		1.00	17.0	7.5	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
25	STATIC	LC	>50.	.	.		.	.	>50.	.	.
26	STATIC	LC	>1000.	.	.		.	.	730.	580.000	920.000
27	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
28	STATIC	LC	>100.	.	.		.	.	110.	63.000	190.000
29	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
30	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
31	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
32	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
33	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
34	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
35	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
36	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
37	STATIC	LC	>50.	.	.	>50.	.	.	>50.	.	.
38	STATIC	LC	>50.	.	.		.	.	>50.	.	.
39	STATIC	LC	>100.	.	.		.	.	>100.	.	.
40	STATIC	LC	>100.	.	.		.	.	>100.	.	.
41	STATIC	LC	>100.	.	.		.	.	>100.	.	.
42	STATIC	LC	>100.	.	.		.	.	>100.	.	.
43	STATIC	LC	>100.	.	.		.	.	>100.	.	.
44	STATIC	LC	>100.	.	.		.	.	>100.	.	.
45	STATIC	LC	>100.	.	.		.	.	>100.	.	.
46	STATIC	LC	>100.	.	.		.	.	>100.	.	.
47	STATIC	LC	>100.	.	.	>100.	.	.	>100.	.	.
48	STATIC	LC	>100.	.	.	>100.	.	.	>100.	.	.

ED_005172C_00000035-00076

45	STATIC	LC	>100.	.	.		>100.	.	.	>100.	.	.
46	STATIC	LC	>100.	.	.		>100.	.	.	>100.	.	.
47	STATIC	LC	>100.	.	.		>100.	.	.			
48	STATIC	LC	>100.	.	.							

ACEPHATE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
49	75.00% WETTABLE POWDER	FATHEAD MINNOW		1.00	20.0	7.4	40
50	75.00% WETTABLE POWDER	FATHEAD MINNOW		1.00	20.0	7.4	40
51	94.00% TECHNICAL MATERIAL	CHANNEL CATFISH		2.00	22.0	7.4	40
52	75.00% WETTABLE POWDER	CHANNEL CATFISH		0.50	22.0	7.5	40
53	94.00% TECHNICAL MATERIAL	BLUEGILL		0.40	20.0	7.4	44
54	94.00% TECHNICAL MATERIAL	BLUEGILL		0.40	12.0	7.4	44
55	94.00% TECHNICAL MATERIAL	BLUEGILL		0.80	17.0	6.5	40
56	94.00% TECHNICAL MATERIAL	BLUEGILL		0.80	17.0	9.5	40
57	94.00% TECHNICAL MATERIAL	BLUEGILL		0.60	17.0	7.4	40
58	94.00% TECHNICAL MATERIAL	BLUEGILL		0.60	17.0	7.4	320
59	75.00% WETTABLE POWDER	BLUEGILL		0.40	20.0	7.4	40
60	94.00% TECHNICAL MATERIAL	YELLOW PERCH		2.00	12.0	7.5	42
61	75.00% WETTABLE POWDER	YELLOW PERCH		1.80	12.0	7.5	42
62	75.00% WETTABLE POWDER	YELLOW PERCH		1.80	12.0	6.5	42
63	75.00% WETTABLE POWDER	YELLOW PERCH		1.80	12.0	9.0	42
64	75.00% WETTABLE POWDER	YELLOW PERCH		1.60	12.0	8.0	12
65	75.00% WETTABLE POWDER	YELLOW PERCH		1.80	12.0	8.0	44
66	75.00% WETTABLE POWDER	YELLOW PERCH		1.80	12.0	8.0	300

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
49	STATIC	LC	>1000.	.	.		.	.	>1000.	.	.
50	STATIC	LC	>1000.	.	.		.	.	>1000.	.	.
51	STATIC	LC	>1000.	.	.		.	.	>1000.	.	.
52	STATIC	LC		560.000	1000.000		.	.		560.000	1000.000
53	STATIC	LC	>1000.	.	.		.	.	>1000.	.	.
54	STATIC	LC	>50.	.	.		.	.	>50.	.	.
55	STATIC	LC	>50.	.	.		.	.	>50.	.	.
56	STATIC	LC	>50.	.	.		.	.	>50.	.	.
57	STATIC	LC	>50.	.	.		.	.	>50.	.	.
58	STATIC	LC	>50.	.	.		.	.	>50.	.	.
59	STATIC	LC	>200.	.	.		.	.	>200.	.	.
60	STATIC	LC	>50.	.	.		.	.	>50.	.	.
61	STATIC	LC	>100.	.	.		.	.	>100.	.	.
62	STATIC	LC	>100.	.	.		.	.	>100.	.	.
63	STATIC	LC	>100.	.	.		.	.	>100.	.	.
64	STATIC	LC	>100.	.	.		.	.	>100.	.	.
65	STATIC	LC	>100.	.	.		.	.	>100.	.	.
66	STATIC	LC	>100.	.	.		.	.	>100.	.	.

BEST AVAILABLE COPY

ED_005172C_00000035-00077

ACETONE

CAS # 67-64-1 PRINCIPAL USE: SOLVENT

MEASURED IN ML /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	12.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	6.01	4.660	7.750	.	.	.	5.54	4.740	6.330

ED_005172C_00000035-00078

AEROFROTH 71
MEASURED IN MG /L

PRINCIPAL USE: FLOTATION AGENT

OBS	CAS #	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0%	COMMERCIAL FORMULATION	CHANNEL CATFISH		1.10	15.0	7.4	272
2	100.0%	COMMERCIAL FORMULATION	BLUEGILL		0.60	15.0	7.4	272

OBS	TEST TYPE		24 HOUR CONFIDENCE INTERVAL		48 HOUR CONFIDENCE INTERVAL		96 HOUR CONFIDENCE INTERVAL	
1	STATIC	LC	100.000	1000.000	.	.	10.000	100.000
2	STATIC	LC	100.000	1000.000	.	.	10.000	100.000

ED_005172C_00000035-00079

AERO XANTHATE 343
MEASURED IN MG /L

CAS # PRINCIPAL USE: FLOTATION AGENT

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% COMMERCIAL FORMULATION	CHANNEL CATFISH		1.10	15.0	7.4	272
2	100.0% COMMERCIAL FORMULATION	BLUEGILL		0.60	15.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>10.	:	:		:	:	>10.	:	:
2	STATIC	LC	>10.	:	:		:	:	>10.	:	:

ED_005172C_00000035-00080

CAS # 1757-18-2 PRINCIPAL USE: INSECTICIDE

AKTON

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	26.00% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		1.00	12.0	7.4	40
2	26.00% EMULSIFIABLE CONCENTRATE	FATHEAD MINNOW		1.00	12.0	7.4	40
3	80.00% TECHNICAL MATERIAL	CHANNEL CATFISH		0.70	18.0	7.4	40
4	80.00% TECHNICAL MATERIAL	BLUEGILL		1.10	18.0	7.4	40
5	80.00% TECHNICAL MATERIAL	REDEAR SUNFISH		2.50	18.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	8200.	.	.		.	.	1200.	1080.000	1370.000
2	STATIC	LC	1350.	.	.		.	.	270.	.	.
3	STATIC	LC	4200.	3119.000	5657.000		.	.	400.	295.000	542.000
4	STATIC	LC	590.	350.000	990.000		.	.	170.	100.000	270.000
5	STATIC	LC	>2400.	.	.		.	.	380.	170.000	850.000

ED_005172C_00000035-00081

ALACHLOR

CAS # 15972-60-3 PRINCIPAL USE: HERBICIDE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	93.00% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.4	44
2	45.00% EMULSIFIABLE CONCENTRATE	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.4	44
3	93.00% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	22.0	7.4	44
4	45.00% EMULSIFIABLE CONCENTRATE	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	22.0	7.4	44
5	43.00% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		0.80	12.0	7.4	44
6	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	12.0	7.4	44
7	43.00% EMULSIFIABLE CONCENTRATE	BLUEGILL		1.00	22.0	7.4	44
8	100.0% TECHNICAL MATERIAL	BLUEGILL		1.00	22.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	21.0	15.900	27.800		.	.
2	STATIC	EC		.	.	7.7	6.000	9.900		.	.
3	STATIC	EC		.	.	3.2	2.300	4.400		.	.
4	STATIC	EC		.	.	2.5	1.600	3.600		.	.
5	STATIC	LC	4.3	3.610	5.110		.	.	1.4	1.080	1.820
6	STATIC	LC		.	.		.	.	2.4	1.840	3.120
7	STATIC	LC	7.8	6.010	10.110		.	.	3.2	2.290	4.460
8	STATIC	LC	11.5	8.580	15.410		.	.	4.3	3.530	5.500

ED_005172C_00000035-00082

ALDICARB MEASURED IN UG /L

CAS # 116-06-3 PRINCIPAL USE: INSECTICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		0.50	12.0	7.4	44
2	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		2.70	18.0	7.4	44
3	100.0% TECHNICAL MATERIAL	BLUEGILL		1.30	24.0	7.4	44
4	100.0% TECHNICAL MATERIAL	BLUEGILL		3.90	18.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	1000.	727.000	1376.000	.	.	.	560.	394.000	796.000
2	STATIC	LC	780.	592.000	1027.000	.	.	.	660.	472.000	921.000
3	STATIC	LC	103.	66.000	161.000	.	.	.	52.	34.000	79.000
4	STATIC	LC	160.	120.000	214.000	.	.	.	71.	54.000	93.000

ED_005172C_00000035-00083

ALDRIN

CAS # 309-00-2 PRINCIPAL USE: INSECTICIDE MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	90.00% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	15.0	7.1	40
2	90.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	15.0	7.1	40
3	90.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	21.0	7.1	40
4	90.00% TECHNICAL MATERIAL	CYPRIDOPSIS VIDUA	MATURE	.	21.0	7.4	40
5	90.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	IMMATURE	.	21.0	7.1	40
6	90.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	IMMATURE	.	16.0	7.1	40
7	90.00% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	IMMATURE	.	21.0	7.1	40
8	90.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	40
9	90.00% TECHNICAL MATERIAL	CHINOOK SALMON		0.80	15.0	7.2	81
10	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	2.0	7.1	40
11	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	7.0	7.1	40
12	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	13.0	7.1	40
13	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	18.0	7.1	40
14	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	13.0	7.1	40
15	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	13.0	7.7	135
16	90.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.60	18.0	7.1	40
17	90.00% TECHNICAL MATERIAL	BLACK BULLHEAD		1.50	24.0	7.1	40
18	90.00% TECHNICAL MATERIAL	CHANNEL CATFISH		5.20	18.0	7.4	40
19	90.00% TECHNICAL MATERIAL	BLUEGILL		0.80	18.0	7.1	40
20	90.00% TECHNICAL MATERIAL	BLUEGILL		0.80	18.0	7.7	135
21	90.00% TECHNICAL MATERIAL	BLUEGILL		0.70	7.0	7.4	40
22	90.00% TECHNICAL MATERIAL	BLUEGILL		0.70	13.0	7.4	40
23	90.00% TECHNICAL MATERIAL	BLUEGILL		0.70	18.0	7.4	40
24	90.00% TECHNICAL MATERIAL	BLUEGILL		0.70	24.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	28.	20.000	39.000		.	.
2	STATIC	EC		.	.	23.	17.000	30.000		.	.
3	STATIC	EC		.	.	32.	22.000	36.000		.	.
4	STATIC	EC		.	.	18.	15.000	21.000		.	.
5	STATIC	LC	56000.	42400.000	73900.000		.	.	4300.	3500.000	5300.000
6	STATIC	LC	52000.	38800.000	69700.000	12000.	7300.000	19800.000	5600.	3600.000	8700.000
7	STATIC	LC	120.	90.000	160.000		.	.	50.	38.000	65.000
8	STATIC	LC	30.	19.000	48.000		.	.	1.3	0.750	2.200
9	FLOW THROUGH	LC		.	.		.	.	14.3	8.820	23.200
10	STATIC	LC	24.0	19.000	31.000		.	.	3.40	1.600	7.200
11	STATIC	LC	7.50	6.100	9.200		.	.	3.2	2.800	3.600
12	STATIC	LC	6.30	5.700	7.500		.	.	2.60	2.300	2.900
13	STATIC	LC	4.50	4.000	5.000		.	.	2.60	2.500	2.900
14	STATIC	LC	19.0	11.000	32.000		.	.	8.2	6.000	11.100
15	STATIC	LC	19.2	11.400	32.400		.	.	9.3	6.800	10.300
16	STATIC	LC	15.	.	.		.	.	8.2	.	.
17	STATIC	LC	22.	.	.		.	.	19.	.	.
18	STATIC	LC	53.	.	.		.	.	53.	.	.
19	STATIC	LC	40.0	30.000	54.000		.	.	12.0	8.000	18.000
20	STATIC	LC	32.0	20.000	50.000		.	.	10.0	5.500	19.000
21	STATIC	LC	130.	105.000	160.000		.	.	9.7	8.400	11.300
22	STATIC	LC	36.8	31.400	43.000		.	.	7.7	6.600	9.000
23	STATIC	LC	16.4	14.000	19.000		.	.	6.2	5.200	7.700
24	STATIC	LC	9.3	7.400	11.700		.	.	5.6	4.700	6.700

23	STATIC	LC	9.3
24	STATIC	LC	

ALDRIN MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
25	90.00% TECHNICAL MATERIAL	LARGEMOUTH BASS		2.50	18.0	8.0	135
26	90.00% TECHNICAL MATERIAL	FOWLERS TOAD	TADPOLE	.	16.0	7.1	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
25	STATIC	LC	19.0	.	.	148.	29.000	768.000	5.0	17.000	270.000
26	STATIC	LC	>180.	.	.				68.		

BEST AVAILABLE COPY

83

ED_005172C_00000035-00085

CAS # PRINCIPAL USE: INSECTICIDE

ALLETHRIN RACEMIC MIXTURE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	90.00% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	15.0	7.1	40
2	90.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	15.0	7.1	40
3	90.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	21.0	7.1	40
4	90.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	40
5	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.90	13.0	7.1	40
6	90.00% TECHNICAL MATERIAL	BLUEGILL		0.90	24.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	21.0	19.000	35.000		.	.
2	STATIC	EC		.	.	56.0	40.000	78.000		.	.
3	STATIC	LC	38.0	30.000	49.000		.	.	11.0	8.000	15.000
4	STATIC	LC	9.0	7.000	11.000		.	.	5.6	4.900	6.400
5	STATIC	LC	20.0	.	.		.	.	19.0	.	.
6	STATIC	LC	65.0	.	.		.	.	56.0	.	.

84

ED_005172C_00000035-00086

ALUMINUM NITRATE
MEASURED IN MG /L

CAS # 13473-90-0 PRINCIPAL USE:

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	7.20 % TECHNICAL MATERIAL	FATHEAD MINNOW		0.45	22.0	7.4	38

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL	48 HOUR	CONFIDENCE INTERVAL	96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	.	.	.	.	4.25	3.300	5.500

ED_005172C_00000035-00087

CAS # 10043-01-3 PRINCIPAL USE:

ALUMINUM SULFATE
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	8.10 % TECHNICAL MATERIAL	FATHEAD MINNOW		0.45	22.0	7.4	38

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC		.	.		.	.	4.4	3.400	5.600

ED_005172C_00000035-00088

A'.JRO MEASURED IN UG /L

CAS # 67485-29-4 PRINCIPAL USE: INSECTICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	97.70% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	22.0	7.5	43
2	97.70% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	17.0	7.5	43
3	97.70% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	12.0	7.4	39
4	97.70% TECHNICAL MATERIAL	FATHEAD MINNOW		0.75	22.0	7.2	40
5	97.70% TECHNICAL MATERIAL	CHANNEL CATFISH		0.60	22.0	7.3	42
6	97.70% TECHNICAL MATERIAL	BLUEGILL		0.30	22.0	7.2	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC	.	.	.	130.	60.000	280.000	.	.	.
2	STATIC	EC	.	.	.	140.	100.000	180.000	76.	59.000	97.000
3	STATIC	LC	240.	171.000	337.000	.	.	.	75.	59.000	96.000
4	STATIC	LC	130.	90.000	190.000	.	.	.	70.	53.000	92.000
5	STATIC	LC	.	.	.	.	.	.	120.	87.000	166.000
6	STATIC	LC									

AMETRYN

MEASURED IN MG /L

CAS # 834-12-8 PRINCIPAL USE: HERBICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	13.0	7.1	40
2	99.00% TECHNICAL MATERIAL	BLUEGILL		0.80	24.0	7.1	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	8.0	:	:	3.4	:	:	3.2	:	:
2	STATIC	LC	10.0	:	:	5.0	:	:	3.7	:	:

ED_005172C_00000035-00090

AMINOCARB

CAS # 2032-59-9 PRINCIPAL USE: INSECTICIDE MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
1	98.00% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.1	42
2	17.00% OIL SOLUBLE	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.1	42
3	17.00% OIL SOLUBLE	DAPHNIA MAGNA	1ST INSTAR	.	21.0	7.4	40
4	17.00% OIL SOLUBLE	GAMMARUS FASCIATUS	MATURE	.	21.0	7.4	40
5	17.00% OIL SOLUBLE	GAMMARUS PSEUDOLIMNAEUS	IMMATURE	.	17.0	7.1	42
6	98.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	IMMATURE	.	17.0	7.1	42
7	98.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	6.5	40
8	98.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	7.5	40
9	98.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	8.5	40
10	98.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	8.5	40
11	98.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	6.5	40
12	98.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	7.5	40
13	98.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	8.5	40
14	98.00% TECHNICAL MATERIAL	SKWALA SP.	NAIAD	0.12	7.0	7.0	35
15	17.00% OIL SOLUBLE	SKWALA SP.	NAIAD	0.16	7.0	7.5	42
16	98.00% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	0.07	12.0	6.5	42
17	98.00% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	0.06	12.0	7.5	40
18	98.00% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	0.04	12.0	8.5	45
19	17.00% OIL SOLUBLE	PTERONARCELLA BADIA	1ST YEAR CLASS	0.07	12.0	6.5	42
20	17.00% OIL SOLUBLE	PTERONARCELLA BADIA	1ST YEAR CLASS	0.06	12.0	7.5	38
21	17.00% OIL SOLUBLE	PTERONARCELLA BADIA	1ST YEAR CLASS	0.05	12.0	8.5	38
22	17.00% OIL SOLUBLE	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	22.0	7.4	42
23	17.00% OIL SOLUBLE	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	21.0	7.4	40
24	98.00% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	20.0	7.2	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC	>1000.	.	.	320.	271.000	378.000		.	.
2	STATIC	EC		.	.	19.	16.000	23.000		.	.
3	STATIC	EC		.	.		10.000	100.000		.	.
4	STATIC	LC	39.	30.000	51.000	29.	21.000	39.000	12.	8.200	17.500
5	STATIC	LC	>100.	.	.		.	.	30.	17.000	55.000
6	STATIC	LC	>1000.	.	.		.	.	145.	96.000	220.000
7	STATIC	EC	2500.	1610.000	3880.000		.	.		.	.
8	STATIC	EC	4300.	3060.000	6040.000	3100.	1890.000	5080.000		.	.
9	STATIC	EC	2900.	2150.000	3920.000		.	.		.	.
10	STATIC	EC	4100.	3020.000	5560.000	1800.	1340.000	2410.000		.	.
11	STATIC	EC	5000.	3250.000	7680.000		.	.		.	.
12	STATIC	EC	3300.	2530.000	4300.000	2630.	2000.000	3460.000	2200.	1480.000	3270.000
13	STATIC	EC	2850.	2010.000	4040.000	1700.	1170.000	2460.000		.	.
14	STATIC	LC	58.	41.000	82.000		.	.	41.	27.000	62.000
15	STATIC	LC		.	.		.	.	23.	18.000	29.000
16	STATIC	LC	35.	30.000	41.000		.	.	28.	18.000	43.000
17	STATIC	LC	34.	23.000	49.000		.	.	19.	14.000	26.000
18	STATIC	LC		.	.		.	.	26.	20.000	33.000
19	STATIC	LC	60.	44.000	82.000		.	.	21.	16.000	28.000
20	STATIC	LC	56.	36.000	87.000		.	.	16.	12.000	20.000
21	STATIC	LC	37.	.	.		.	.	23.	17.000	31.000
22	STATIC	EC	>100.	.	.	30.	22.000	41.000		.	.
23	STATIC	EC		.	.	270.	150.000	490.000		.	.
24	STATIC	EC		.	.	13500.	5100.000	36000.000		.	.

ED_005172C_00000035-00091

AMINOCARB

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
25	98.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.80	10.0	7.4	206
26	17.00% OIL SOLUBLE	CUTTHROAT TROUT		1.00	10.0	7.4	206
27	98.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.70	12.0	7.5	42
28	98.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.70	7.0	7.5	42
29	98.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.90	12.0	6.5	40
30	98.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.90	12.0	8.5	40
31	98.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.80	12.0	7.8	40
32	98.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00	12.0	7.8	340
33	17.00% OIL SOLUBLE	CUTTHROAT TROUT		0.60	12.0	7.5	40
34	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	10.0	7.2	40
35	17.00% OIL SOLUBLE	RAINBOW TROUT		1.50	10.0	7.2	40
36	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.20	7.0	7.4	40
37	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.20	12.0	7.4	40
38	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.20	17.0	7.4	40
39	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.30	12.0	7.4	40
40	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	12.0	7.4	320
41	17.00% OIL SOLUBLE	RAINBOW TROUT		0.40	10.0	8.1	10
42	17.00% OIL SOLUBLE	RAINBOW TROUT		0.40	10.0	8.1	40
43	17.00% OIL SOLUBLE	RAINBOW TROUT		0.40	10.0	8.1	160
44	17.00% OIL SOLUBLE	RAINBOW TROUT		0.40	10.0	8.1	320
45	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.40	12.0	8.5	40
46	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.40	12.0	7.5	40
47	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.40	12.0	8.5	40
48	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.40	12.0	9.5	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
25	STATIC	LC	38500.	26200.000	56500.000	.	.	.	31000.	18000.000	52400.000
26	STATIC	LC	1400.	858.000	2290.000	.	.	.	64.	31.000	134.000
27	STATIC	LC	32400.	27500.000	38200.000	.	.	.	28000.	24000.000	32600.000
28	STATIC	LC	28000.	24200.000	32400.000	.	.	.	27000.	21500.000	34000.000
29	STATIC	LC	29800.	24200.000	36700.000	.	.	.	26500.	21500.000	32600.000
30	STATIC	LC		.	.	.	.	.	1300.	1090.000	1560.000
31	STATIC	LC		.	.	.	.	.	22000.	18800.000	25700.000
32	STATIC	LC	29000.	23100.000	36500.000	.	.	.	24500.	19400.000	31000.000
33	STATIC	LC	200.	151.000	264.000	.	.	.	88.	73.000	106.000
34	STATIC	LC	32000.	23000.000	44000.000	.	.	.	13500.	11300.000	16200.000
35	STATIC	LC	135.	113.000	162.000	.	.	.	130.	103.000	164.000
36	STATIC	LC	>32000	.	.	.	.	.	27000.	19100.000	38200.000
37	STATIC	LC	>32000	.	.	.	.	.	18000.	12600.000	25700.000
38	STATIC	LC	30000.	18700.000	48000.000	.	.	.	17000.	12100.000	23900.000
39	STATIC	LC	>32000	.	.	.	.	.	25000.	17800.000	35100.000
40	STATIC	LC	>32000	.	.	.	.	.	12000.	9420.000	15300.000
41	STATIC	LC	135.	113.000	162.000	.	.	.	79.	63.000	98.000
42	STATIC	LC	125.	100.000	157.000	.	.	.	88.	71.000	109.000
43	STATIC	LC	150.	118.000	190.000	.	.	.	75.	63.000	90.000
44	STATIC	LC	170.	140.000	206.000	.	.	.	85.	69.000	104.000
45	STATIC	LC	33000.	26200.000	41600.000	.	.	.	17500.	15600.000	19600.000
46	STATIC	LC	28000.	26000.000	30100.000	.	.	.	21000.	18700.000	23600.000
47	STATIC	LC	25000.	22700.000	27500.000	.	.	.	21200.	19400.000	23100.000
48	STATIC	LC	5100.	4250.000	6120.000	.	.	.	<5000.	.	.

ED_005172C_00000035-00092

47 STATIC
48 STATIC LC 51000.

AMINOCARB
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
49	17.00% OIL SOLUBLE	RAINBOW TROUT		1.00	12.0	7.2	40
50	17.00% OIL SOLUBLE	RAINBOW TROUT		1.20	10.0	7.2	40
51	17.00% OIL SOLUBLE	RAINBOW TROUT		0.80	10.0	7.2	40
52	17.00% OIL SOLUBLE	RAINBOW TROUT		0.80	10.0	7.2	40
53	17.00% OIL SOLUBLE	RAINBOW TROUT		0.80	10.0	7.2	40
54	17.00% OIL SOLUBLE	RAINBOW TROUT		3.70	10.0	7.4	272
55	17.00% OIL SOLUBLE	RAINBOW TROUT	EYED EGG	.	12.0	7.2	40
56	17.00% OIL SOLUBLE	RAINBOW TROUT	EYED EGG	.	10.0	7.4	272
57	17.00% OIL SOLUBLE	RAINBOW TROUT	SWIMUP FRY	.	12.0	7.4	40
58	17.00% OIL SOLUBLE	RAINBOW TROUT	YOLK-SAC FRY	.	12.0	7.2	40
59	17.00% OIL SOLUBLE	RAINBOW TROUT	YOLK-SAC FRY	.	10.0	7.4	272
60	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40	12.0	7.5	40
61	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40	7.0	7.5	40
62	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.60	17.0	7.5	40
63	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40	12.0	6.5	40
64	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40	12.0	8.5	40
65	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40	12.0	8.0	10
66	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40	12.0	8.0	40
67	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		1.00	12.0	7.5	40
68	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		1.00	12.0	7.5	40
69	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		1.00	12.0	7.5	40
70	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		1.00	12.0	7.5	40
71	98.00% TECHNICAL MATERIAL	ATLANTIC SALMON		1.00	12.0	7.5	40
72	98.00% TECHNICAL MATERIAL	BROWN TROUT		0.50	12.0	7.5	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
49	STATIC	LC	56.	40.000	79.000	.	.	.	46.	38.000	56.000
50	0 DAY DEGRA	LC	280.	230.000	340.000	.	.	.	140.	120.000	170.000
51	1 DAY DEGRA	LC	130.	103.000	164.000	.	.	.	76.	56.000	103.000
52	3 DAY DEGRA	LC	130.	103.000	164.000	.	.	.	86.	68.000	108.000
53	7 DAY DEGRA	LC	130.	103.000	164.000	.	.	.	115.	92.000	144.000
54	FLOW THROUGH	LC		.	.	.	.	.	54.	49.000	59.000
55	STATIC	LC	100000	.	.	.	.	.	>32000	.	.
56	FLOW THROUGH	LC		.	.	.	.	.	>130.	.	.
57	STATIC	LC	95.	76.000	120.000	.	.	.	48.	36.000	63.000
58	STATIC	LC	80.	64.000	100.000	.	.	.	36.	30.000	44.000
59	FLOW THROUGH	LC		.	.	.	.	.	38.	36.000	40.000
60	STATIC	LC	7600.	5880.000	9820.000	.	.	.	7600.	5880.000	9820.000
61	STATIC	LC	4250.	3170.000	5780.000	.	.	.	4200.	2240.000	7880.000
62	STATIC	LC	10500.	8740.000	12600.000	.	.	.	7000.	6010.000	8150.000
63	STATIC	LC	11300.	7470.000	17100.000	.	.	.	8550.	6310.000	11600.000
64	STATIC	LC	6850.	5520.000	8500.000	.	.	.	6030.	4550.000	7990.000
65	STATIC	LC	4750.	3020.000	7460.000	.	.	.	4750.	3020.000	7460.000
66	STATIC	LC	5700.	3490.000	9300.000	.	.	.	4050.	2090.000	7830.000
67	0 DAY DEGRA	LC		.	.	.	.	.	2350.	1690.000	3260.000
68	7 DAY DEGRA	LC		.	.	.	.	.	3500.	2980.000	4110.000
69	14 DAY DEGRA	LC		.	.	.	.	.	4700.	3040.000	7270.000
70	21 DAY DEGRA	LC		.	.	.	.	.	3500.	2640.000	4980.000
71	28 DAY DEGRA	LC		.	.	.	.	.	2800.	1770.000	4440.000
72	STATIC	LC	20000.	17000.000	23500.000	.	.	.	15000.	11500.000	19500.000

ED_005172C_00000035-00093

AMINOCARB
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
73	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.00	7.0	7.5	40
74	98.00% TECHNICAL MATERIAL	BROWN TROUT		0.50	12.0	6.5	40
75	98.00% TECHNICAL MATERIAL	BROWN TROUT		0.50	12.0	8.5	40
76	98.00% TECHNICAL MATERIAL	BROWN TROUT		0.50	12.0	9.0	40
77	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	8.1	10
78	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	8.1	40
79	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	8.1	160
80	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	8.1	280
81	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	7.5	40
82	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	7.5	40
83	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	7.5	40
84	98.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	7.5	40
85	98.00% TECHNICAL MATERIAL	BROOK TROUT		0.10	12.0	7.5	40
86	98.00% TECHNICAL MATERIAL	BROOK TROUT		0.80	12.0	7.5	40
87	98.00% TECHNICAL MATERIAL	BROOK TROUT		1.00	7.0	7.5	40
88	98.00% TECHNICAL MATERIAL	BROOK TROUT		1.00	17.0	7.5	40
89	98.00% TECHNICAL MATERIAL	BROOK TROUT		0.80	12.0	6.5	40
90	98.00% TECHNICAL MATERIAL	BROOK TROUT		0.80	12.0	7.5	40
91	98.00% TECHNICAL MATERIAL	BROOK TROUT		1.20	12.0	8.5	40
92	98.00% TECHNICAL MATERIAL	BROOK TROUT		0.80	12.0	9.0	40
93	98.00% TECHNICAL MATERIAL	BROOK TROUT		1.00	12.0	8.0	10
94	98.00% TECHNICAL MATERIAL	BROOK TROUT		1.00	12.0	8.0	40
95	98.00% TECHNICAL MATERIAL	BROOK TROUT		1.00	12.0	8.0	160
96	98.00% TECHNICAL MATERIAL	BROOK TROUT		0.20	12.0	8.0	280

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
73	STATIC	LC	35000.	24400.000	50100.000	.	.	.	27500.	21400.000	35300.000
74	STATIC	LC	9200.	5760.000	12500.000	.	.	.	8500.	5950.000	12100.000
75	STATIC	LC	13500.	10700.000	17000.000	.	.	.	8800.	6870.000	11300.000
76	STATIC	LC	17000.	9700.000	29800.000	.	.	.	3150.	2280.000	4340.000
77	STATIC	LC	16000.	12600.000	25800.000	.	.	.	14000.	11400.000	17200.000
78	STATIC	LC	14300.	11100.000	18400.000	.	.	.	11500.	9730.000	13600.000
79	STATIC	LC		.	.	.	.	.	9000.	6200.000	13000.000
80	STATIC	LC	17100.	14600.000	20000.000	.	.	.	14000.	10400.000	18900.000
81	0 DAY DEGRA	LC		.	.	.	.	.	17000.	13500.000	21400.000
82	7 DAY DEGRA	LC		.	.	.	.	.	8500.	6910.000	10700.000
83	14 DAY DEGRA	LC		.	.	.	.	.	7000.	5210.000	9410.000
84	21 DAY DEGRA	LC		.	.	.	.	.	6100.	.	.
85	STATIC	LC	23000.	16900.000	31300.000	.	.	.	16000.	12700.000	20100.000
86	STATIC	LC	13000.	10500.000	16000.000	.	.	.	12100.	9600.000	15200.000
87	STATIC	LC	12400.	7930.000	19400.000	.	.	.	9400.	6870.000	12900.000
88	STATIC	LC	3150.	2570.000	3860.000	.	.	.	2600.	2090.000	3230.000
89	STATIC	LC		.	.	.	.	.	11000.	7700.000	15600.000
90	STATIC	LC		.	.	.	.	.	10500.	6100.000	18100.000
91	STATIC	LC	7400.	6200.000	8800.000	.	.	.	4300.	3400.000	5400.000
92	STATIC	LC		.	.	.	.	.	2200.	1800.000	2700.000
93	STATIC	LC	14800.	10200.000	21500.000	.	.	.	7700.	5300.000	11200.000
94	STATIC	LC	6100.	4120.000	9040.000	.	.	.	4850.	4020.000	5850.000
95	STATIC	LC	11600.	8310.000	16200.000	.	.	.	8000.	5800.000	11000.000
96	STATIC	LC	19000.	13900.000	25900.000	.	.	.	10600.	8660.000	13000.000

ED_005172C_00000035-00094

95 STATIC
96 STATIC LC

AMINOCARB
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
97	17.00% OIL SOLUBLE	BROOK TROUT		1.20	12.0	7.5	40
98	17.00% OIL SOLUBLE	BROOK TROUT		1.20	12.0	6.5	40
99	17.00% OIL SOLUBLE	BROOK TROUT		1.20	12.0	8.5	40
100	17.00% OIL SOLUBLE	BROOK TROUT		1.20	12.0	9.0	40
101	98.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.70	20.0	7.2	40
102	17.00% OIL SOLUBLE	FATHEAD MINNOW		0.70	20.0	7.2	40
103	98.00% TECHNICAL MATERIAL	CHANNEL CATFISH		0.60	20.0	7.4	40
104	17.00% OIL SOLUBLE	CHANNEL CATFISH		4.50	20.0	7.2	40
105	98.00% TECHNICAL MATERIAL	CHANNEL CATFISH		4.10	20.0	7.2	40
106	17.00% OIL SOLUBLE	CHANNEL CATFISH		0.90	17.0	7.4	40
107	17.00% OIL SOLUBLE	CHANNEL CATFISH		2.00	22.0	7.2	40
108	98.00% TECHNICAL MATERIAL	BLUEGILL		0.60	20.0	7.2	40
109	17.00% OIL SOLUBLE	BLUEGILL		0.60	20.0	7.2	40
110	98.00% TECHNICAL MATERIAL	BLUEGILL		0.90	12.0	7.5	40
111	98.00% TECHNICAL MATERIAL	BLUEGILL		1.90	12.0	7.5	40
112	98.00% TECHNICAL MATERIAL	BLUEGILL		0.40	17.0	7.4	40
113	98.00% TECHNICAL MATERIAL	BLUEGILL		0.40	17.0	7.4	320
114	98.00% TECHNICAL MATERIAL	BLUEGILL		0.40	12.0	7.4	40
115	98.00% TECHNICAL MATERIAL	BLUEGILL		0.40	17.0	7.4	40
116	98.00% TECHNICAL MATERIAL	BLUEGILL		0.40	22.0	7.4	40
117	98.00% TECHNICAL MATERIAL	BLUEGILL		0.70	12.0	6.5	40
118	98.00% TECHNICAL MATERIAL	BLUEGILL		0.70	12.0	7.5	40
119	98.00% TECHNICAL MATERIAL	BLUEGILL		0.70	12.0	8.5	40
120	98.00% TECHNICAL MATERIAL	BLUEGILL		0.70	12.0	9.5	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
97	STATIC	LC	162.	128.000	204.000	.	.	.	151.	122.000	187.000
98	STATIC	LC	180.	134.000	241.000	.	.	.	140.	113.000	174.000
99	STATIC	LC	180.	138.000	235.000	.	.	.	142.	114.000	176.000
100	STATIC	LC	142.	115.000	176.000	.	.	.	142.	115.000	176.000
101	STATIC	LC	10000.	7400.000	13400.000	.	.	.	8500.	6500.000	11100.000
102	STATIC	LC	75.	63.000	90.000	.	.	.	75.	63.000	90.000
103	STATIC	LC	>32000	.	.	.	.	.	10000.	5970.000	14330.000
104	STATIC	LC	245.	159.000	377.000	.	.	.	130.	105.000	160.000
105	STATIC	LC	47000.	28000.000	80000.000	.	.	.	4200.	3500.000	5000.000
106	STATIC	LC	75.	50.000	113.000	.	.	.	75.	63.000	89.000
107	STATIC	LC	240.	200.000	286.000	.	.	.	140.	104.000	187.000
108	STATIC	LC	7500.	4500.000	12400.000	.	.	.	3100.	1900.000	5100.000
109	STATIC	LC	118.	91.000	153.000	.	.	.	100.	68.000	148.000
110	STATIC	LC	25000.	18600.000	33700.000	.	.	.	20000.	14500.000	27600.000
111	STATIC	LC	20000.	15700.000	25400.000	.	.	.	12500.	9130.000	17100.000
112	STATIC	LC	8200.	5990.000	11220.000	.	.	.	3300.	2300.000	4730.000
113	STATIC	LC	7200.	5240.000	9890.000	.	.	.	4600.	3380.000	6270.000
114	STATIC	LC		.	.	.	.	.	4200.	3190.000	5530.000
115	STATIC	LC		.	.	.	.	.	6000.	4310.000	8360.000
116	STATIC	LC		.	.	.	.	.	3500.	2410.000	5070.000
117	STATIC	LC	13200.	10900.000	16100.000	.	.	.	10500.	8510.000	13000.000
118	STATIC	LC	14800.	12600.000	17300.000	.	.	.	8400.	6470.000	10900.000
119	STATIC	LC	16000.	12800.000	20000.000	.	.	.	5200.	3830.000	7070.000
120	STATIC	LC	3200.	2510.000	4080.000	.	.	.	<1000.	.	.

ED_005172C_00000035-00095

AMINOCARB

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
121	17.00% OIL SOLUBLE	BLUEGILL		0.30	17.0	7.4	40
122	17.00% OIL SOLUBLE	BLUEGILL		0.30	22.0	7.4	40
123	17.00% OIL SOLUBLE	BLUEGILL		0.30	27.0	7.4	40
124	17.00% OIL SOLUBLE	BLUEGILL		1.50	20.0	7.2	40
125	17.00% OIL SOLUBLE	BLUEGILL		1.50	20.0	7.2	40
126	17.00% OIL SOLUBLE	BLUEGILL		1.50	20.0	7.2	40
127	17.00% OIL SOLUBLE	BLUEGILL		1.50	20.0	7.2	40
128	17.00% OIL SOLUBLE	BLUEGILL		3.10	20.0	7.4	272
129	98.00% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.70	18.0	7.8	135
130	98.00% TECHNICAL MATERIAL	YELLOW PERCH		0.60	12.0	7.5	40
131	17.00% OIL SOLUBLE	YELLOW PERCH		2.00	12.0	7.5	40
132	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	7.0	7.5	40
133	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	7.5	40
134	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	17.0	7.5	40
135	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	22.0	7.5	40
136	98.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	6.5	40
137	98.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	7.5	40
138	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	8.5	40
139	98.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	9.0	40
140	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.80	12.0	8.0	10
141	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	8.0	40
142	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	8.0	160
143	98.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	8.0	280
144	98.00% TECHNICAL MATERIAL	WALLEYE		0.80	18.0	7.8	135

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
121	STATIC	LC	130.	100.000	170.000	.	.	.	75.	63.000	90.000
122	STATIC	LC	110.	88.000	130.000	.	.	.	57.	49.000	67.000
123	STATIC	LC	69.	57.000	83.000	.	.	.	45.	36.000	56.000
124	0 DAY DEGRA	LC	270.	209.000	348.000	.	.	.	240.	201.000	286.000
125	1 DAY DEGRA	LC	240.	201.000	286.000	.	.	.	200.	144.000	278.000
126	3 DAY DEGRA	LC	250.	201.000	311.000	.	.	.	180.	153.000	212.000
127	7 DAY DEGRA	LC	240.	201.000	286.000	.	.	.	240.	201.000	286.000
128	FLOW THROUGH	LC		.	.	.	.	.	100.	95.000	106.000
129	STATIC	LC	8100.	6720.000	9760.000	.	.	.	3150.	2840.000	3490.000
130	STATIC	LC	6400.	5300.000	7730.000	.	.	.	6400.	5300.000	7730.000
131	STATIC	LC	1920.	1320.000	2800.000	.	.	.	230.	184.000	288.000
132	STATIC	LC	13000.	11600.000	14600.000	.	.	.	11700.	9990.000	13700.000
133	STATIC	LC	5950.	4880.000	7250.000	.	.	.	5600.	4710.000	6660.000
134	STATIC	LC	4700.	3960.000	5580.000	.	.	.	4700.	3960.000	5580.000
135	STATIC	LC	2100.	1600.000	2750.000	.	.	.	1700.	1330.000	2180.000
136	STATIC	LC	6600.	4870.000	8940.000	.	.	.	6400.	4730.000	8650.000
137	STATIC	LC	7000.	5510.000	8890.000	.	.	.	7000.	5510.000	8890.000
138	STATIC	LC		.	.	.	.	.	1430.	950.000	2160.000
139	STATIC	LC	5550.	4330.000	7120.000	.	.	.	425.	350.000	520.000
140	STATIC	LC	7800.	6270.000	9700.000	.	.	.	5400.	3740.000	7790.000
141	STATIC	LC	6700.	5350.000	8390.000	.	.	.	6200.	4950.000	7760.000
142	STATIC	LC	8400.	6410.000	11000.000	.	.	.	7400.	5220.000	10500.000
143	STATIC	LC	6100.	4890.000	7610.000	.	.	.	5600.	4770.000	6580.000
144	STATIC	LC	2000.	1570.000	2540.000	.	.	.	880.	755.000	1030.000

ED_005172C_00000035-00096

141	STATIC	LC	8400.	[illegible]	7610.000
142	STATIC	LC	6100.	4890.000	2540.000
143	STATIC	LC	2000.	1570.000	
144	STATIC	LC			

AMITROLE

MEASURED IN MG /L

CAS # 61-82-5 PRINCIPAL USE: HERBICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	18.0	7.4	40
2	100.0% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	18.0	7.4	272
3	100.0% TECHNICAL MATERIAL	FATHEAD MINNOW		1.20	18.0	7.1	40
4	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		1.80	18.0	7.1	40
5	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		1.80	18.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>10.0	.	.	.	.	.	>10.0	.	.
2	STATIC	LC	>10.0	.	.	.	.	.	>10.0	.	.
3	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
4	STATIC	LC	>160.	.	.	.	.	.	>160.	.	.
5	STATIC	LC	>160.	.	.	.	.	.	>160.	.	.

BEST AVAILABLE COPY

ED_005172C_00000035-00097

ANILAZINE

MEASURED IN UG /L

CAS # 101-05-3 PRINCIPAL USE: FUNGICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	95.50% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.1	44
2	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	13.0	7.1	44
3	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		0.70	18.0	7.1	44
4	95.50% TECHNICAL MATERIAL	BLUEGILL		1.10	18.0	7.1	44
5	95.50% TECHNICAL MATERIAL	REDEAR SUNFISH		2.50	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	1250	950.000	1640.000	.	.	.	270	210.000	350.000
2	STATIC	LC	145.	127.000	166.000	.	.	.	140.	122.000	161.000
3	STATIC	LC	270.	120.000	640.000	.	.	.	240.	110.000	550.000
4	STATIC	LC	600.	319.000	1130.000	.	.	.	320.	142.000	735.000
5	STATIC	LC	<140.	.	.	.	.	.	<140.	.	.

ED_005172C_00000035-00098

ANTIMYCIN A
MEASURED IN NG /L

CAS # 1397-94-0 PRINCIPAL USE: PISCICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	95.50% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INST.	.	21.0	7.1	44
2	95.50% TECHNICAL MATERIAL	ASELLUS BREVICAUDUS	MATURE	.	15.0	7.4	272
3	95.50% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	272
4	95.50% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	MATURE	.	21.0	7.4	272
5	95.50% TECHNICAL MATERIAL	PADDLEFISH		0.01	15.0	7.4	272
6	95.50% TECHNICAL MATERIAL	COHO SALMON		0.90	12.0	7.1	44
7	95.50% TECHNICAL MATERIAL	COHO SALMON		5.60	12.0	7.1	44
8	95.50% TECHNICAL MATERIAL	COHO SALMON		11.10	12.0	7.1	44
9	95.50% TECHNICAL MATERIAL	COHO SALMON		19.40	12.0	7.1	44
10	95.50% TECHNICAL MATERIAL	COHO SALMON		1.50	12.0	7.1	44
11	95.50% TECHNICAL MATERIAL	COHO SALMON		0.50	17.0	7.4	270
12	95.50% TECHNICAL MATERIAL	COHO SALMON		19.50	12.0	7.1	40
13	95.50% TECHNICAL MATERIAL	COHO SALMON		1.80	16.0	7.1	40
14	95.50% TECHNICAL MATERIAL	COHO SALMON		1.50	12.0	7.1	40
15	95.50% TECHNICAL MATERIAL	COHO SALMON		0.75	16.0	7.1	40
16	95.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.10	9.0	7.4	162
17	95.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00	9.0	7.9	170
18	95.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.10	9.0	7.9	170
19	95.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.10	9.0	7.9	170
20	95.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.10	9.0	7.9	170
21	95.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.10	9.0	7.9	170
22	95.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00	9.0	7.9	170
23	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	12.0	7.1	44
24	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		4.70	12.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.		5000.000	10000.000		.	.
2	STATIC	LC	>1000.	.	.		.	.	>1000	.	.
3	STATIC	LC	10.0	7.500	13.000		.	.	8.0	5.800	11.000
4	STATIC	LC	>10000	.	.		.	.		3000.000	6000.000
5	STATIC	LC	3.	1.000	12.000		.	.	1.0	0.400	3.000
6	STATIC	LC	151.	99.000	233.000		.	.	18.	12.000	28.000
7	STATIC	LC	110.	62.000	197.000		.	.	9.	3.000	30.000
8	STATIC	LC	210.	96.000	460.000		.	.	20.	9.000	42.000
9	STATIC	LC	>650.	.	.		.	.	60.	40.000	89.000
10	STATIC	LC	>120.	.	.		.	.	21.	8.000	54.000
11	STATIC	LC	34.3	4.070	288.000	15.1	2.080	110.000	15.1	2.080	110.000
12	STATIC	LC	>420.	.	.	92.7	14.300	603.000		.	.
13	STATIC	LC	60.5	41.300	88.800	31.3	24.300	40.300	9.1	5.900	14.000
14	STATIC	LC	120.	7.800	1860.000	31.7	18.400	54.800		.	.
15	STATIC	LC	35.6	9.000	140.000	7.1	3.300	15.500		.	.
16	STATIC	LC	433.	283.000	660.000		.	.	57.	19.000	166.000
17	STATIC	LC		.	.		.	.	77.6	60.300	99.700
18	STATIC	LC		.	.		.	.	84.1	72.400	97.600
19	STATIC	LC		.	.		.	.	83.1	71.500	96.600
20	STATIC	LC		.	.		.	.	87.0	74.900	101.000
21	STATIC	LC		.	.	159.	124.000	205.000	60.	.	.
22	STATIC	LC		.	.		.	.	112.	78.500	160.000
23	STATIC	LC	71.	24.000	205.000		.	.	14.	7.000	27.000
24	STATIC	LC	126.	52.000	306.000		.	.	22.	8.000	59.000

ED_005172C_00000035-00099

ANTIMYCIN A
MEASURED IN NG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT			DEG. CENT.	PH	HARDNESS MG/L
25	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		10.80			12.0	7.1	44
26	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		23.50			12.0	7.1	44
27	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		46.00			12.0	7.1	44
28	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		107.00			12.0	7.1	44
29	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.70	FISH PROTEIN CONCENTRATE	DIET	10.0	7.4	44
30	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	EWOS	DIET	10.0	7.4	44
31	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	GLENCO	DIET	10.0	7.4	44
32	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	FPRL 50 PERCENT PROTEIN	DIET	10.0	7.4	44
33	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	DONALDSON	STRAIN	12.0	7.4	44
34	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	IOWA	STRAIN	12.0	7.4	44
35	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		3.40	MISSOURI	STRAIN	12.0	7.4	44
36	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.30	MT. WHITNEY	STRAIN	12.0	7.4	44
37	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.90	NEW HAMPSHIRE	STRAIN	12.0	7.4	44
38	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	SOAP LAKE	STRAIN	12.0	7.4	44
39	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		3.60	WYTHEVILLE	STRAIN	12.0	7.4	44
40	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.30			12.0	7.1	40
41	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.60			12.0	7.4	40
42	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00			12.0	7.1	40
43	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00			12.0	7.1	40
44	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.50			12.0	7.1	40
45	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20			12.0	7.1	40
46	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.70			12.0	7.1	40
47	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.70			12.0	7.1	40
48	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.40			12.0	7.1	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
25	STATIC	LC	347.	181.000	669.000		.	.	26.	9.000	80.000
26	STATIC	LC	191.	113.000	321.000		.	.	30.	12.000	74.000
27	STATIC	LC	120.	22.000	660.000		.	.	115.	49.000	272.000
28	STATIC	LC	314.	239.000	413.000		.	.	89.	68.000	117.000
29	STATIC	LC		85.000	120.000		.	.	24.	20.000	29.000
30	STATIC	LC		60.000	120.000		.	.	16.	12.000	21.000
31	STATIC	LC		85.000	120.000		.	.	32.	27.000	38.000
32	STATIC	LC		65.000	120.000		.	.	52.	44.000	62.000
33	STATIC	LC	>87.	.	.		.	.	13.	6.000	28.000
34	STATIC	LC	49.	38.000	64.000		.	.	<8.	.	.
35	STATIC	LC	>67.	.	.		.	.	11.	5.000	23.000
36	STATIC	LC	>87.	.	.		.	.	31.	10.000	93.000
37	STATIC	LC	85.	42.000	173.000		.	.	9.	1.000	61.000
38	STATIC	LC	>75.	.	.		.	.	7.	4.000	15.000
39	STATIC	LC	100.	.	.		.	.	8.	3.000	23.000
40	STATIC	LC	87.5	44.300	173.000	46.7	30.200	72.200	12.2	6.600	22.700
41	STATIC	LC	421.	192.000	920.000	194.	66.800	565.000	91.6	39.800	211.000
42	STATIC	LC	98.4	56.100	173.000	55.4	48.500	63.200	26.1	17.600	38.900
43	STATIC	LC	57.6	47.800	69.500	36.5	30.500	43.500	22.0	18.200	26.600
44	STATIC	LC	54.0	45.000	64.600	23.8	19.900	28.500	9.1	5.600	14.700
45	STATIC	LC	63.5	27.200	148.000	28.0	12.100	64.400	11.7	5.300	25.800
46	STATIC	LC	57.9	34.300	97.800	16.2	6.300	41.400	10.9	6.500	18.300
47	STATIC	LC	96.0	45.400	203.000	29.0	14.900	56.400	10.2	.	.
48	STATIC	LC	68.5	20.400	230.000		.	.		.	.

ED_005172C_00000035-00100

46	STATIC	LC	[illegible]	[illegible]
47	STATIC	LC	68.5	20.400
48	STATIC	LC		

ANTIMYCIN A
MEASURED IN NG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT			DEG. CENT.	PH	HARDNESS MG/L
49	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20			12.0	7.1	40
50	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		2.00	MT. WHITNEY	STRAIN	12.0	7.1	40
51	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	MISSOURI	STRAIN	12.0	7.1	40
52	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.30	MISSOURI	STRAIN	12.0	7.1	40
53	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	MISSOURI	STRAIN	12.0	7.1	40
54	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	MISSOURI	STRAIN	12.0	7.1	40
55	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	MISSOURI	STRAIN	17.0	7.1	40
56	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	MT. WHITNEY	STRAIN	12.0	7.1	40
57	95.50% TECHNICAL MATERIAL	LAKE TROUT	FINGERLING	.			12.0	7.4	44
58	95.50% TECHNICAL MATERIAL	GOLDFISH		0.90			18.0	7.4	44
59	95.50% TECHNICAL MATERIAL	GOLDFISH		1.00			17.0	7.1	40
60	95.50% TECHNICAL MATERIAL	GOLDFISH		1.00			17.0	7.1	40
61	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.80			17.0	7.1	44
62	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.50			12.0	7.6	314
63	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		2.10			20.0	7.4	40
64	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		1.00			12.0	7.1	40
65	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.50			17.0	7.1	40
66	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.90			17.0	7.1	40
67	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		1.20			17.0	7.1	40
68	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.70			17.0	7.1	40
69	95.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.50			8.0	7.5	320
70	95.50% TECHNICAL MATERIAL	BLACK BULLHEAD		1.20			18.0	7.1	44
71	95.50% TECHNICAL MATERIAL	BLACK BULLHEAD		1.20			17.0	7.1	40
72	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		0.10			25.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
49	STATIC	LC	69.5	31.800	152.000	19.6	9.900	38.800	10.4	5.600	19.300
50	STATIC	LC	.	.	.		49.000	87.000	39.2	18.200	84.600
51	STATIC	LC	120.	60.000	241.000	63.7	34.300	118.000	25.	.	.
52	STATIC	LC	81.6	40.100	166.000	.	.	.		.	.
53	STATIC	LC	142.	106.000	191.000	61.8	46.200	82.700		.	.
54	STATIC	LC	89.5	66.100	121.000	37.0	27.700	49.600	13.4	.	.
55	STATIC	LC	54.9	41.300	73.100	21.7	15.100	31.200	.	.	.
56	STATIC	LC	.	.	.	67.1	7.500	600.000	11.1	5.400	230.000
57	STATIC	LC	53.	45.000	63.000		.	.	.	.	.
58	STATIC	LC	1030.	800.000	1300.000		.	.	180.	99.000	348.000
59	STATIC	LC	2100.	1300.000	3200.000	500.	300.000	700.000	200.	90.000	300.000
60	STATIC	LC	510.	310.000	840.000	250.	180.000	320.000	190.	50.000	400.000
61	STATIC	LC	515.	155.000	1710.000		.	.	40.	21.000	77.000
62	FLOW THROUGH	LC		390.000	530.000		.	.	38.	23.000	61.000
63	STATIC	LC	910.	644.000	1290.000	468.	272.000	807.000	265.	102.000	687.000
64	STATIC	LC	194.	141.000	267.000	63.2	39.900	99.900		.	.
65	STATIC	LC	109.	73.100	165.000	69.9	43.200	113.000	32.2	20.300	50.500
66	STATIC	LC	121.	64.200	228.000	75.1	27.500	205.000	25.	8.400	74.000
67	STATIC	LC	250.	170.000	350.000	120.	80.000	170.000	80.	70.000	100.000
68	STATIC	LC		.	.	83.3	45.000	154.000	40.2	20.000	80.700
69	FLOW THROUGH	LC	>300.	.	.		.	.	165.	147.000	185.000
70	STATIC	LC	7500.	4300.000	13100.000		.	.	7500.	4300.000	13100.000
71	STATIC	LC	36800.	20300.000	67000.000		.	.	4800.	3400.000	6800.000
72	STATIC	LC	2070	1530.000	2810.000		.	.	1360	1020.000	1820.000

BEST AVAILABLE COPY

ED_005172C_00000035-00101

ANTIMYCIN A
MEASURED IN NG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
73	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		0.15	21.0	7.1	44
74	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		1.90	17.0	7.1	44
75	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		5.00	17.0	7.1	44
76	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		9.70	17.0	7.1	44
77	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		18.50	17.0	7.1	44
78	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		2.30	17.0	7.1	40
79	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		2.10	17.0	7.1	40
80	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		1.40	17.0	7.1	40
81	95.50% TECHNICAL MATERIAL	WALKING CATFISH		3.80	25.0	7.1	44
82	95.50% TECHNICAL MATERIAL	MOSQUITOFISH		0.60 NON-RESISTANT STRAIN	17.0	7.1	44
83	95.50% TECHNICAL MATERIAL	MOSQUITOFISH		0.60 RESISTANT STRAIN	17.0	7.1	44
84	95.50% TECHNICAL MATERIAL	GREEN SUNFISH		1.10	18.0	7.1	44
85	95.50% TECHNICAL MATERIAL	BLUEGILL		1.20	17.0	7.1	44
86	95.50% TECHNICAL MATERIAL	BLUEGILL		4.80	17.0	7.1	44
87	95.50% TECHNICAL MATERIAL	BLUEGILL		9.70	17.0	7.1	44
88	95.50% TECHNICAL MATERIAL	BLUEGILL		20.00	17.0	7.1	44
89	95.50% TECHNICAL MATERIAL	BLUEGILL		1.10	17.0	7.1	40
90	95.50% TECHNICAL MATERIAL	BLUEGILL		1.20	17.0	7.1	40
91	95.50% TECHNICAL MATERIAL	BLUEGILL		0.80	17.0	7.1	40
92	95.50% TECHNICAL MATERIAL	BLUEGILL		1.00	17.0	7.1	40
93	95.50% TECHNICAL MATERIAL	BLUEGILL		0.70	17.0	7.1	40
94	95.50% TECHNICAL MATERIAL	BLUEGILL		0.70	17.0	7.1	40
95	95.50% TECHNICAL MATERIAL	BLUEGILL		0.60	17.0	7.1	40
96	95.50% TECHNICAL MATERIAL	BLUEGILL		1.10	17.0	7.1	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
73	STATIC	LC	1580.	1120.000	2220.000		.	.	1580.	1120.000	2220.000
74	STATIC	LC	6230.	3010.000	12900.000		.	.	3300.	1530.000	7110.000
75	STATIC	LC	7270.	787.000	67000.000		.	.	4610.	2260.000	9380.000
76	STATIC	LC	8180.	5310.000	12600.000		.	.	>3200.	.	.
77	STATIC	LC	9310.	4500.000	19300.000		.	.	6860.	3140.000	15000.000
78	STATIC	LC	5700.	3030.000	10700.000	5570.	2960.000	10500.000	4230.	2830.000	6340.000
79	STATIC	LC	4360.	.	.	4360.	.	.	4360.	3140.000	6060.000
80	STATIC	LC	4910.	1950.000	12400.000	3670.	828.000	16300.000	3420.	749.000	15600.000
81	STATIC	LC	20000.	.	.		.	.	15000.	.	.
82	STATIC	LC	870.	.	.		.	.	192.	114.000	324.000
83	STATIC	LC	1010.	660.000	1540.000		.	.	564.	358.000	884.000
84	STATIC	LC	579.	465.000	721.000		.	.	220.	128.000	416.000
85	STATIC	LC	226.	131.000	392.000		.	.	38.1	19.300	75.000
86	STATIC	LC	484.	211.000	1110.000		.	.	68.	27.000	169.000
87	STATIC	LC	1000.	416.000	2400.000		.	.	180.	138.000	234.000
88	STATIC	LC	1340.	360.000	5000.000		.	.	197.	66.000	590.000
89	STATIC	LC	130.	90.000	180.000	60.	30.000	110.000	<50.	.	.
90	STATIC	LC	180.	110.000	300.000	110.	80.000	140.000	80.	60.000	110.000
91	STATIC	LC	200.	93.700	423.000	101.	58.500	174.000	36.9	22.500	60.500
92	STATIC	LC	284.	87.300	926.000	110.	61.300	198.000	61.7	34.200	111.000
93	STATIC	LC	446.	192.000	1040.000	216.	86.000	540.000	159.	99.800	251.000
94	STATIC	LC		.	.	211.	62.400	716.000	135.	68.800	264.000
95	STATIC	LC	923.	63.100	13500.000	218.	16.800	2840.000	89.4	1.100	894.000
96	STATIC	LC	444.	301.000	657.000	362.	247.000	530.000	81.7	48.600	139.000

ED_005172C_00000035-00102

95 STATIC LC
96 STATIC LC 448.

ANTIMYCIN A
MEASURED IN NG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
97	95.50% TECHNICAL MATERIAL	BLUEGILL		0.90	17.0	7.1	40
98	95.50% TECHNICAL MATERIAL	BLUEGILL		1.50	17.0	7.1	40
99	95.50% TECHNICAL MATERIAL	BLUEGILL		1.20	17.0	7.1	40
100	95.50% TECHNICAL MATERIAL	BLUEGILL		1.10	17.0	7.1	40
101	95.50% TECHNICAL MATERIAL	BLUEGILL		1.40	17.0	7.1	40
102	95.50% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.60	17.0	7.1	44
103	95.50% TECHNICAL MATERIAL	LARGEMOUTH BASS		5.70	17.0	7.1	44
104	95.50% TECHNICAL MATERIAL	LARGEMOUTH BASS		6.10	17.0	7.1	44
105	95.50% TECHNICAL MATERIAL	LARGEMOUTH BASS		11.70	17.0	7.1	44
106	95.50% TECHNICAL MATERIAL	LARGEMOUTH BASS		20.50	17.0	7.1	44
107	95.50% TECHNICAL MATERIAL	WHITE CRAPPIE		1.50	17.0	7.1	44
108	95.50% TECHNICAL MATERIAL	YELLOW PERCH		0.70	12.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
97	STATIC	LC	280.	162.000	481.000	132.	45.400	241.000	103.	43.900	239.000
98	STATIC	LC	328.	133.000	813.000	69.2	31.400	153.000	33.9	8.100	141.000
99	STATIC	LC	280.	190.000	410.000	150.	97.000	230.000	50.	20.000	170.000
100	STATIC	LC	146.	99.800	214.000	64.3	46.300	89.400	44.6	30.400	65.500
101	STATIC	LC	518.	262.000	1030.000	159.	117.000	217.000	94.3	67.300	132.000
102	STATIC	LC	1020.	593.000	1760.000		.	.	237.	159.000	354.000
103	STATIC	LC	369.	160.000	849.000		.	.	>180.	.	.
104	STATIC	LC	>1000.	.	.		.	.	252.	124.000	515.000
105	STATIC	LC	562.	143.000	2220.000		.	.	88.	50.000	153.000
106	STATIC	LC	535.	136.000	2110.000		.	.	63.	40.000	100.000
107	STATIC	LC	420.	356.000	495.000		.	.	340.	273.000	424.000
108	STATIC	LC	54.	42.000	70.000		.	.	40.	31.000	52.000

BEST AVAILABLE COPY

101

ED_005172C_00000035-00103

APHOLATE

CAS # 52-46-0 PRINCIPAL USE: INSECTICIDE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	13.0	7.1	44
2	100.0% TECHNICAL MATERIAL	BLUEGILL		8.50	13.0	7.1	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>40.	:	:		:	:	>40.	:	:
2	STATIC	LC	>30.	:	:		:	:	>30.	:	:

102

AQUA COP

CAS # PRINCIPAL USE: ALGICIDE

MEASURED IN MG /L

ED_005172C_00000035-00104

CAS # PRINCIPAL USE: ALGICIDE

AQUA COP

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	8.00 % LIQUID	RAINBOW TROUT		1.00	12.0	7.4	40
2	8.00 % LIQUID	CHANNEL CATFISH		0.90	22.0	7.6	44
3	8.00 % LIQUID	BLUEGILL		0.80	22.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC		.	.		.	.	24.	19.000	31.000
2	STATIC	LC		.	.		.	.	24.	16.000	36.000
3	STATIC	LC	29.	20.000	42.000		.	.	29.	20.000	42.000

103

ED_005172C_00000035-00105

CAS # PRINCIPAL USE: HERBICIDE

AQUA-VATOR
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	25.00% LIQUID	APPLE SNAIL	2 DAY	.	22.0	7.2	42
2	25.00% LIQUID	RAINBOW TROUT		1.00	12.0	7.4	40
3	25.00% LIQUID	CHANNEL CATFISH		1.00	22.0	7.6	42
4	25.00% LIQUID	BLUEGILL		0.80	22.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC		.	.		.	.	0.25	0.110	0.560
2	STATIC	LC		.	.		.	.	18.	10.000	32.000
3	STATIC	LC	>100.	.	.		.	.	43.	36.000	52.000
4	STATIC	LC		.	.		.	.	92.	69.000	122.000

104

ED_005172C_00000035-00106

ARAMITE

CAS # 140-57-8 PRINCIPAL USE: ACARICIDE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	90.00% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	15.0	7.4	272
2	90.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	15.0	7.4	272
3	90.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	21.0	7.1	44
4	90.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
5	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.90	13.0	7.4	44
6	90.00% TECHNICAL MATERIAL	BLUEGILL		0.70	24.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	160.	113.000	225.000		.	.
2	STATIC	EC		.	.	230.	140.000	390.000		.	.
3	STATIC	LC	350.	250.000	495.000		.	.	60.	43.000	84.000
4	STATIC	LC	>1000.	.	.		.	.	>1000.	.	.
5	STATIC	LC	720.	616.000	842.000		.	.	320.	265.000	387.000
6	STATIC	LC	480.	.	.		.	.	350.	.	.

105

ED_005172C_00000035-00107

ATRAZINE 4L

CAS # 1912-24-9 PRINCIPAL USE: HERBICIDE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	43.00% LIQUID	RAINBOW TROUT		0.70	12.0	7.3	40
2	43.00% LIQUID	BLUEGILL		0.60	22.0	7.1	43

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC		.	.		.	.	24.	18.000	32.000
2	STATIC	LC	48.	42.000	55.000		.	.	42.	36.000	49.000

106

ED_005172C_00000035-00108

AZIDE POTASSIUM
MEASURED IN MG /L

CAS # 20762-60-1 PRINCIPAL USE: HERBICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	98.00% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	15.0	7.1	44
2	98.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	15.0	7.1	44
3	98.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	21.0	7.1	44
4	98.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
5	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	13.0	7.1	44
6	98.00% TECHNICAL MATERIAL	BLUEGILL		0.40	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	7.50	6.300	9.100		.	.
2	STATIC	EC		.	.	8.40	6.100	12.200		.	.
3	STATIC	LC	15.0	11.000	18.000		.	.	6.40	4.600	8.900
4	STATIC	LC	22.0	17.000	29.000		.	.	8.00	5.700	11.000
5	STATIC	LC	1.70	.	.		.	.	1.60	.	.
6	STATIC	LC	2.70	.	.		.	.	0.78	.	.

107

ED_005172C_00000035-00109

CAS # 26628-22-8 PRINCIPAL USE: HERBICIDE

AZIDE SODIUM
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	98.00% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	15.0	7.1	44
2	98.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	15.0	7.1	44
3	98.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	21.0	7.1	44
4	98.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
5	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	13.0	7.1	44
6	98.00% TECHNICAL MATERIAL	BLUEGILL		0.60	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	4.20	2.800	6.200		.	.
2	STATIC	EC		.	.	6.40	4.600	8.900		.	.
3	STATIC	LC	14.0	13.000	15.000		.	.	5.00	3.700	6.800
4	STATIC	LC	16.0	11.000	22.000		.	.	9.00	6.400	13.000
5	STATIC	LC	1.00	.	.		.	.	0.80	.	.
6	STATIC	LC	2.10	.	.		.	.	0.68	.	.

108

ED_005172C_00000035-00110

AZINPHOS-ETHYL
MEASURED IN UG /L

CAS # 2642-71-9 PRINCIPAL USE: INSECTICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	88.00% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	21.0	7.1	44
2	88.00% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	15.0	7.1	44
3	88.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	15.0	7.1	44
4	88.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
5	88.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	13.0	7.1	44
6	88.00% TECHNICAL MATERIAL	BLUEGILL		0.80	24.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	4.0	3.000	5.300		.	.
2	STATIC	EC		.	.	3.2	1.800	5.800		.	.
3	STATIC	EC		.	.	4.2	2.900	6.100		.	.
4	STATIC	LC	>2.	.	.		.	.	1.5	0.800	2.700
5	STATIC	LC	50.	44.000	57.000		.	.	20.	17.000	22.000
6	STATIC	LC	3.7	3.200	4.300		.	.	1.1	0.900	1.200

ED_005172C_00000035-00111

CAS # 86-50-0 PRINCIPAL USE: INSECTICIDE

AZINPHOS-METHYL
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	93.00% TECHNICAL MATERIAL	ASELLUS BREVICAUDUS	MATURE	.	15.0	7.1	44
2	93.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	21.0	7.1	44
3	93.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	272
4	93.00% TECHNICAL MATERIAL	PROCAMBARUS SP.	IMMATURE	.	12.0	7.5	44
5	93.00% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	MATURE	.	21.0	7.4	272
6	93.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
7	93.00% TECHNICAL MATERIAL	COHO SALMON		0.70	12.0	7.5	44
8	93.00% TECHNICAL MATERIAL	COHO SALMON		4.70	12.0	7.1	44
9	93.00% TECHNICAL MATERIAL	COHO SALMON		9.50	12.0	7.1	44
10	93.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	12.0	7.1	44
11	93.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	2.0	7.1	44
12	93.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	7.0	7.1	44
13	93.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	12.0	7.1	44
14	93.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	18.0	7.1	44
15	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.50	12.0	7.5	44
16	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.80	12.0	7.5	44
17	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.80	12.0	7.5	44
18	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.80	12.0	7.5	44
19	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.80	12.0	7.5	44
20	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.50	12.0	7.5	44
21	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON		0.50	12.0	7.5	40
22	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	FINGERLING	.	12.0	7.5	314
23	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	GREEN EGG	.	7.0	6.6	12
24	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	GREEN EGG	.	7.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	162.	117.000	225.000		.	.	21.	15.000	28.000
2	STATIC	LC	0.56	0.390	0.810	0.25	0.190	0.350	0.15	0.110	0.200
3	STATIC	LC	0.44	0.260	0.750	0.16	0.080	0.320	0.10	0.070	0.140
4	STATIC	LC	>200.	.	.		.	.	56.	41.000	77.000
5	FLOW THROUGH	LC	2.5	0.820	3.700		.	.	1.2	.	.
6	STATIC	LC	25.	21.000	29.000	8.0	6.000	10.000	1.9	1.500	2.400
7	STATIC	LC	7.6	5.800	9.900		.	.	6.1	5.000	7.400
8	STATIC	LC	23.	17.000	31.000		.	.	3.2	2.500	4.000
9	STATIC	LC	4.0	2.800	5.700		.	.	3.2	2.400	4.200
10	STATIC	LC	10.	8.000	13.000		.	.	4.3	3.000	6.400
11	STATIC	LC	26.	23.000	29.000		.	.	7.1	6.400	7.900
12	STATIC	LC	16.	15.000	17.000		.	.	5.8	4.800	7.000
13	STATIC	LC	12.	10.000	15.000		.	.	6.3	5.500	7.200
14	STATIC	LC	13.	10.000	16.000		.	.	2.9	2.300	3.700
15	STATIC	LC	>9.0	.	.		.	.	2.1	1.700	2.600
16	STATIC	LC	7.5	5.800	9.700		.	.	2.7	2.200	3.300
17	STATIC	LC	10.	7.800	13.000		.	.	3.2	2.800	3.700
18	STATIC	LC	9.0	7.200	11.300		.	.	3.5	3.000	4.100
19	STATIC	LC	>15.	.	.		.	.	>15.	.	.
20	STATIC	LC	7.6	6.300	9.200		.	.	3.6	3.200	4.100
21	STATIC	LC		.	.		.	.	2.5	2.000	3.100
22	FLOW THROUGH	LC		.	.		.	.	2.5	2.000	3.200
23	STATIC	LC	>50.	.	.		.	.	>50.	.	.
24	STATIC	LC	>50.	.	.		.	.	>50.	.	.

24 STATIC LC

AZINPHOS-METHYL
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
25	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	GREEN EGG	.	7.0	7.8	170
26	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	YOLK-SAC FRY	.	7.0	6.6	12
27	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	YOLK-SAC FRY	.	7.0	7.4	44
28	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	YOLK-SAC FRY	.	7.0	7.8	170
29	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	YOLK-SAC FRY	.	12.0	6.6	12
30	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	YOLK-SAC FRY	.	12.0	7.4	44
31	93.00% TECHNICAL MATERIAL	ATLANTIC SALMON	YOLK-SAC FRY	.	12.0	7.8	170
32	93.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	7.5	44
33	93.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	6.5	44
34	93.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	9.5	44
35	93.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	7.5	12
36	93.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	7.5	170
37	93.00% TECHNICAL MATERIAL	BROWN TROUT		1.50	12.0	7.5	40
38	93.00% TECHNICAL MATERIAL	BROOK TROUT		1.20	12.0	7.5	40
39	93.00% TECHNICAL MATERIAL	NORTHERN PIKE	YOLK-SAC FRY	.	12.0	7.5	44
40	93.00% TECHNICAL MATERIAL	GOLDFISH		0.90	18.0	7.1	44
41	93.00% TECHNICAL MATERIAL	CARP		0.60	18.0	7.1	44
42	93.00% TECHNICAL MATERIAL	FATHEAD MINNOW		1.20	18.0	7.1	44
43	93.00% TECHNICAL MATERIAL	FATHEAD MINNOW		1.20	18.0	7.4	272
44	93.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.80	17.0	7.1	40
45	93.00% TECHNICAL MATERIAL	BLACK BULLHEAD		1.70	18.0	7.1	4[illegible]
46	93.00% TECHNICAL MATERIAL	BLACK BULLHEAD		1.20	18.0	7.4	272
47	93.00% TECHNICAL MATERIAL	BLACK BULLHEAD		1.20	16.0	7.7	135
48	93.00% TECHNICAL MATERIAL	CHANNEL CATFISH		1.50	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
25	STATIC	LC	>50.	.	.		.	.	>50.	.	.
26	STATIC	LC	>15.	.	.		.	.	>15.	.	.
27	STATIC	LC	>15.	.	.		.	.	18.	13.000	25.000
28	STATIC	LC	>15.	.	.		.	.	15.	12.000	19.000
29	STATIC	LC	>15.	.	.		.	.	3.5	.	.
30	STATIC	LC	>15.	.	.		.	.	2.3	.	.
31	STATIC	LC	>15	.	.		.	.	1.8	1.300	2.600
32	STATIC	LC	7.0	5.200	9.400		.	.	4.6	3.500	6.100
33	STATIC	LC	7.0	5.500	8.900		.	.	4.3	3.400	5.500
34	STATIC	LC	8.0	6.900	9.200		.	.	3.5	2.700	4.500
35	STATIC	LC	8.0	5.500	11.700		.	.	6.0	4.900	7.300
36	STATIC	LC	6.0	4.900	7.300		.	.	5.1	4.400	5.900
37	STATIC	LC	7.10	5.560	9.030		.	.	6.60	5.390	8.090
38	STATIC	LC	2.3	1.900	2.700		.	.	1.2	0.870	1.660
39	STATIC	LC	0.67	0.490	0.910		.	.	0.36	0.270	0.480
40	STATIC	LC	7050.	5460.000	9120.000		.	.	4270.	3030.000	6010.000
41	STATIC	LC	1240.	853.000	1770.000		.	.	695.	410.000	1280.000
42	STATIC	LC	255.	150.000	438.000		.	.	235.	135.000	410.000
43	STATIC	LC	524.	277.000	986.000		.	.	293.	110.000	780.000
44	STATIC	LC	160.	119.000	213.000	160.	119.000	213.000	148.	104.000	204.000
45	STATIC	LC	5410.	3960.000	7750.000		.	.	3500.	2920.000	4950.000
46	STATIC	LC	10000.	7420.000	13900.000		.	.	4600.	2730.000	7770.000
47	STATIC	LC	7230.	5400.000	9700.000	7130.	5360.000	9490.000	4810.	3900.000	5910.000
48	STATIC	LC	4530.	3470.000	5900.000		.	.	3290.	2490.000	4340.000

ED_005172C_00000035-00113

AZINPHOS-METHYL
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
49	93.00% TECHNICAL MATERIAL	GREEN SUNFISH		1.10	18.0	7.1	44
50	93.00% TECHNICAL MATERIAL	BLUEGILL		1.50	18.0	7.1	44
51	93.00% TECHNICAL MATERIAL	BLUEGILL		0.90	12.0	7.1	44
52	93.00% TECHNICAL MATERIAL	BLUEGILL		0.90	18.0	7.1	44
53	93.00% TECHNICAL MATERIAL	BLUEGILL		0.90	24.0	7.1	44
54	93.00% TECHNICAL MATERIAL	BLUEGILL		0.50	12.0	6.5	44
55	93.00% TECHNICAL MATERIAL	BLUEGILL		0.50	12.0	8.5	44
56	93.00% TECHNICAL MATERIAL	BLUEGILL		2.20	12.0	7.5	314
57	93.00% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.90	18.0	7.1	44
58	93.00% TECHNICAL MATERIAL	BLACK CRAPPIE		1.00	18.0	7.1	44
59	93.00% TECHNICAL MATERIAL	YELLOW PERCH		1.40	18.0	7.1	44
60	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	7.0	7.5	44
61	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	17.0	7.5	44
62	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	22.0	7.5	44
63	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	6.5	44
64	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	7.5	44
65	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	8.5	44
66	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	9.0	44
67	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	8.0	12
68	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	8.0	44
69	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	8.0	170
70	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	8.0	300
71	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	7.5	44
72	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	7.5	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
49	STATIC	LC	130.	84.000	201.000		.	.	52.	28.000	94.000
50	STATIC	LC	26.	23.000	29.000		.	.	22.	20.000	25.000
51	STATIC	LC	14.	12.000	16.000	11.0	9.300	13.000	8.2	7.000	9.600
52	STATIC	LC	16.	14.000	18.000		.	.	8.0	6.800	9.400
53	STATIC	LC	17.	14.000	20.000	9.0	7.700	10.500	4.1	3.700	4.500
54	STATIC	LC	42.	31.000	56.000		.	.	17.	13.000	23.000
55	STATIC	LC	52.	40.000	68.000		.	.	34.	24.000	48.000
56	FLOW THROUGH	LC		.	.		.	.	4.8	3.700	6.700
57	STATIC	LC	23.	18.000	29.000		.	.	4.8	3.100	7.400
58	STATIC	LC	4.7	3.700	6.000		.	.	3.0	2.300	3.800
59	STATIC	LC	52.	46.000	59.000		.	.	15.	11.000	19.000
60	STATIC	LC	100.	80.000	126.000		.	.	40.	27.000	53.000
61	STATIC	LC	51.	37.000	70.000		.	.	5.6	3.000	11.000
62	STATIC	LC	>15.	.	.		.	.	2.4	1.800	3.300
63	STATIC	LC	54.	45.000	64.000		.	.	17.	12.000	25.000
64	STATIC	LC	>50.	.	.		.	.	29.	23.000	36.000
65	STATIC	LC	70.	56.000	88.000		.	.	8.5	6.000	12.000
66	STATIC	LC	77.	63.000	95.000		.	.	29.	21.000	39.000
67	STATIC	LC	>50.	.	.		.	.	18.	14.000	28.000
68	STATIC	LC	>50.	.	.		.	.	36.	28.000	46.000
69	STATIC	LC	>50.	.	.		.	.	11.	8.600	14.000
70	STATIC	LC	>50.	.	.		.	.	27.	20.000	36.000
71	0 DAY DEGRA	LC	>40.	.	.		.	.	10.	7.000	14.000
72	7 DAY DEGRA	LC	>40.	.	.		.	.	24.	18.000	31.000

24. 18.000 31.000

AZINPHOS-METHYL
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
73	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	7.5	44
74	93.00% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	7.5	44
75	93.00% TECHNICAL MATERIAL	YELLOW PERCH		15.00	12.0	7.5	314
76	93.00% TECHNICAL MATERIAL	FOWLERS TOAD	TADPOLE	.	15.0	7.1	44
77	93.00% TECHNICAL MATERIAL	WESTERN CHORUS FROG	TADPOLE	.	15.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
73	14 DAY DEGRA	LC	>40.	.	.	.	.	.	20.	15.000	26.000
74	21 DAY DEGRA	LC	>40	.	.	.	.	.	33.	19.000	59.000
75	FLOW THROUGH	LC	>4.	.	.	.	.	.	6.5	3.600	11.000
76	STATIC	LC	710.	563.000	895.000	.	.	.	109.	72.000	164.000
77	STATIC	LC	>3200.	.	.	.	.	.	3200.	2243.000	4565.000

ED_005172C_00000035-00115

CAS # PRINCIPAL USE: INSECTICIDE

BACILLUS THURINGIENSIS
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% COMMERCIAL FORMULATION	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
2	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		1.50	13.0	7.1	44
3	100.0% COMMERCIAL FORMULATION	BLUEGILL		1.10	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	80.	.	.		.	.	10.	.	.
2	STATIC	LC	>10.	.	.		.	.	>10.	.	.
3	STATIC	LC	>100.	.	.		.	.	95.	.	.

ED_005172C_00000035-00116

CAS # 101-27-9 PRINCIPAL USE: HERBICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES
1	22.40% EMULSIFIABLE CONCENTRATE	DAPHNIA MAGNA
2	22.40% EMULSIFIABLE CONCENTRATE	CHIRONOMUS PLUMOSUS

BARBAN MEASURED IN MG /L

SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1ST INSTAR	:	17.0	7.2	42
3RD INSTAR	:	22.0	7.2	42

OBS	TEST TYPE		24 HOUR CONFIDENCE INTERVAL		48 HOUR CONFIDENCE INTERVAL			96 HOUR CONFIDENCE INTERVAL	
1	STATIC	EC	:	:	0.43	0.360	0.510	:	:
2	STATIC	EC			0.32	0.230	0.440		

ED_005172C_00000035-00117

BENEFIN

CAS # 1861-40-1 PRINCIPAL USE: HERBICIDE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	IMMATURE	.	15.0	7.1	44
2	100.0% TECHNICAL MATERIAL	GOLDFISH		1.00	18.0	7.1	44
3	100.0% TECHNICAL MATERIAL	FATHEAD MINNOW		0.90	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	8.22	2.360	28.600	4.	2.720	5.820	1.10	0.630	1.930
2	STATIC	LC	0.88	0.720	1.080	0.85	0.680	10.500	0.81	0.650	1.020
3	STATIC	LC		1.000	10.000		.	.		0.100	1.000

116

ED_005172C_00000035-00118

CAS # 17804-35-2 PRINCIPAL USE: FUNGICIDE

BENOMYL

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	99.00% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.4	40
2	99.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	17.0	7.4	40
3	99.00% TECHNICAL MATERIAL	ORCONECTES NAIS	EARLY INSTAR	.	22.0	7.4	40
4	99.00% TECHNICAL MATERIAL	PROCAMBARUS SP.	IMMATURE	.	22.0	7.4	40
5	99.00% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	22.0	7.4	40
6	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	12.0	7.4	44
7	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	7.0	7.4	44
8	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	12.0	7.4	44
9	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	17.0	7.4	44
10	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	12.0	6.5	44
11	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	12.0	7.5	44
12	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	12.0	8.5	44
13	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	12.0	7.4	44
14	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	12.0	7.4	320
15	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	12.0	7.4	44
16	50.00% WETTABLE POWDER	RAINBOW TROUT		1.00	12.0	7.4	44
17	99.00% TECHNICAL MATERIAL	RAINBOW TROUT	FINGERLING	0.30	12.0	7.4	44
18	99.00% TECHNICAL MATERIAL	RAINBOW TROUT	SWIMUP FRY	.	12.0	7.4	44
19	99.00% TECHNICAL MATERIAL	RAINBOW TROUT	YOLK-SAC FRY	.	12.0	7.4	44
20	99.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.90	22.0	7.4	44
21	50.00% WETTABLE POWDER	FATHEAD MINNOW		0.50	22.0	7.4	44
22	99.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.50	22.0	7.1	45
23	50.00% WETTABLE POWDER	FATHEAD MINNOW		0.50	22.0	7.1	45
24	99.00% TECHNICAL MATERIAL	CHANNEL CATFISH		1.20	22.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	2800.	2100.000	3700.000		.	.
2	STATIC	LC		.	.		.	.	750.	500.000	1100.000
3	STATIC	LC	>10000	.	.		.	.	>10000	.	.
4	STATIC	LC	>100000	.	.		.	.	>100000	.	.
5	STATIC	EC		.	.	7000.	5300.000	9200.000		.	.
6	STATIC	LC	560.	380.000	800.000		.	.	170.	120.000	230.000
7	STATIC	LC	700.	530.000	930.000		.	.	170.	110.000	250.000
8	STATIC	LC		.	.		.	.	200.	140.000	280.000
9	STATIC	LC	560.	370.000	840.000		.	.	280.	200.000	390.000
10	STATIC	LC	640.	470.000	870.000		.	.	160.	100.000	240.000
11	STATIC	LC		.	.		.	.	190.	120.000	290.000
12	STATIC	LC	4000.	2800.000	5600.000		.	.	880.	630.000	1240.000
13	STATIC	LC	800.	520.000	1220.000		.	.	230.	170.000	310.000
14	STATIC	LC		.	.		.	.	600.	420.000	850.000
15	STATIC	LC		.	.		.	.	150.	110.000	200.000
16	STATIC	LC		.	.		.	.	310.	250.000	390.000
17	STATIC	LC		.	.		.	.	120.	90.000	160.000
18	STATIC	LC		.	.		.	.	160.	127.000	189.000
19	STATIC	LC		.	.		.	.	280.	220.000	360.000
20	STATIC	LC	3300.	2340.000	4640.000		.	.	2200.	1590.000	3040.000
21	STATIC	LC	2200.	1660.000	2920.000		.	.	1900.	1430.000	2530.000
22	STATIC	LC	1700.	1180.000	2450.000		.	.	1300.	880.000	1920.000
23	STATIC	LC	2200.	1660.000	2920.000		.	.	1900.	1430.000	2530.000
24	STATIC	LC		.	.		.	.	29.	22.000	37.000

ED_005172C_00000035-00119

BENOMYL

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
25	50.00% WETTABLE POWDER	CHANNEL CATFISH		1.20	22.0	7.4	44
26	99.00% TECHNICAL MATERIAL	CHANNEL CATFISH	FINGERLING	0.05	22.0	7.4	44
27	99.00% TECHNICAL MATERIAL	CHANNEL CATFISH	FINGERLING	0.15	22.0	7.4	44
28	99.00% TECHNICAL MATERIAL	CHANNEL CATFISH	SWIMUP FRY	.	22.0	7.4	44
29	99.00% TECHNICAL MATERIAL	CHANNEL CATFISH	YOLK-SAC FRY	.	22.0	7.4	44
30	99.00% TECHNICAL MATERIAL	BLUEGILL		0.90	22.0	7.4	44
31	99.00% TECHNICAL MATERIAL	BLUEGILL		0.90	22.0	7.4	44
32	99.00% TECHNICAL MATERIAL	BLUEGILL		0.90	12.0	7.4	44
33	99.00% TECHNICAL MATERIAL	BLUEGILL		0.90	17.0	7.4	44
34	99.00% TECHNICAL MATERIAL	BLUEGILL		0.90	22.0	7.4	44
35	99.00% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	6.5	44
36	99.00% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	7.5	44
37	99.00% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	8.5	44
38	99.00% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	7.4	44
39	99.00% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	7.4	320
40	50.00% WETTABLE POWDER	BLUEGILL		0.60	22.0	7.4	44
41	99.00% TECHNICAL MATERIAL	BLUEGILL		0.80	22.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
25	STATIC	LC	34.	26.000	44.000	.	.	.	27.5	21.000	34.000
26	STATIC	LC	.	.	.	.	.	.	13.	11.000	15.000
27	STATIC	LC	.	.	.	.	.	.	24.	19.000	29.000
28	STATIC	LC	43.	37.000	50.000	.	.	.	12.	8.800	16.400
29	STATIC	LC	18.	14.000	22.000	.	.	.	5.6	4.600	6.800
30	STATIC	LC	1400	1160.000	1680.000	.	.	.	850.	550.000	1300.000
31	STATIC	LC	800.	517.000	1230.000	.	.	.	560.	410.000	750.000
32	STATIC	LC	.	.	.	.	.	.	750.	560.000	1010.000
33	STATIC	LC	3000.	2100.000	4200.000	.	.	.	1300.	960.000	1760.000
34	STATIC	LC	1600.	1130.000	2260.000	.	.	.	1300.	950.000	1780.000
35	STATIC	LC	1300.	1000.000	1680.000	.	.	.	1300.	1000.000	1680.000
36	STATIC	LC	1300.	815.000	2070.000	.	.	.	1200.	829.000	1736.000
37	STATIC	LC	47000.	29000.000	74000.000	.	.	.	6400.	4185.000	9787.000
38	STATIC	LC	1400.	1030.000	1901.000	.	.	.	1300.	964.000	1752.000
39	STATIC	LC	2900.	2037.000	4128.000	.	.	.	2300.	1702.000	3107.000
40	STATIC	LC	1300.	910.000	1840.000	.	.	.	1200.	900.000	1590.000
41	STATIC	LC	520.	266.000	1010.000	.	.	.	390.	207.000	732.000

BENSULIDE MEASURED IN MG /L

CAS # 741-58-2 PRINCIPAL USE: HERBICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	95.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.1	44
2	95.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.60	13.0	7.4	44
3	95.00% TECHNICAL MATERIAL	BLUEGILL		1.00	24.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	7.60	3.900	15.000	.	.	.	1.40	0.390	5.100
2	STATIC	LC	0.96	.	.	.	.	.	0.72	.	.
3	STATIC	LC	0.97	.	.	.	.	.	0.81	.	.

ED_005172C_00000035-00121

BENTHIOCARB

CAS # 28249-77-6 PRINCIPAL USE: HERBICIDE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
1	95.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	12.0	7.4	44
2	85.20% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		0.60	12.0	7.4	44
3	95.50% TECHNICAL MATERIAL	CHANNEL CATFISH		1.20	20.0	7.4	44
4	85.20% EMULSIFIABLE CONCENTRATE	CHANNEL CATFISH		1.20	20.0	7.4	44
5	95.50% TECHNICAL MATERIAL	BLUEGILL		0.60	20.0	7.4	44
6	85.20% EMULSIFIABLE CONCENTRATE	BLUEGILL		0.60	20.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	1960.	1720.000	2240.000	.	.	.	1150.	738.000	1780.000
2	STATIC	LC	1340.	1160.000	1700.000	.	.	.	1050.	690.000	1590.000
3	STATIC	LC	8190.	6700.000	10000.000	.	.	.	2280.	1190.000	4380.000
4	STATIC	LC	4250.	3170.000	5690.000	.	.	.	2290.	1190.000	4380.000
5	STATIC	LC	5790.	3430.000	9770.000	.	.	.	2480.	1650.000	3710.000
6	STATIC	LC	2490.	1780.000	3470.000	.	.	.	1660.	1220.000	2260.000

130

ED_005172C_00000035-00122

BENZALKONIUM CHLORIDE
MEASURED IN UG /L

CAS # 8001-54-5 PRINCIPAL USE: DISINFECTANT

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	10.00% LIQUID	RAINBOW TROUT		1.80	12.0	7.[illegible]	44
2	10.00% LIQUID	BLUEGILL		0.30	24.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	1300.	990.000	1710.000	.	.	.	1150.	823.000	1610.000
2	STATIC	LC	580.	390.000	867.000	.	.	.	320.	223.000	460.000

121

ED_005172C_00000035-00123

BENZENE

CAS # 71-43-2 PRINCIPAL USE: SOLVENT MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% LIQUID	RAINBOW TROUT		2.40	12.0	7.4	44
2	100.0% LIQUID	CHANNEL CATFISH		0.10	22.0	7.4	44
3	100.0% LIQUID	BLUEGILL		0.20	17.0	7.4	44
4	100.0% LIQUID	BLUEGILL		0.20	22.0	7.4	44
5	100.0% LIQUID	BLUEGILL		0.20	19.0	6.5	44
6	100.0% LIQUID	BLUEGILL		0.20	19.0	7.4	44
7	100.0% LIQUID	BLUEGILL		0.20	19.0	8.5	44
8	100.0% LIQUID	BLUEGILL		0.50	17.0	8.0	12
9	100.0% LIQUID	BLUEGILL		0.50	17.0	8.0	44
10	100.0% LIQUID	BLUEGILL		0.50	17.0	8.0	162
11	100.0% LIQUID	BLUEGILL		0.50	17.0	8.0	300

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	9.2	7.200	11.700	.	.	.	9.2	7.200	11.700
2	STATIC	LC	425.	357.000	505.000	.	.	.	425.	357.000	505.000
3	STATIC	LC	400.	285.000	570.000	.	.	.	230.	181.000	291.000
4	STATIC	LC	140.	107.000	183.000	.	.	.	100.	70.000	142.000
5	STATIC	LC	910.	654.000	1267.000	.	.	.	600.	481.000	748.000
6	STATIC	LC	740.	584.000	937.000	.	.	.	450.	349.000	580.000
7	STATIC	LC	580.	343.000	981.000	.	.	.	290.	178.000	473.000
8	STATIC	LC	370.	285.000	481.000	.	.	.	370.	285.000	481.000
9	STATIC	LC	260.	213.000	311.000	.	.	.	260.	213.000	311.000
10	STATIC	LC	102.	86.000	120.000	.	.	.	102.	86.000	120.000
11	STATIC	LC	165.	116.000	234.000	.	.	.	165.	116.000	234.000

122

ED_005172C_00000035-00124

CAS # 608-73-1 PRINCIPAL USE: INSECTICIDE

BENZENE HEXACHLORIDE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	41.50% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	16.0	7.1	44
2	41.50% TECHNICAL MATERIAL	GAMMARUS LACUSTRIS	MATURE	.	21.0	7.1	44
3	41.50% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
4	41.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00	13.0	7.1	44
5	41.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00	13.0	7.4	272
6	41.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	13.0	7.4	272
7	41.50% TECHNICAL MATERIAL	GOLDFISH		1.00	18.0	7.1	44
8	41.50% TECHNICAL MATERIAL	FATHEAD MINNOW		1.00	18.0	7.1	44
9	41.50% TECHNICAL MATERIAL	CHANNEL CATFISH		1.10	18.0	7.1	44
10	41.50% TECHNICAL MATERIAL	BLUEGILL		1.50	18.0	7.1	44
11	41.50% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.80	18.0	7.1	44
12	41.50% TECHNICAL MATERIAL	FOWLERS TOAD	TADPOLE	.	15.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	680.	.	.		.	.
2	STATIC	LC	450.	330.000	600.000		.	.	78.	54.000	113.000
3	STATIC	LC	<18.	.	.		.	.	<18.	.	.
4	STATIC	LC	9.6	8.400	11.000		.	.	9.0	8.000	10.000
5	STATIC	LC	43.	41.000	45.000		.	.	35.	33.000	37.000
6	STATIC	LC	20.	17.000	23.000		.	.	18.	15.000	20.000
7	STATIC	LC	664.	531.000	832.000		.	.	348.	261.000	466.000
8	STATIC	LC	1000.	.	.		.	.	125.	96.000	164.000
9	STATIC	LC	150.	124.000	181.000		.	.	105.	85.000	129.000
10	STATIC	LC	148.	120.000	182.000		.	.	67.	51.000	96.000
11	STATIC	LC	44.	41.000	48.000		.	.	41.	36.000	47.000
12	STATIC	LC	13200.	4320.000	40300.000		.	.	4350	2990.000	6310.000

133

ED_005172C_00000035-00125

BENZOTHIAZOLE 2-MERCAPTO
MEASURED IN MG /L

CAS # 149-30-4 PRINCIPAL USE: INDUSTRIAL

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	12.0	7.5	41
2	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		1.00	22.0	7.3	42
3	100.0% TECHNICAL MATERIAL	BLUEGILL		0.80	22.0	7.5	41
4	100.0% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	6.5	40
5	100.0% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	7.5	40
6	100.0% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	8.5	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	0.76	0.600	0.960	.	.		0.76	0.600	0.960
2	STATIC	LC	2.35	1.800	3.060	.	.		1.65	1.160	2.350
3	STATIC	LC	2.25	29.600	1.710	.	.		2.25	2.960	1.710
4	STATIC	LC	2.35	1.830	3.020	.	.		1.90	1.320	2.730
5	STATIC	LC	4.70	3.540	6.240	.	.		4.20	3.310	5.320
6	STATIC	LC	>10.	.	.	.	.		7.6	5.940	9.720

124

ED_005172C_00000035-00126

CAS # 64036-43-7 PRINCIPAL USE: INDUSTRIAL

BENZOTHIAZOLE 2-METHYL MERCAPTO
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	16.0	7.5	44
2	100.0% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	17.0	7.5	44
3	100.0% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	22.0	7.5	44
4	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		0.60	12.0	7.5	41
5	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		1.00	22.0	7.3	42
6	100.0% TECHNICAL MATERIAL	BLUEGILL		0.80	22.0	7.5	41
7	100.0% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	6.5	44
8	100.0% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	7.5	44
9	100.0% TECHNICAL MATERIAL	BLUEGILL		0.60	22.0	8.5	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	>10.	.	.		.	.
2	STATIC	LC		.	.		.	.	4.2	3.500	5.000
3	STATIC	EC		.	.	1.3	0.900	1.900		.	.
4	STATIC	LC	7.60	5.940	9.720		.	.	3.70	2.850	4.810
5	STATIC	LC	13.0	10.200	16.600		.	.	7.60	5.940	9.720
6	STATIC	LC	13.5	10.600	17.300		.	.	5.70	4.050	8.030
7	STATIC	LC	13.5	10.600	17.200		.	.	4.2	3.300	5.300
8	STATIC	LC	10.5	7.300	15.100		.	.	4.3	3.400	5.400
9	STATIC	LC	10.5	7.300	15.100		.	.	6.3	4.700	8.400

ED_005172C_00000035-00127

CAS # 33878-50-1 PRINCIPAL USE: HERBICIDE

BENZOYLPROP ETHYL
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	19.00% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		1.10	12.0	7.4	44
2	19.00% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		1.10	12.0	7.4	44
3	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	12.0	7.4	44
4	19.00% EMULSIFIABLE CONCENTRATE	FATHEAD MINNOW		0.50	17.0	7.4	44
5	25.00% EMULSIFIABLE CONCENTRATE	FATHEAD MINNOW		0.50	17.0	7.4	44
6	100.0% TECHNICAL MATERIAL	FATHEAD MINNOW		0.50	17.0	7.4	44
7	19.00% EMULSIFIABLE CONCENTRATE	BLUEGILL		0.80	17.0	7.4	44
8	25.00% EMULSIFIABLE CONCENTRATE	BLUEGILL		0.80	17.0	7.4	44
9	100.0% TECHNICAL MATERIAL	BLUEGILL		0.80	17.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	560.	341.000	920.000	.	.		<32.	.	.
2	STATIC	LC	1300.	1100.000	1540.000	.	.		<32.	.	.
3	STATIC	LC	1100.	841.000	1439.000	.	.		.	.	.
4	STATIC	LC	3000.	2187.000	4114.000	.	.		460.	330.000	640.000
5	STATIC	LC	3500.	.	.	.	.		760.	525.000	1100.000
6	STATIC	LC	4000.	2901.000	5514.000	.	.		860.	660.000	1120.000
7	STATIC	LC	1080.	923.000	1262.000	.	.		1000.	808.000	1238.000
8	STATIC	LC	1100.	941.000	1286.000	.	.		720.	554.000	919.000
9	STATIC	LC	1750.	1570.000	1950.000	.	.		1200.	1073.000	1341.000

126

ED_005172C_00000035-00128

BENZYL BENZOATE
MEASURED IN MG /L

CAS # 120-51-4 PRINCIPAL USE: ACARICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	99.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	16.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	9.8	.	.	.	.	.	4.8	.	.

127

ED_005172C_00000035-00129

BINAPACRYL

CAS # 485-31-4 PRINCIPAL USE: ACARICIDE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	99.90% TECHNICAL MATERIAL	ASELLUS BREVICAUDUS	MATURE	.	16.0	7.4	272
2	99.90% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	13.0	7.1	44
3	99.90% TECHNICAL MATERIAL	CHANNEL CATFISH		1.40	18.0	7.1	44
4	99.90% TECHNICAL MATERIAL	GREEN SUNFISH		0.80	18.0	7.1	44
5	99.90% TECHNICAL MATERIAL	BLUEGILL		1.30	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	130.	.	.	.	.	.	29.	.	.
2	STATIC	LC	50.	46.000	55.000	.	.	.	50.	46.000	55.000
3	STATIC	LC	62.	39.000	99.000	.	.	.	15.	.	.
4	STATIC	LC	53.	44.000	63.000	.	.	.	43.	37.000	50.000
5	STATIC	LC	42.	34.000	51.000	.	.	.	40.	34.000	47.000

ED_005172C_00000035-00130

BLACK CRUDE OIL
MEASURED IN MG /L

CAS # 8002-05-9 PRINCIPAL USE: PETROLEUM

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% WHOLE MATERIAL	CUTTHROAT TROUT		0.70	12.0	7.4	162
2	100.0% WATER SOLUBLE	CUTTHROAT TROUT		0.70	12.0	7.4	162
3	100.0% WHOLE MATERIAL	CUTTHROAT TROUT	EYED EGG	.	10.0	7.2	170
4	100.0% WHOLE MATERIAL	CUTTHROAT TROUT	SWIMUP FRY	.	10.0	7.2	170
5	100.0% WHOLE MATERIAL	CUTTHROAT TROUT	YOLK-SAC FRY	.	10.0	7.2	170

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL	48 HOUR	CONFIDENCE INTERVAL	96 HOUR	CONFIDENCE INTERVAL
1	STATIC	LC	.	.	.	.	>100.	. .
2	STATIC	LC	.	.	.	.	>11.3	. .
3	STATIC	LC	.	.	.	.	>100.	. .
4	STATIC	LC	.	.	.	.	>100.	. .
5	STATIC	LC	.	.	.	.	>100.	. .

129

ED_005172C_00000035-00131

CAS # 122-10-1 PRINCIPAL USE: INSECTICIDE

BOHYL

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	50.00% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		0.60	18.0	7.1	44
2	25.00% WETTABLE POWDER	RAINBOW TROUT		0.60	18.0	7.1	44
3	94.00% TECHNICAL MATERIAL	BLUEGILL		0.40	24.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	0.17	.	.		.	.		.	.
2	STATIC	LC	0.10	.	.		.	.		.	.
3	STATIC	LC	0.60	.	.		.	.	0.51	.	.

130

ED_005172C_00000035-00132

BUFENCARB

MEASURED IN UG /L

CAS # 2282-34-0 PRINCIPAL USE: INSECTICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	72.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	44
2	72.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	272
3	72.00% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	MATURE	.	21.0	7.4	272
4	72.00% TECHNICAL MATERIAL	GOLDFISH		1.00	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	5.3	3.700	7.400	.	.	.	1.0	0.600	1.600
2	STATIC	LC	3.2	.	.	.	.	.	1.0	.	.
3	STATIC	LC	40.	16.000	100.000	.	.	.	0.96	0.590	1.600
4	STATIC	LC	322.	239.000	434.000	.	.	.	294.	222.000	389.000

131

ED_005172C_00000035-00133

CAS # 9003-13-8 PRINCIPAL USE: FLY REPELLENT

BUTOXY POLYPROPYLENE GLYCOL
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE		15.0	7.4	4[illegible]

OBS	TEST TYPE		24 HOUR CONFIDENCE INTERVAL		48 HOUR CONFIDENCE INTERVAL		96 HOUR CONFIDENCE INTERVAL	
1	STATIC	LC					17.000	7.060	40.900

ED_005172C_00000035-00134

CAS # 2008-41-5 PRINCIPAL USE: HERBICIDE

BUTYLATE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
1	98.70% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	262
2	98.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	44
3	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.30	12.0	7.4	44
4	98.00% TECHNICAL MATERIAL	BLUEGILL		1.20	24.0	7.4	44
5	98.00% TECHNICAL MATERIAL	BLUEGILL		1.10	24.0	7.4	44
6	98.00% TECHNICAL MATERIAL	BLUEGILL		1.00	24.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	26000.	20000.000	34000.000	.	.	.	11000.	7600.000	16000.000
2	STATIC	LC	17000.	10500.000	27500.000	.	.	.	12000.	8320.000	17300.000
3	STATIC	LC	4000.	3050.000	5250.000	.	.	.	2100.	1180.000	3740.000
4	STATIC	LC	5000.	2340.000	9850.000	.	.	.		300.000	700.000
5	STATIC	LC	4000.	2180.000	7340.000	.	.	.	210.	88.000	500.000
6	STATIC	LC	900.	700.000	1160.000	.	.	.	470.	.	.

ED_005172C_00000035-00135

CAS # 124-65-2 PRINCIPAL USE: HERBICIDE

CACODYLIC ACID
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	272
2	100.0% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	44
3	100.0% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	MATURE	.	21.0	7.4	272
4	100.0% TECHNICAL MATERIAL	BLUEGILL		0.80	18.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC		.	.	.	.	.	135.	80.000	227.000
2	STATIC	LC		.	.	.	.	.	140.	40.000	486.000
3	STATIC	LC		.	.	.	.	.	28.	14.000	58.000
4	STATIC	LC	21.	19.000	23.000	.	.	.	17.	15.000	19.000

134

ED_005172C_00000035-00136

CALCIUM HYPOCHLORITE
MEASURED IN UG /L

CAS # 7778-54-3 PRINCIPAL USE: DISINFECTANT

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% WETTABLE POWDER	FATHEAD MINNOW		0.90	18.0	7.1	44
2	100.0% WETTABLE POWDER	FATHEAD MINNOW		0.90	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	1120.	817.000	1536.000	.	.	.	840.	641.000	1101.000
2	STATIC	LC	1170.	376.000	3660.000	.	.	.	889.	561.000	1410.000

ED_005172C_00000035-00137

CAPTAFOL

CAS # 2425-06-1 PRINCIPAL USE: FUNGICIDE MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	80.00% WETTABLE POWDER	GAMMARUS LACUSTRIS	MATURE	.	21.0	7.1	44
2	80.00% WETTABLE POWDER	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
3	80.00% WETTABLE POWDER	RAINBOW TROUT		1.00	13.0	7.1	44
4	80.00% WETTABLE POWDER	RAINBOW TROUT		1.00	13.0	7.4	272
5	80.00% WETTABLE POWDER	GOLDFISH		0.90	18.0	7.1	44
6	80.00% WETTABLE POWDER	GOLDFISH		0.90	18.0	7.4	272
7	80.00% WETTABLE POWDER	CHANNEL CATFISH		1.00	18.0	7.1	44
8	80.00% WETTABLE POWDER	BLUEGILL		1.10	18.0	7.1	44
9	80.00% WETTABLE POWDER	BLUEGILL		1.10	18.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	2000.	1700.000	2800.000	.	.	.	800.	500.000	1300.000
2	STATIC	LC	480.	370.000	660.000	.	.	.	40.0	20.000	60.000
3	STATIC	LC	30.2	19.500	46.500	.	.	.	20.6	13.300	31.700
4	STATIC	LC	49.4	32.000	76.400	.	.	.	41.2	30.600	55.100
5	STATIC	LC	189.	156.000	229.000	.	.	.	178.	146.000	217.000
6	STATIC	LC	350.	258.000	474.000	.	.	.	350.	258.000	474.000
7	STATIC	LC	36.	.	.	.	.	.	28.	.	.
8	STATIC	LC	34.0	21.000	54.000	.	.	.	34.	21.000	54.000
9	STATIC	LC	59.0	49.000	70.000	.	.	.	59.0	49.000	70.000

CAPTAN MEASURED IN UG /L

ED_005172C_00000035-00138

CAS # 133-06-2 PRINCIPAL USE: FUNGICIDE

CAPTAN

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
1	90.00% TECHNICAL MATERIAL	COHO SALMON		0.80	12.0	7.5	44
2	90.00% TECHNICAL MATERIAL	COHO SALMON		0.80	12.0	7.5	314
3	90.00% TECHNICAL MATERIAL	CHINOOK SALMON	FINGERLING	.	12.0	7.5	44
4	90.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.40	12.0	7.4	44
5	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	12.0	7.4	44
6	90.00% TECHNICAL MATERIAL	BROWN TROUT		0.70	12.0	7.5	44
7	90.00% TECHNICAL MATERIAL	BROWN TROUT		0.60	12.0	7.5	314
8	90.00% TECHNICAL MATERIAL	LAKE TROUT		0.42	12.0	7.5	44
9	90.00% TECHNICAL MATERIAL	LAKE TROUT		2.30	12.0	7.4	162
10	90.00% TECHNICAL MATERIAL	LAKE TROUT	FINGERLING	.	12.0	7.5	314
11	90.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.30	12.0	7.5	44
12	90.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.40	12.0	7.5	314
13	90.00% TECHNICAL MATERIAL	CHANNEL CATFISH		1.20	20.0	7.4	44
14	90.00% TECHNICAL MATERIAL	BLUEGILL		1.10	17.0	7.1	44
15	90.00% TECHNICAL MATERIAL	YELLOW PERCH		1.00	17.0	7.5	314

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	137.	117.000	160.000	.	.	.	137.	117.000	160.000
2	FLOW THROUGH	LC	75.0	64.900	85.500	.	.	.	56.5	52.300	61.000
3	STATIC	LC	139.	115.000	168.000	.	.	.	120.	103.000	140.000
4	STATIC	LC	74.1	55.400	99.100	.	.	.	56.4	42.200	75.400
5	STATIC	LC	76.4	69.500	83.900	.	.	.	73.2	66.600	80.400
6	STATIC	LC	81.0	69.800	94.000	.	.	.	80.0	63.800	100.000
7	FLOW THROUGH	LC	26.2	21.800	31.300	.	.	.	26.2	21.800	31.300
8	STATIC	LC	53.0	44.500	63.100	.	.	.	49.0	40.100	59.900
9	STATIC	LC	63.2	49.600	80.500	.	.	.	63.2	49.600	80.500
10	FLOW THROUGH	LC	75.0	51.700	109.000	.	.	.	51.0	39.200	66.200
11	STATIC	LC	290.	211.000	398.000	.	.	.	200.	168.000	238.000
12	FLOW THROUGH	LC	152.	123.000	186.000	.	.	.	134.	100.000	178.000
13	STATIC	LC	79.8	72.600	87.800	.	.	.	77.5	70.500	85.200
14	STATIC	LC	145.	122.000	172.000	.	.	.	141.	119.000	167.000
15	FLOW THROUGH	LC	>154.	.	.	.	.	.	120.	97.300	147.000

ED_005172C_00000035-00139

CAS # 63-25-2 PRINCIPAL USE: INSECTICIDE

CARBARYL

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
1	99.50% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	16.0	7.4	42
2	99.50% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	16.0	7.1	44
3	99.50% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	10.0	7.1	44
4	99.50% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	16.0	7.1	44
5	99.50% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	21.0	7.1	44
6	99.50% TECHNICAL MATERIAL	CYPRIDOPSIS VIDUA	MATURE	.	21.0	7.4	272
7	99.50% TECHNICAL MATERIAL	ASELLUS BREVICAUDUS	MATURE	.	18.0	7.1	44
8	99.50% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	21.0	7.1	44
9	99.50% TECHNICAL MATERIAL	GAMMARUS LACUSTRIS	MATURE	.	21.0	7.1	44
10	99.50% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	6.5	40
11	99.50% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	6.5	40
12	99.50% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	7.5	40
13	99.50% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	12.0	8.5	40
14	99.50% TECHNICAL MATERIAL	PROCAMBARUS SP.	IMMATURE	.	12.0	7.5	42
15	99.50% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	MATURE	.	21.0	7.4	272
16	99.50% TECHNICAL MATERIAL	CLAASSENIA SABULOSA	2ND YEAR CLASS	.	16.0	7.1	44
17	99.50% TECHNICAL MATERIAL	ISOGENUS SP.	1ST YEAR CLASS	.	7.0	7.0	35
18	99.50% TECHNICAL MATERIAL	ISOGENUS SP.	1ST YEAR CLASS	.	12.0	7.5	42
19	99.50% TECHNICAL MATERIAL	ISOGENUS SP.	1ST YEAR CLASS	.	12.0	7.5	42
20	99.50% TECHNICAL MATERIAL	ISOGENUS SP.	1ST YEAR CLASS	.	12.0	7.5	42
21	99.50% TECHNICAL MATERIAL	ISOGENUS SP.	1ST YEAR CLASS	.	12.0	7.5	42
22	49.00% OIL DISPERSION	ISOGENUS SP.	1ST YEAR CLASS	.	7.0	7.5	42
23	99.50% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	.	16.0	7.1	44
24	99.50% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	.	12.0	6.5	38

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	5.6	2.700	12.000		.	.
2	STATIC	LC		.	.	6.4	4.500	8.900		.	.
3	STATIC	EC		.	.	11.	7.800	16.000		.	.
4	STATIC	EC		.	.	7.6	6.200	9.300		.	.
5	STATIC	EC		.	.	8.1	6.700	9.800		.	.
6	STATIC	EC	1920.	994.000	3709.000	115.0	74.000	179.000		.	.
7	STATIC	LC	340.	216.000	534.000		.	.	280.	214.000	367.000
8	STATIC	LC	50.	32.000	70.000		.	.	26.	16.000	39.000
9	STATIC	LC	40.	32.000	49.000		.	.	22.	16.000	30.000
10	STATIC	LC	19.0	14.400	25.000	13.0	8.890	19.010		.	.
11	STATIC	LC	12.6	8.510	18.600	8.0	4.930	13.000		.	.
12	STATIC	LC	22.0	14.300	33.800	11.5	7.510	17.600	7.0	4.120	11.900
13	STATIC	LC	10.5	6.640	16.600	7.8	5.950	10.200	7.2	5.600	9.260
14	STATIC	LC	>4000.	.	.		.	.	1900.	1161.000	3108.000
15	STATIC	LC	120.	100.000	140.000		.	.	5.6	3.600	8.300
16	STATIC	LC	12.	9.100	16.000		.	.	5.6	3.900	8.100
17	STATIC	LC	8.0	5.300	12.000		.	.	2.8	2.000	4.000
18	0 DAY DEGRA	LC	8.7	4.900	9.200		.	.	3.6	2.400	5.500
19	7 DAY DEGRA	LC	8.0	6.300	10.000		.	.	6.6	5.000	8.900
20	14 DAY DEGRA	LC	9.	6.900	12.000		.	.	6.6	5.300	8.300
21	21 DAY DEGRA	LC	15.	11.000	21.000		.	.	12.	9.700	15.000
22	STATIC	LC	15.	8.900	24.000		.	.	9.2	7.400	12.000
23	STATIC	LC	5.0	3.600	7.000		.	.	1.7	1.400	2.400
24	STATIC	LC	19.	16.000	22.000		.	.	11.	9.700	13.000

23	STATIC	LC	[illegible]	[illegible]	[illegible]
24	STATIC	LC	19.	16.000	22.00

CARBARYL
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT		DEG. CENT.	PH	HARDNESS MG/L
25	99.50% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	.		12.0	7.5	38
26	99.50% TECHNICAL MATERIAL	PTERONARCELLA BADIA	1ST YEAR CLASS	.		12.0	8.5	36
27	99.50% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	1ST YEAR CLASS	.		16.0	7.1	44
28	99.50% TECHNICAL MATERIAL	COHO SALMON		1.00		13.0	7.1	44
29	99.50% TECHNICAL MATERIAL	COHO SALMON		4.60		13.0	7.1	42
30	99.50% TECHNICAL MATERIAL	COHO SALMON		5.10		13.0	7.1	42
31	99.50% TECHNICAL MATERIAL	COHO SALMON		10.10		13.0	7.1	42
32	99.50% TECHNICAL MATERIAL	COHO SALMON		19.10		13.0	7.1	42
33	99.00% TECHNICAL MATERIAL	CHINOOK SALMON	FINGERLING	.		12.0	7.5	314
34	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50		12.0	7.5	40
35	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.60		7.0	7.5	42
36	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.70		12.0	8.5	40
37	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.60		12.0	8.5	40
38	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50		12.0	7.8	330
39	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50		12.0	7.3	40
40	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50		12.0	7.3	40
41	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50		12.0	7.2	42
42	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50		12.0	7.2	42
43	99.50% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.90		12.0	7.5	42
44	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.50		12.0	7.1	42
45	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.50		12.0	7.4	272
46	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	FISH PROTEIN CONCENTRATE DIET	10.0	7.4	40
47	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	EWOS DIET	10.0	7.4	40
48	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	GLENCO DIET	10.0	7.4	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
25	STATIC	LC	18.	14.000	24.000	.	.	.	13.	12.000	16.000
26	STATIC	LC	32.	26.000	36.000	.	.	.	29.	21.000	41.000
27	STATIC	LC	30.	22.000	40.000	.	.	.	4.8	3.000	7.700
28	STATIC	LC	6520.	5040.000	8260.000	.	.	.	4340.	3310.000	5690.000
29	STATIC	LC	4300.	2766.000	6685.000	.	.	.	2400.	1865.000	3088.000
30	STATIC	LC	2400.	1865.000	3088.000	.	.	.	1750.	1258.000	2434.000
31	STATIC	LC	3100.	2311.000	4158.000	.	.	.	2700.	1570.000	4643.000
32	STATIC	LC	1350.	949.000	1920.000	.	.	.	1150.	845.000	1564.000
33	FLOW THROUGH	LC		.	.	.	.	.	2400.	1623.000	3549.000
34	STATIC	LC	7400.	5700.000	9600.000	.	.	.	7100.	5240.000	9620.000
35	STATIC	LC	7300.	5750.000	9270.000	.	.	.	6000.	4630.000	7770.000
36	STATIC	LC	5700.	4870.000	6670.000	.	.	.	5000.	4100.000	6100.000
37	STATIC	LC	2750.	2340.000	3230.000	.	.	.	970.	770.000	1220.000
38	STATIC	LC	7300.	5190.000	10300.000	.	.	.	3950.	3370.000	4630.000
39	0 DAY DEGRA	LC	6800.	5230.000	8840.000	.	.	.	6800.	5230.000	8840.000
40	7 DAY DEGRA	LC	>6000.	.	.	.	.	.	4000.	3130.000	5110.000
41	14 DAY DEGRA	LC	>3600.	.	.	.	.	.	3380.	2300.000	4960.000
42	21 DAY DEGRA	LC	>3600.	.	.	.	.	.	2300.	1830.000	2890.000
43	STATIC	LC	6700.	5460.000	8220.000	.	.	.	6700.	5230.000	8580.000
44	STATIC	LC	1990.	1480.000	2660.000	.	.	.	1950.	1450.000	2630.000
45	STATIC	LC	2100.	1240.000	3520.000	.	.	.	1200.	800.000	1800.000
46	STATIC	LC		.	.	.	.	.	1360.	992.000	1860.000
47	STATIC	LC		.	.	.	.	.	2080.	1580.000	2730.000
48	STATIC	LC		.	.	.	.	.	1900.	1590.000	2263.000

ED_005172C_00000035-00141

CARBARYL

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT		DEG. CENT.	PH	HARDNESS MG/L
49	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	FPRL 50 PERCENT PROTEIN DIET	10.0	7.4	40
50	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.50		12.0	7.5	314
51	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		0.80	DONALDSON STRAIN	12.0	7.1	40
52	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	IOWA STRAIN	12.0	7.1	40
53	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	MISSOURI STRAIN	12.0	7.1	40
54	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.80	NEW HAMPSHIRE STRAIN	12.0	7.1	40
55	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	SOAP LAKE STRAIN	12.0	7.1	40
56	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		2.20	WYTHEVILLE STRAIN	12.0	7.1	40
57	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20		12.0	7.4	40
58	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.20		12.0	7.4	320
59	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00		12.0	7.4	40
60	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00		17.0	7.4	40
61	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00		12.0	6.5	40
62	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00		12.0	7.5	40
63	99.50% TECHNICAL MATERIAL	RAINBOW TROUT		1.00		12.0	8.5	40
64	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40		12.0	7.5	42
65	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.80		7.0	7.5	42
66	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.80		17.0	7.5	42
67	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40		12.0	6.5	42
68	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.40		12.0	8.5	42
69	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.80		12.0	8.0	12
70	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.80		12.0	8.0	42
71	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20		12.0	6.5	42
72	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20		12.0	7.5	42

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
49	STATIC	LC		.	.		.	.	2300.	1500.000	3500.000
50	FLOW THROUGH	LC	1340.	1039.000	1729.000		.	.	1330.	1030.000	1716.000
51	STATIC	LC	6220.	2460.000	17700.000		.	.	<750.	.	.
52	STATIC	LC	2760.	1890.000	4040.000		.	.	<320.	.	.
53	STATIC	LC	3940.	3090.000	5020.000		.	.	1090.	.	.
54	STATIC	LC	4620.	2470.000	8640.000		.	.		.	.
55	STATIC	LC	4600.	3020.000	6990.000		.	.	1460.	1010.000	2110.000
56	STATIC	LC	6090.	2480.000	15000.000		.	.		.	.
57	STATIC	LC	4700.	3320.000	6660.000		.	.	3500.	2450.000	4990.000
58	STATIC	LC	8200.	6660.000	10100.000		.	.	3000.	2080.000	4320.000
59	STATIC	LC	7500.	5500.000	10200.000		.	.	1600.	1070.000	2370.000
60	STATIC	LC	1500.	1000.000	2230.000		.	.	1100.	760.000	1500.000
61	STATIC	LC	1650.	1110.000	2450.000		.	.	1200.	806.000	1790.000
62	STATIC	LC	3000.	2320.000	3878.000		.	.	780.	649.000	937.000
63	STATIC	LC	2400.	2115.000	2722.000		.	.	1450.	1169.000	1797.000
64	STATIC	LC	4900.	4110.000	5840.000		.	.	4500.	3820.000	5310.000
65	STATIC	LC	2600.	2240.000	3010.000		.	.	2070.	1850.000	2310.000
66	STATIC	LC	1900.	1640.000	2200.000		.	.	1180.	988.000	1410.000
67	STATIC	LC	1400.	.	.		.	.	1270.	902.000	1790.000
68	STATIC	LC	905.	569.000	1440.000		.	.	905.	569.000	1440.000
69	STATIC	LC	2100.	1620.000	2720.000		.	.	2010.	1540.000	2630.000
70	STATIC	LC	2050.	1680.000	2510.000		.	.	1430.	1060.000	1930.000
71	STATIC	LC	700.	474.000	1033.000		.	.	500.	352.000	710.000
72	STATIC	LC	1300.	997.000	1690.000		.	.	1000.	754.000	1420.000

CARBARYL

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	HARDNESS PH	HARDNESS MG/L
73	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	8.5	42
74	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	8.0	12
75	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	8.0	42
76	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	7.0	7.5	42
77	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	12.0	7.5	42
78	99.50% TECHNICAL MATERIAL	ATLANTIC SALMON		0.20	17.0	7.5	42
79	99.50% TECHNICAL MATERIAL	BROWN TROUT		0.60	12.0	7.5	42
80	99.50% TECHNICAL MATERIAL	BROWN TROUT	FINGERLING	.	12.0	7.5	314
81	99.50% TECHNICAL MATERIAL	BROOK TROUT		0.80	12.0	7.5	42
82	99.50% TECHNICAL MATERIAL	BROOK TROUT		0.80	7.0	7.5	42
83	99.50% TECHNICAL MATERIAL	BROOK TROUT		1.00	17.0	7.5	42
84	99.50% TECHNICAL MATERIAL	BROOK TROUT		0.70	12.0	6.5	42
85	99.50% TECHNICAL MATERIAL	BROOK TROUT		0.70	12.0	8.5	42
86	99.50% TECHNICAL MATERIAL	BROOK TROUT		0.70	12.0	9.0	42
87	99.50% TECHNICAL MATERIAL	BROOK TROUT		0.80	12.0	7.5	42
88	99.50% TECHNICAL MATERIAL	BROOK TROUT		0.80	12.0	7.5	300
89	49.00% OIL DISPERSION	BROOK TROUT		1.30	12.0	7.5	42
90	99.50% TECHNICAL MATERIAL	LAKE TROUT		1.70	12.0	7.5	40
91	99.50% TECHNICAL MATERIAL	LAKE TROUT		1.70	12.0	6.0	40
92	99.50% TECHNICAL MATERIAL	LAKE TROUT		1.70	12.0	9.0	40
93	99.50% TECHNICAL MATERIAL	LAKE TROUT		0.50	12.0	7.4	162
94	99.50% TECHNICAL MATERIAL	LAKE TROUT		2.60	12.0	7.4	162
95	99.50% TECHNICAL MATERIAL	GOLDFISH		0.90	18.0	7.1	40
96	99.50% TECHNICAL MATERIAL	GOLDFISH		0.90	18.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
73	STATIC	LC	2300.	1630.000	3240.000	.	.	.	1150.	910.000	1450.000
74	STATIC	LC	2000.	1410.000	2840.000	.	.	.	1100.	884.000	1370.000
75	STATIC	LC	2200.	1760.000	2750.000	.	.	.	1350.	996.000	1830.000
76	STATIC	LC	500.	318.000	787.000	.	.	.	250.	118.000	767.000
77	STATIC	LC	1300.	899.000	1880.000	.	.	.	900.	721.000	1120.000
78	STATIC	LC	1150.	769.000	1720.000	.	.	.	1000.	822.000	1220.000
79	STATIC	LC	6840.	5010.000	9350.000	.	.	.	6300.	5520.000	7190.000
80	FLOW THROUGH	LC	2140.	1769.000	2588.000	.	.	.	2000.	1656.000	2414.000
81	STATIC	LC	5000.	3538.000	7066.000	.	.	.	2100.	1684.000	2618.000
82	STATIC	LC	10000.	.	.	.	.	.	3000.	2009.000	4480.000
83	STATIC	LC	770.	648.000	915.000	.	.	.	680.	585.000	790.000
84	STATIC	LC	4600.	3542.000	5974.000	.	.	.	4560.	3497.000	5947.000
85	STATIC	LC	3900.	.	.	.	.	.	2130.	1719.000	2640.000
86	STATIC	LC	2600.	.	.	.	.	.	1130.	792.000	1612.000
87	STATIC	LC	2200.	1548.000	3127.000	.	.	.	1200.	868.000	1658.000
88	STATIC	LC	3000.	2087.000	4312.000	.	.	.	1290.	887.000	1876.000
89	STATIC	LC	4500.	3948.000	5066.000	.	.	.	4500.	3948.000	5066.000
90	STATIC	LC	1300.	970.000	1830.000	.	.	.	690.	520.000	910.000
91	STATIC	LC	760.	630.000	920.000	.	.	.	740.	600.000	910.000
92	STATIC	LC	2000.	1600.000	2510.000	.	.	.	920.	700.000	1200.000
93	STATIC	LC	1420.	994.000	2040.000	.	.	.	872.	466.000	1630.000
94	FLOW THROUGH	LC	3000.	2434.000	3698.000	.	.	.	2300.	1950.000	2710.000
95	STATIC	LC	15600.	11700.000	20700.000	.	.	.	13200.	8310.000	20800.000
96	STATIC	LC	15700.	11700.000	21000.000	.	.	.	12800.	8100.000	20300.000

ED_005172C_00000035-00143

CARBARYL

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
97	99.50% TECHNICAL MATERIAL	CARP		0.60	18.0	7.1	40
98	99.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.50	12.0	7.5	42
99	99.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.80	18.0	7.1	40
100	99.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.80	18.0	7.4	272
101	99.50% TECHNICAL MATERIAL	BLACK BULLHEAD		1.20	18.0	7.1	40
102	99.50% TECHNICAL MATERIAL	CHANNEL CATFISH		1.50	18.0	7.1	40
103	99.50% TECHNICAL MATERIAL	CHANNEL CATFISH		1.50	18.0	7.4	272
104	99.50% TECHNICAL MATERIAL	CHANNEL CATFISH	FINGERLING	.	12.0	7.5	314
105	99.50% TECHNICAL MATERIAL	GREEN SUNFISH		1.10	18.0	7.1	40
106	99.50% TECHNICAL MATERIAL	GREEN SUNFISH		1.10	18.0	7.4	272
107	99.50% TECHNICAL MATERIAL	BLUEGILL		1.20	18.0	7.1	40
108	99.50% TECHNICAL MATERIAL	BLUEGILL		1.20	18.0	7.4	272
109	99.50% TECHNICAL MATERIAL	BLUEGILL		0.60	12.0	7.5	314
110	49.00% OIL DISPERSION	BLUEGILL		0.70	17.0	7.4	40
111	99.50% TECHNICAL MATERIAL	BLUEGILL		0.40	12.0	7.4	44
112	99.50% TECHNICAL MATERIAL	BLUEGILL		0.40	22.0	7.4	44
113	99.50% TECHNICAL MATERIAL	BLUEGILL		0.80	12.0	7.4	40
114	99.50% TECHNICAL MATERIAL	BLUEGILL		0.80	17.0	7.4	40
115	99.50% TECHNICAL MATERIAL	BLUEGILL		0.80	22.0	7.4	40
116	99.50% TECHNICAL MATERIAL	BLUEGILL		0.40	17.0	7.4	320
117	99.50% TECHNICAL MATERIAL	BLUEGILL		0.70	17.0	6.5	40
118	99.50% TECHNICAL MATERIAL	BLUEGILL		0.70	17.0	7.5	40
119	99.50% TECHNICAL MATERIAL	BLUEGILL		0.70	17.0	8.5	40
120	99.50% TECHNICAL MATERIAL	BLUEGILL		0.70	17.0	9.5	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
97	STATIC	LC	6140.	5370.000	7150.000	.	.	.	5280.	4620.000	6050.000
98	STATIC	LC	33200.	26800.000	41200.000	.	.	.	14000.	11400.000	17200.000
99	STATIC	LC	18000.	11700.000	30400.000	.	.	.	14600.	11700.000	19800.000
100	STATIC	LC	17300.	14500.000	20700.000	.	.	.	7700.	4800.000	12200.000
101	STATIC	LC	20000.	18000.000	24000.000	.	.	.	20000.	18000.000	24000.000
102	STATIC	LC	21300.	18300.000	24700.000	.	.	.	15800.	13900.000	18000.000
103	STATIC	LC	15400.	11400.000	20900.000	.	.	.	7790.	4750.000	12300.000
104	FLOW THROUGH	LC	>20000	.	.	.	.	.	17300.	16360.000	18290.000
105	STATIC	LC	11200.	8140.000	15500.000	.	.	.	11200.	8140.000	15500.000
106	STATIC	LC	11000.	7990.000	15200.000	.	.	.	9460.	7000.000	12800.000
107	STATIC	LC	10000.	8330.000	12500.000	.	.	.	6760.	5220.000	8760.000
108	STATIC	LC	9880.	8050.000	12100.000	.	.	.	5230.	4010.000	6820.000
109	FLOW THROUGH	LC	5340.	4447.000	6412.000	.	.	.	5047.	4357.000	5847.000
110	STATIC	LC	42000.	34971.000	50442.000	.	.	.	39000.	29732.000	51157.000
111	STATIC	LC	23000.	14940.000	35410.000	.	.	.	7400.	5420.000	10100.000
112	STATIC	LC	7400.	5500.000	10000.000	.	.	.	5200.	3660.000	7390.000
113	STATIC	LC		.	.	.	.	.	16000.	11410.000	22430.000
114	STATIC	LC		.	.	.	.	.	7000.	5020.000	9760.000
115	STATIC	LC		.	.	.	.	.	8200.	6170.000	10890.000
116	STATIC	LC	14500.	11100.000	19010.000	.	.	.	6200.	4310.000	8920.000
117	STATIC	LC		.	.	.	.	.	5400.	4260.000	6830.000
118	STATIC	LC		.	.	.	.	.	5200.	4060.000	6640.000
119	STATIC	LC		.	.	.	.	.	1800.	1390.000	2320.000
120	STATIC	LC		.	.	.	.	.	2600.	2070.000	3250.000

CARBARYL
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
121	99.50% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.90	18.0	7.1	40
122	99.50% TECHNICAL MATERIAL	BLACK CRAPPIE		1.00	18.0	7.1	40
123	99.50% TECHNICAL MATERIAL	YELLOW PERCH		1.40	18.0	7.1	40
124	99.50% TECHNICAL MATERIAL	YELLOW PERCH		0.60	12.0	7.5	42
125	99.50% TECHNICAL MATERIAL	YELLOW PERCH		1.00	7.0	7.5	42
126	99.50% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	7.5	42
127	99.50% TECHNICAL MATERIAL	YELLOW PERCH		1.00	17.0	7.5	42
128	99.50% TECHNICAL MATERIAL	YELLOW PERCH		1.00	22.0	7.5	42
129	99.50% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	6.5	42
130	99.50% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	7.5	42
131	99.50% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	8.5	42
132	99.50% TECHNICAL MATERIAL	YELLOW PERCH		0.90	12.0	9.0	42
133	99.50% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	8.0	42
134	99.50% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	8.0	170
135	99.50% TECHNICAL MATERIAL	YELLOW PERCH		1.00	12.0	8.0	300
136	99.50% TECHNICAL MATERIAL	YELLOW PERCH	FINGERLING	.	12.0	7.5	314

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
121	STATIC	LC	16900.	15800.000	17300.000	.		.	6400.	4400.000	9200.000
122	STATIC	LC	3740.	2250.000	6190.000	.		.	2600.	1180.000	5700.000
123	STATIC	LC	1030.	826.000	1270.000	.		.	745.	611.000	904.000
124	STATIC	LC	8600.	7260.000	10190.000	.		.	5100.	4520.000	5760.000
125	STATIC	LC	14400.	12150.000	17070.000	.		.	13900.	11700.000	16510.000
126	STATIC	LC	6100.	4800.000	7740.000	.		.	5400.	4140.000	7050.000
127	STATIC	LC	3400.	2660.000	4350.000	.		.	3400.	2690.000	4290.000
128	STATIC	LC	2250.	1580.000	3210.000	.		.	1200.	1000.000	1440.000
129	STATIC	LC	5100.	4230.000	6150.000	.		.	4000.	3030.000	5290.000
130	STATIC	LC	4700.	3650.000	6060.000	.		.	4200.	3100.000	5690.000
131	STATIC	LC	>10000	.	.	.		.	480.	304.000	757.000
132	STATIC	LC	4950.	3850.000	6360.000	.		.	350.	280.000	430.000
133	STATIC	LC	5400.	4020.000	7250.000	.		.	3800.	2730.000	5500.000
134	STATIC	LC	7000.	5540.000	8840.000	.		.	5000.	4030.000	6200.000
135	STATIC	LC	7800.	6180.000	9850.000	.		.	3750.	2850.000	4930.000
136	FLOW THROUGH	LC	2320.	1839.000	2926.000	.		.	1420.	1040.000	1938.000

ED_005172C_00000035-00145

CAS # 1563-66-2 PRINCIPAL USE: INSECTICIDE

CARBOFURAN
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT		DEG. CENT.	PH	HARDNESS MG/L
1	99.00% TECHNICAL MATERIAL	COHO SALMON		0.60		12.0	7.5	44
2	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.50		12.0	7.4	44
3	99.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.30	STEELHEAD STRAIN	12.0	7.5	44
4	99.00% TECHNICAL MATERIAL	BROWN TROUT		0.50		12.0	7.5	44
5	99.00% TECHNICAL MATERIAL	BROWN TROUT		0.60		12.0	7.5	314
6	99.00% TECHNICAL MATERIAL	LAKE TROUT		0.50		12.0	7.5	314
7	99.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.50		12.0	7.5	44
8	99.00% TECHNICAL MATERIAL	FATHEAD MINNOW		1.30		17.0	7.1	44
9	99.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.50		17.0	7.5	314
10	99.00% TECHNICAL MATERIAL	CHANNEL CATFISH		1.00		20.0	7.4	44
11	50.00% WETTABLE POWDER	BLUEGILL		0.80		18.0	7.1	44
12	100.0% TECHNICAL MATERIAL	BLUEGILL		1.10		17.0	7.1	40
13	99.00% TECHNICAL MATERIAL	YELLOW PERCH		0.60		12.0	7.5	44
14	99.00% TECHNICAL MATERIAL	YELLOW PERCH		0.60		12.0	9.5	44
15	99.00% TECHNICAL MATERIAL	YELLOW PERCH		1.30		12.0	7.5	42

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	530.	432.000	650.000	.	.		530.	432.000	650.000
2	STATIC	LC	680.	536.000	863.000	.	.		380.	272.000	531.000
3	STATIC	LC	1020.	635.000	1640.000	.	.		600.	436.000	826.000
4	STATIC	LC	842.	705.000	1010.000	.	.		560.	475.000	660.000
5	FLOW THROUGH	LC	355.	241.000	521.000	.	.		280.	204.000	382.000
6	FLOW THROUGH	LC	164.	119.000	226.000	.	.		164.	119.000	226.000
7	STATIC	LC	2240.	1590.000	3150.000	.	.		1990.	1385.000	2859.000
8	STATIC	LC	883.	494.000	1580.000	.	.		872.	479.000	1590.000
9	FLOW THROUGH	LC	1320.	990.000	1758.000	.	.		1180.	813.000	1711.000
10	STATIC	LC	372.	215.000	642.000	.	.		248.	94.000	649.000
11	STATIC	LC	370.	289.000	473.000	.	.		240.	186.000	310.000
12	STATIC	LC	101.	87.000	117.000	.	.		88.	79.000	104.000
13	STATIC	LC	275.	226.000	335.000	.	.		240.	208.000	277.000
14	STATIC	LC		.	.	.	.		120.	82.000	176.000
15	STATIC	LC	540.	426.000	685.000	.	.		400.	289.000	553.000

ED_005172C_00000035-00146

CARBOPHENOTHION
MEASURED IN UG /L

CAS # 786-19-6 PRINCIPAL USE: INSECTICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	95.30% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	16.0	7.1	40
2	95.30% TECHNICAL MATERIAL	GAMMARUS LACUSTRIS	MATURE	.	21.0	7.1	40
3	95.30% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	MATURE	.	21.0	7.4	272
4	95.30% TECHNICAL MATERIAL	CHANNEL CATFISH		1.10	18.0	7.1	40
5	95.30% TECHNICAL MATERIAL	GREEN SUNFISH		0.80	18.0	7.1	40
6	95.30% TECHNICAL MATERIAL	BLUEGILL		1.10	18.0	7.1	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC		10.000	100.000	.	.			10.000	100.000
2	STATIC	LC	45.	30.000	62.000	.	.		5.2	4.100	6.500
3	STATIC	LC	18.	12.000	27.000	.	.		1.2	0.800	1.400
4	STATIC	LC	11800.	10900.000	12800.000	.	.		6000.	5740.000	6270.000
5	STATIC	LC	3950.	.	.	.	.		280.	186.000	422.000
6	STATIC	LC	27.	21.000	34.000	.	.		13.	10.000	16.000

145

ED_005172C_00000035-00147

CAS # PRINCIPAL USE: WOOD PRESERVATIVE CCA TYPE III

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	50.00% LIQUID	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.4	41
2	50.00% LIQUID	GAMMARUS PSEUDOLIMNAEUS	IMMATURE	.	17.0	7.4	41
3	50.00% LIQUID	RAINBOW TROUT		0.90	12.0	7.4	42
4	50.00% LIQUID	CHANNEL CATFISH		0.40	22.0	7.4	40
5	50.00% LIQUID	BLUEGILL		0.34	22.0	7.5	35

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	0.04	0.020	0.050			
2	STATIC	LC		.	.		.	.	0.05	0.040	0.070
3	STATIC	LC		.	.		.	.	<0.32	.	.
4	STATIC	LC		.	.		.	.	1.8	1.340	2.420
5	STATIC	LC	23.0	17.200	30.700		.	.	12.5	9.300	16.700

ED_005172C_00000035-00148

CHLORAMBEN
MEASURED IN MG /L

CAS # 133-90-4 PRINCIPAL USE: HERBICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	10.00% WETTABLE POWDER	RAINBOW TROUT		0.30	12.0	7.6	40
2	10.00% WETTABLE POWDER	BLUEGILL		0.34	22.0	7.5	38

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC		:	:		:	:	>10.	:	:
2	STATIC	LC		:	:		:	:	>10.	:	:

ED_005172C_00000035-00149

CHLORBROMURON

CAS # 13360-45-7 PRINCIPAL USE: HERBICIDE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	90.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.70	12.0	7.1	44
2	90.00% TECHNICAL MATERIAL	CHANNEL CATFISH		0.70	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	6.6	3.200	13.400	.	.		1.4	0.700	3.100
2	STATIC	LC	10.2	8.400	12.400	.	.		10.2	8.400	12.400

CHLORDANE

MEASURED IN UG /L

ED_005172C_00000035-00150

CAS # 57-74-9 PRINCIPAL USE: INSECTICIDE

CHLORDANE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT		DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.		15.0	7.1	44
2	100.0% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.		15.0	7.1	44
3	100.0% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.		21.0	7.1	44
4	100.0% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.		21.0	7.1	44
5	100.0% TECHNICAL MATERIAL	ORCONECTES NAIS	EARLY INSTAR	.		22.0	7.4	272
6	100.0% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.		15.0	7.1	44
7	100.0% TECHNICAL MATERIAL	COHO SALMON		0.6		12.0	7.5	44
8	100.0% TECHNICAL MATERIAL	COHO SALMON		0.9		12.0	7.1	44
9	100.0% TECHNICAL MATERIAL	COHO SALMON		4.50		12.0	7.1	44
10	100.0% TECHNICAL MATERIAL	COHO SALMON		11.10		12.0	7.1	44
11	100.0% TECHNICAL MATERIAL	COHO SALMON		19.40		12.0	7.1	44
12	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00		13.0	7.1	44
13	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00		13.0	7.4	272
14	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		0.80		7.0	7.1	44
15	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.00		12.0	7.1	44
16	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		0.80		18.0	7.1	44
17	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.20		12.0	7.1	44
18	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		4.70		12.0	7.1	44
19	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		10.80		12.0	7.1	44
20	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		23.50		12.0	7.1	44
21	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		46.00		12.0	7.1	44
22	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	FPRL 75 PERCENT PROTEIN DIET	12.0	7.1	44
23	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	FPRL 50 PERCENT PROTEIN DIET	12.0	7.1	44
24	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	FPRL 25 PERCENT PROTEIN DIET	12.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	29.0	23.000	36.000		.	.
2	STATIC	EC		.	.	20.0	12.000	32.000		.	.
3	STATIC	EC		.	.	24.0	20.000	28.000		.	.
4	STATIC	LC	100.	60.000	190.000		.	.	40.0	21.000	60.000
5	FLOW THROUGH	EC	>50.0	.	.		.	.	>50.0	.	.
6	STATIC	LC	170.	85.000	340.000		.	.	15.0	9.000	24.000
7	STATIC	LC	>20.0	.	.		.	.	14.0	11.400	17.200
8	STATIC	LC	19.7	9.800	39.500		.	.	13.9	9.490	20.500
9	STATIC	LC	>87.0	.	.		.	.	16.3	4.820	54.800
10	STATIC	LC	151.	95.100	239.000		.	.	73.5	55.000	98.300
11	STATIC	LC	155.	43.500	555.000		.	.	80.0	57.600	111.000
12	STATIC	LC	43.0	37.000	49.000		.	.	27.0	24.000	31.000
13	STATIC	LC	48.0	43.000	53.000		.	.	23.0	20.000	26.000
14	STATIC	LC	91.	81.000	102.000		.	.	45.	41.000	49.000
15	STATIC	LC	60.	54.000	67.000		.	.	42.	37.000	48.000
16	STATIC	LC	34.	27.000	41.000		.	.	29.	24.000	35.000
17	STATIC	LC	20.	15.000	26.000		.	.	10.	6.500	16.000
18	STATIC	LC	74.	62.000	89.000		.	.	45.	36.000	56.000
19	STATIC	LC	110.	91.000	133.000		.	.	59.	49.000	71.000
20	STATIC	LC	145.	112.000	188.000		.	.	72.	55.000	94.000
21	STATIC	LC	200.	130.000	309.000		.	.	135.	107.000	170.000
22	STATIC	LC	58.0	37.700	87.200		.	.	<40.0	.	.
23	STATIC	LC	73.0	60.900	87.500		.	.	47.0	37.700	58.500
24	STATIC	LC	66.0	48.700	89.400		.	.	28.5	19.800	41.000

ED_005172C_00000035-00151

CHLORDANE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT			DEG. CENT.	PH	HARDNESS MG/L
25	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	EWOS		DIET 12.0	7.1	44
26	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	SILVER CUP		DIET 12.0	7.1	44
27	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	GLENCO		DIET 12.0	7.1	44
28	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	OREGON MOIST		DIET 12.0	7.1	44
29	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	MISSOURI	STRAIN	12.0	7.1	44
30	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		2.00	NEW HAMPSHIRE	STRAIN	12.0	7.1	44
31	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	SOAP LAKE	STRAIN	12.0	7.1	44
32	72.00% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		0.80			12.0	7.1	44
33	40.00% WETTABLE POWDER	RAINBOW TROUT		0.80			12.0	7.1	44
34	10.00% GRANULAR FORMULATION	RAINBOW TROUT		1.20			12.0	7.1	44
35	25.00% GRANULAR FORMULATION	RAINBOW TROUT		0.80			12.0	7.1	44
36	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		0.25	STEELHEAD	STRAIN	12.0	7.5	42
37	100.0% TECHNICAL MATERIAL	BROWN TROUT		0.60			12.0	7.5	44
38	100.0% TECHNICAL MATERIAL	BROWN TROUT		0.60			12.0	7.5	314
39	100.0% TECHNICAL MATERIAL	FATHEAD MINNOW		0.70			17.0	7.1	44
40	100.0% TECHNICAL MATERIAL	FATHEAD MINNOW		0.40			17.0	7.5	314
41	100.0% TECHNICAL MATERIAL	WHITE SUCKER		0.40			12.0	7.5	314
42	100.0% TECHNICAL MATERIAL	WHITE SUCKER		0.40			12.0	7.5	320
43	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		1.90			17.0	7.1	44
44	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		4.30			17.0	7.1	44
45	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		9.70			17.0	7.1	44
46	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		18.50			17.0	7.1	44
47	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		0.90			18.0	7.4	44
48	100.0% TECHNICAL MATERIAL	BLUEGILL		1.40			17.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
25	STATIC	LC	61.0	48.400	76.800	.	.	.	31.0	22.500	42.600
26	STATIC	LC	40.0	28.200	56.700	.	.	.	20.0	14,200	28.000
27	STATIC	LC	40.	26.000	61.400	.	.	.	9.10	4.800	17.200
28	STATIC	LC	27.0	20.700	35.100	.	.	.	8.2	6.100	11.000
29	STATIC	LC	43.6	28.200	67.400	.	.	.	27.0	19.000	38.600
30	STATIC	LC	77.7	38.200	158.000	.	.	.	19.5	9.020	42.000
31	STATIC	LC	62.4	42.500	91.600	.	.	.	12.6	7.340	21.600
32	STATIC	LC	28.1	21.900	36.100	.	.	.	4.43	2.630	7.480
33	STATIC	LC	72.1	63.000	82.500	.	.	.	19.0	11.200	32.000
34	STATIC	LC	28.6	21.500	38.000	.	.	.	10.2	4.970	30.500
35	STATIC	LC	15.4	11.200	21.100	.	.	.	7.11	3.710	13.600
36	STATIC	LC	12.9	10.700	15.600	.	.	.	2.9	1.500	5.600
37	STATIC	LC	30.4	26.500	34.900	.	.	.	11.1	9.380	13.100
38	FLOW THROUGH	LC	38.0	31.300	46.100	.	.	.	11.1	8.890	13.900
39	STATIC	LC	420.	293.000	601.000	.	.	.	115.	62.000	214.000
40	STATIC	LC	320.	241.000	424.000	.	.	.	56.0	28.100	112.000
41	FLOW THROUGH	LC	>30.0	.	.	.	.	.	17.	11.000	27.000
42	FLOW THROUGH	LC	>30.	.	.	.	.	.	17.1	10.700	27.100
43	STATIC	LC	28.	12.000	63.000	.	.	.	6.7	3.100	15.000
44	STATIC	LC	194.	136.000	277.000	.	.	.	70.	28.000	176.000
45	STATIC	LC	260.	206.000	328.000	.	.	.	230.	186.000	285.000
46	STATIC	LC	370.	311.000	440.000	.	.	.	225.	192.000	263.000
47	STATIC	LC	1.6	.	.	.	.	.	0.8	.	.
48	STATIC	LC	145.	119.000	176.000	.	.	.	57.	40.000	81.000

ED_005172C_00000035-00152

CHLORDANE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
49	100.0% TECHNICAL MATERIAL	BLUEGILL		4.80	17.0	7.1	44
50	100.0% TECHNICAL MATERIAL	BLUEGILL		20.00	17.0	7.1	44
51	100.0% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.80	18.0	7.4	272
52	100.0% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.90	17.0	7.1	44
53	100.0% TECHNICAL MATERIAL	LARGEMOUTH BASS		4.20	17.0	7.1	44
54	100.0% TECHNICAL MATERIAL	LARGEMOUTH BASS		6.10	17.0	7.1	44
55	100.0% TECHNICAL MATERIAL	LARGEMOUTH BASS		19.50	17.0	7.1	44
56	100.0% TECHNICAL MATERIAL	YELLOW PERCH		0.60	12.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
49	STATIC	LC	157.	124.000	226.000	.	.	.	92.	64.000	133.000
50	STATIC	LC	170.	135.000	215.000	.	.	.	128.	107.000	153.000
51	STATIC	LC	27.	23.000	32.000	.	.	.	12.	10.000	14.000
52	STATIC	LC	94.	57.000	160.000	.	.	.	32.	.	.
53	STATIC	LC	180.	140.000	220.000	.	.	.	56.	42.000	74.000
54	STATIC	LC	270.	216.000	337.000	.	.	.	140.	108.000	182.000
55	STATIC	LC	320.	271.000	378.000	.	.	.	130.	98.000	172.000
56	FLOW THROUGH	LC	18.5	14.700	23.200	.	.	.	9.60	8.050	11.400

151

ED_005172C_00000035-00153

CHLORDANE CIS
MEASURED IN UG /L

CAS # 5103-71-9 PRINCIPAL USE: INSECTICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	BLUEGILL		1.00	17.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	28.3	19.400	41.400	.	.		7.09	4.270	11.800

ED_005172C_00000035-00154

CHLORDANE-HCS-3260
MEASURED IN UG /L

CAS # PRINCIPAL USE: INSECTICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	RAINBOW TROUT		1.50	12.0	7.1	44
2	45.00% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		1.50	12.0	7.1	44
3	33.30% GRANULAR FORMULATION	RAINBOW TROUT		1.50	12.0	7.1	44
4	100.0% TECHNICAL MATERIAL	FATHEAD MINNOW		0.70	17.0	7.1	44
5	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		1.80	17.0	7.1	44
6	100.0% TECHNICAL MATERIAL	BLUEGILL		1.30	17.0	7.1	44
7	45.00% EMULSIFIABLE CONCENTRATE	BLUEGILL		1.50	17.0	7.1	44
8	33.30% GRANULAR FORMULATION	BLUEGILL		1.50	17.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	36.2	23.400	56.000	.	.		24.9	16.100	38.500
2	STATIC	LC	55.6	10.400	299.000	.	.		24.9	16.100	38.500
3	STATIC	LC	32.6	22.900	46.500	.	.		.	.	.
4	STATIC	LC	80.7	.	.	.	.		24.8	.	.
5	STATIC	LC	47.3	.	.	.	.		45.8	.	.
6	STATIC	LC	49.8	.	.	.	.		29.3	.	.
7	STATIC	LC	27.0	.	.	.	.		23.4	.	.
8	STATIC	LC	52.0	.	.	.	.		19.1	.	.

ED_005172C_00000035-00155

CAS # 5103-74-2 PRINCIPAL USE: INSECTICIDE

CHLORDANE TRANS
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	99.80% TECHNICAL MATERIAL	BLUEGILL		1.00	17.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	210.	137.000	321.000	.	.	.	50.5	30.800	82.700

BEST AVAILABLE COPY

CAS # 143-50-0 PRINCIPAL USE: INSECTICIDE

CHLORDECONE
MEASURED IN UG /L

ED_005172C_00000035-00156

CAS # 143-50-0 PRINCIPAL USE: INSECTICIDE

CHLORDECONE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	90.70% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.2	43
2	90.70% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	17.0	7.7	38
3	90.70% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	22.0	7.7	38
4	90.70% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	12.0	7.1	44
5	90.70% TECHNICAL MATERIAL	RAINBOW TROUT		1.10	12.0	7.4	272
6	94.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.03	22.0	7.1	44
7	94.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.30	22.0	7.4	272
8	90.70% TECHNICAL MATERIAL	CHANNEL CATFISH		1.60	18.0	7.1	44
9	90.70% TECHNICAL MATERIAL	BLUEGILL		3.50	24.0	7.1	44
10	90.70% TECHNICAL MATERIAL	REDEAR SUNFISH		1.00	7.0	7.1	44
11	90.70% TECHNICAL MATERIAL	REDEAR SUNFISH		1.00	12.0	7.1	44
12	90.70% TECHNICAL MATERIAL	REDEAR SUNFISH		1.00	18.0	7.1	44
13	90.70% TECHNICAL MATERIAL	REDEAR SUNFISH		1.00	24.0	7.1	44
14	90.70% TECHNICAL MATERIAL	REDEAR SUNFISH		1.00	29.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	260.	200.000	345.000			
2	STATIC	LC		.	.		.	.	180.	110.000	290.000
3	STATIC	EC		.	.	320.	230.000	450.000		.	.
4	STATIC	LC	64.	50.000	82.000		.	.	30.	24.000	38.000
5	STATIC	LC	100.	87.000	115.000		.	.	29.	23.000	37.000
6	STATIC	LC		.	.		.	.	420.	373.000	473.000
7	FLOW THROUGH	LC		.	.		.	.	340.	289.000	400.000
8	STATIC	LC	225.	.	.		.	.	225.	.	.
9	STATIC	LC	280.	.	.		.	.	72.	.	.
10	STATIC	LC	620.	570.000	680.000		.	.	140.	130.000	150.000
11	STATIC	LC	540.	490.000	590.000		.	.	96.	87.000	105.000
12	STATIC	LC	340.	310.000	370.000		.	.	64.	59.000	70.000
13	STATIC	LC	240.	220.000	260.000		.	.	44.	41.000	47.000
14	STATIC	LC	120.	110.000	130.000		.	.	29.	26.000	32.000

155

ED_005172C_00000035-00157

CAS # 6164-98-3 PRINCIPAL USE: INSECTICIDE

CHLORDIMEFORM
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	95.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.70	12.0	7.1	44
2	95.00% TECHNICAL MATERIAL	CHANNEL CATFISH		0.70	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	29.0	20.400	41.400	.	.		13.2	8.100	21.300
2	STATIC	LC	20.7	14.900	28.600	.	.		20.2	14.800	27.600

ED_005172C_00000035-00158

CAS # 1770-80-5 PRINCIPAL USE: PLASTICIZER

CHLORENDATE DIBUTYL
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		1.00	6.0	7.4	44
2	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		0.80	16.0	7.4	272
3	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		1.40	16.0	7.4	272
4	100.0% COMMERCIAL FORMULATION	FATHEAD MINNOW		1.00	20.0	7.4	44
5	100.0% COMMERCIAL FORMULATION	CHANNEL CATFISH		1.00	20.0	7.4	44
6	100.0% COMMERCIAL FORMULATION	BLUEGILL		1.40	20.0	7.4	44
7	100.0% COMMERCIAL FORMULATION	BLUEGILL		0.80	12.0	7.6	314

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>150.	.	.	.	.	.	170.	132.000	210.000
2	FLOW THROUGH	LC	>0.150	.	.	.	.	.	0.050	0.038	0.066
3	FLOW THROUGH	LC	>0.020	.	.	.	.	.	>0.020	.	.
4	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
5	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
6	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
7	FLOW THROUGH	LC	>10.	.	.	.	.	.	10.4	8.900	12.000

157

ED_005172C_00000035-00159

CAS # 1773-89-3 PRINCIPAL USE: PLASTICIZER

CHLORENDATE DIMETHYL
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		1.00	6.0	7.4	44
2	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		1.00	11.0	7.4	44
3	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		0.80	16.0	7.4	272
4	100.0% COMMERCIAL FORMULATION	FATHEAD MINNOW		1.00	20.0	7.4	44
5	100.0% COMMERCIAL FORMULATION	FATHEAD MINNOW		1.60	16.0	7.4	272
6	100.0% COMMERCIAL FORMULATION	CHANNEL CATFISH		1.00	20.0	7.4	44
7	100.0% COMMERCIAL FORMULATION	CHANNEL CATFISH		1.20	16.0	7.4	272
8	100.0% COMMERCIAL FORMULATION	BLUEGILL		1.40	20.0	7.4	44
9	100.0% COMMERCIAL FORMULATION	BLUEGILL		1.50	20.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	3.25	2.770	3.810	.	.	.	1.75	1.510	2.020
2	STATIC	LC	3.00	2.420	3.720	.	.	.	1.40	1.040	1.890
3	FLOW THROUGH	LC	>0.70	.	.	.	.	.	0.42	0.360	0.480
4	STATIC	LC	2.21	0.960	5.090	.	.	.	1.70	1.250	2.300
5	FLOW THROUGH	LC	>2.5	.	.	.	.	.	1.75	1.610	1.900
6	STATIC	LC		.	.	.	.	.	2.00	1.590	2.510
7	FLOW THROUGH	LC	>2.50	.	.	.	.	.	2.30	1.390	3.830
8	STATIC	LC	4.12	3.000	5.660	.	.	.	2.24	1.830	2.740
9	FLOW THROUGH	LC	>2.50	.	.	.	.	.	2.70	2.250	3.230

CHLORFENETHOL
MEASURED IN MG /L

CAS # 80-06-8 PRINCIPAL USE: ACARICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	0.50	21.0	7.1	44
2	100.0% TECHNICAL MATERIAL	FATHEAD MINNOW		1.20	18.0	7.1	44
3	100.0% TECHNICAL MATERIAL	FATHEAD MINNOW		1.20	18.0	7.4	272
4	100.0% TECHNICAL MATERIAL	CHANNEL CATFISH		1.40	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC	1.74	1.410	2.150	0.27	.	.	1.39	0.950	2.050
2	STATIC	LC	1.88	1.400	2.520		.	.	1.52	0.960	2.410
3	STATIC	LC	1.70	1.600	1.800		.	.	0.90	0.650	1.200
4	STATIC	LC									

ED_005172C_00000035-00161

CAS # 470-90-6 PRINCIPAL USE: INSECTICIDE

CHLORFENVINPHOS
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	92.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	21.0	7.4	44
2	92.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.4	44
3	92.00% TECHNICAL MATERIAL	RAINBOW TROUT		4.90	12.0	7.4	44
4	92.00% TECHNICAL MATERIAL	BLUEGILL		0.30	24.0	7.4	44
5	92.00% TECHNICAL MATERIAL	BLUEGILL		0.90	18.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	27	18.000	41.000	.	.	.	9.6	7.300	12.700
2	STATIC	LC	5.8	3.700	9.200	.	.	.	0.7	0.500	1.100
3	STATIC	LC	1650	1240.000	2190.000	.	.	.	510	420.000	620.000
4	STATIC	LC	2.8	2.200	3.600	.	.	.	.	.	.
5	STATIC	LC	50	40.000	62.000	.	.	.	23	17.000	31.000

ED_005172C_00000035-00162

CHLORINATED ALPHA OLEFIN 22/44
MEASURED IN MG /L

PRINCIPAL USE: PLASTICIZER

OBS	CAS #	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0%	COMMERCIAL FORMULATION	RAINBOW TROUT		0.70	10.0	7.4	44
2	100.0%	COMMERCIAL FORMULATION	BLUEGILL		0.50	20.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL	48 HOUR	CONFIDENCE INTERVAL	96 HOUR	CONFIDENCE INTERVAL
1	STATIC	LC		:		:	>300.	:
2	STATIC	LC		:		:	>300.	:

ED_005172C_00000035-00163

CAS # 999-81-5 PRINCIPAL USE: GROWTH REGULATOR

CHLORMEQUAT CHLORIDE
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	98.10% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	12.0	7.4	272
2	98.10% TECHNICAL MATERIAL	FATHEAD MINNOW		1.50	17.0	7.4	272
3	98.10% TECHNICAL MATERIAL	CHANNEL CATFISH		1.50	17.0	7.4	272
4	98.10% TECHNICAL MATERIAL	BLUEGILL		1.50	17.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
2	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
3	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
4	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.

162

ED_005172C_00000035-00164

CHLOROBENZILATE
MEASURED IN UG /L

CAS # 510-15-6 PRINCIPAL USE: ACARICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	95.50% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	15.0	7.1	48
2	95.50% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	0.80	15.0	7.1	44
3	95.50% TECHNICAL MATERIAL	RAINBOW TROUT			13.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	870.	690.000	1100.000		.	.
2	STATIC	EC	750.	.	.	550.	460.000	650.000	670.	.	.
3	STATIC	LC					.	.			

103

ED_005172C_00000035-00165

CAS # 2425-66-3 PRINCIPAL USE: FUNGICIDE

CHLORONITROPROPANE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	45.00% EMULSIFIABLE CONCENTRATE	GAMMARUS LACUSTRIS	MATURE	.	21.0	7.1	44
2	45.00% EMULSIFIABLE CONCENTRATE	RAINBOW TROUT		1.50	12.0	7.1	44
3	45.00% EMULSIFIABLE CONCENTRATE	BLUEGILL		1.10	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	2800.	1090.000	3836.000	.	.	.	640.	460.000	889.000
2	STATIC	LC	120.	111.000	129.000	.	.	.	110.	96.000	126.000
3	STATIC	LC	152.	140.000	165.000	.	.	.	152.	143.000	162.000

ED_005172C_00000035-00166

6-CHLORO-2-PICOLINIC ACID
MEASURED IN MG /L

CAS # 4684-94-0 PRINCIPAL USE: NITROGEN STABILIZER

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	99.00% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	21.0	7.4	272
2	99.00% TECHNICAL MATERIAL	ASELLUS BREVICAUDUS	MATURE	.	16.0	7.4	272
3	99.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	16.0	7.4	272
4	99.00% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	MATURE	.	21.0	7.4	272
5	99.00% TECHNICAL MATERIAL	RAINBOW TROUT	FINGERLING	1.40	12.0	7.1	44
6	99.00% TECHNICAL MATERIAL	CHANNEL CATFISH			18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC	7.0	.	.	1.6	.	.	1.3	.	.
2	STATIC	LC	>8.0	.	.		.	.	10.0	.	.
3	STATIC	LC	3.2	.	.		.	.	0.4	.	.
4	STATIC	LC	88.	.	.		.	.	82.	.	.
5	STATIC	LC	145.	.	.		.	.	53.	.	.
6	STATIC	LC									

165

ED_005172C_00000035-00167

CAS # PRINCIPAL USE: INDUSTRIAL CHLOROWAX LV
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		0.70	10.0	7.4	44
2	100.0% COMMERCIAL FORMULATION	BLUEGILL		0.50	20.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>300.	:	:		:	:	>300.	:	:
2	STATIC	LC	>300.	:	:		:	:	>300.	:	:

ED_005172C_00000035-00168

CHLOROWAX ND
MEASURED IN MG /L

CAS # 51990-12-6 PRINCIPAL USE: INDUSTRIAL

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		0.70	10.0	7.4	44
2	100.0% COMMERCIAL FORMULATION	BLUEGILL		0.70	20.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>300.	:	:		:	:	>300.	:	:
2	STATIC	LC	>300.	:	:		:	:	>300.	:	:

ED_005172C_00000035-00169

CAS # 56509-64-9 PRINCIPAL USE: INDUSTRIAL

CHLOROWAX 50
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		0.70	10.0	7.4	44
2	100.0% COMMERCIAL FORMULATION	BLUEGILL		0.80	20.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>300.	:	:		:	:	>300.	:	:
2	STATIC	LC	>300.	:	:		:	:	>300.	:	:

ED_005172C_00000035-00170

CHLOROWAX 70
MEASURED IN MG /L

OBS	CAS #	CHEMICAL DESCRIPTION	PRINCIPAL USE: INDUSTRIAL	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0%	COMMERCIAL FORMULATION		RAINBOW TROUT		0.70	10.0	7.4	44
2	100.0%	COMMERCIAL FORMULATION		BLUEGILL		0.50	20.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>300.	:	:		:	:	>300.	:	:
2	STATIC	LC	>300.	:	:		:	:	>300.	:	:

ED_005172C_00000035-00171

CAS # 63449-39-8 PRINCIPAL USE: INDUSTRIAL

CHLOROWAX 500C
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		1.10	5.0	7.4	44
2	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		0.50	10.0	7.4	44
3	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		1.10	15.0	7.4	44
4	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT		1.00	10.0	7.4	272
5	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT	SWIMUP FRY	.	10.0	7.4	272
6	100.0% COMMERCIAL FORMULATION	RAINBOW TROUT	YOLK-SAC FRY	.	10.0	7.4	272
7	100.0% COMMERCIAL FORMULATION	FATHEAD MINNOW		0.80	20.0	7.4	44
8	100.0% COMMERCIAL FORMULATION	CHANNEL CATFISH		1.10	20.0	7.4	44
9	100.0% COMMERCIAL FORMULATION	CHANNEL CATFISH		2.10	20.0	7.4	272
10	100.0% COMMERCIAL FORMULATION	BLUEGILL		1.20	15.0	7.4	44
11	100.0% COMMERCIAL FORMULATION	BLUEGILL		0.70	20.0	7.4	44
12	100.0% COMMERCIAL FORMULATION	BLUEGILL		1.20	25.0	7.4	44
13	100.0% COMMERCIAL FORMULATION	BLUEGILL		1.40	20.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>300.	.	.	.	.	.	>300.	.	.
2	STATIC	LC	>300.	.	.	.	.	.	>300.	.	.
3	STATIC	LC	>300.	.	.	.	.	.	>300.	.	.
4	FLOW THROUGH	LC	>0.10	.	.	.	.	.	>0.10	.	.
5	FLOW THROUGH	LC	>0.10	.	.	.	.	.	>0.10	.	.
6	FLOW THROUGH	LC	>0.10	.	.	.	.	.	.	0.060	0.080
7	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
8	STATIC	LC	>300.	.	.	.	.	.	>300.	.	.
9	FLOW THROUGH	LC	>0.10	.	.	.	.	.	>0.10	.	.
10	STATIC	LC	>300.	.	.	.	.	.	>300.	.	.
11	STATIC	LC	>300.	.	.	.	.	.	>300.	.	.
12	STATIC	LC	>300.	.	.	.	.	.	>300.	.	.
13	FLOW THROUGH	LC	>0.10	.	.	.	.	.	>0.10	.	.

ED_005172C_00000035-00172

CHLOROXURON

CAS # 1982-47-4 PRINCIPAL USE: HERBICIDE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	95.00% TECHNICAL MATERIAL	RAINBOW TROUT		0.70	12.0	7.4	44
2	95.00% TECHNICAL MATERIAL	CHANNEL CATFISH		1.30	18.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>3720.	.	.	.	.	.	430.	362.000	510.000
2	STATIC	LC	800.	698.000	917.000	.	.	.	450.	163.000	1239.000

ED_005172C_00000035-00173

CAS # 2921-86-2 PRINCIPAL USE: INSECTICIDE

CHLORPYRIFOS
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	97.00% TECHNICAL MATERIAL	GAMMARUS LACUSTRIS	MATURE	.	21.0	7.1	44
2	97.00% TECHNICAL MATERIAL	CLAASSENIA SABULOSA	2ND YEAR CLASS	.	15.0	7.1	44
3	97.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	15.0	7.1	44
4	97.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.40	10.0	7.5	44
5	97.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.90	10.0	9.0	44
6	97.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.90	10.0	7.4	162
7	97.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		2.30	10.0	7.4	162
8	97.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	2.0	7.1	44
9	97.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	7.0	7.1	44
10	97.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	13.0	7.1	44
11	97.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	18.0	7.1	44
12	97.00% TECHNICAL MATERIAL	LAKE TROUT		2.30	12.0	6.0	44
13	97.00% TECHNICAL MATERIAL	LAKE TROUT		2.30	12.0	7.5	44
14	97.00% TECHNICAL MATERIAL	LAKE TROUT		2.30	12.0	9.0	44
15	97.00% TECHNICAL MATERIAL	LAKE TROUT		0.30	12.0	7.4	162
16	97.00% TECHNICAL MATERIAL	LAKE TROUT		2.90	12.0	7.4	162
17	97.00% TECHNICAL MATERIAL	LAKE TROUT		2.30	12.0	7.4	162
18	97.00% TECHNICAL MATERIAL	CHANNEL CATFISH		0.80	18.0	7.1	44
19	97.00% TECHNICAL MATERIAL	BLUEGILL		0.60	18.0	7.1	44
20	97.00% TECHNICAL MATERIAL	BLUEGILL		0.50	13.0	7.4	272
21	97.00% TECHNICAL MATERIAL	BLUEGILL		0.50	18.0	7.4	272
22	97.00% TECHNICAL MATERIAL	BLUEGILL		0.50	24.0	7.4	272
23	97.00% TECHNICAL MATERIAL	BLUEGILL		0.50	29.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	0.76	0.470	1.200	.	.	.	0.11	0.070	0.170
2	STATIC	LC	8.2	4.900	14.000	.	.	.	0.57	0.400	0.800
3	STATIC	LC	50.	38.000	65.000	.	.	.	10.	7.000	13.000
4	STATIC	LC	>32.	.	.	.	.	.	18.4	15.600	21.700
5	STATIC	LC	>32.	.	.	.	.	.	5.4	4.100	7.100
6	STATIC	LC	120.	89.500	160.000	.	.	.	26.0	16.800	40.300
7	STATIC	LC	95.9	71.000	130.000	.	.	.	13.4	9.900	18.000
8	STATIC	LC	550.	480.000	630.000	.	.	.	51.	43.000	60.000
9	STATIC	LC	110.	90.000	130.000	.	.	.	15.	13.000	17.000
10	STATIC	LC	53.	45.000	62.000	.	.	.	7.1	6.000	8.400
11	STATIC	LC	15.	12.000	19.000	.	.	.	<1.	.	.
12	STATIC	LC	230.	.	.	.	.	.	140.	90.000	217.000
13	STATIC	LC	230.	.	.	.	.	.	98.	54.000	180.000
14	STATIC	LC		.	.	.	.	.	205.	130.000	320.000
15	STATIC	LC	419.	278.000	632.000	.	.	.	227.	171.000	303.000
16	STATIC	LC	195.	91.000	413.000	.	.	.	73.	45.000	121.000
17	FLOW THROUGH	LC		.	.	.	.	.	244.	205.000	290.000
18	STATIC	LC	410.	286.000	588.000	.	.	.	280.	206.000	381.000
19	STATIC	LC	>10.	.	.	.	.	.	2.4	1.100	5.100
20	STATIC	LC		.	.	.	.	.	4.2	.	.
21	STATIC	LC		.	.	.	.	.	1.8	.	.
22	STATIC	LC		.	.	.	.	.	2.5	.	.
23	STATIC	LC		.	.	.	.	.	1.7	.	.

172

ED_005172C_00000035-00174

CAS # 5598-13-0 PRINCIPAL USE: INSECTICIDE

CHLORPYRIFOS-METHYL
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	13.0	7.1	44
2	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	13.0	7.4	272
3	96.00% TECHNICAL MATERIAL	BROOK TROUT		1.15	12.0	7.5	40
4	61.00% EMULSIFIABLE CONCENTRATE	BROOK TROUT		1.15	12.0	7.5	40
5	98.00% TECHNICAL MATERIAL	FATHEAD MINNOW		0.90	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	567.	464.000	684.000	.	.	.	301.	233.000	389.000
2	STATIC	LC	151.	113.000	201.000	.	.	.	120.	91.000	159.000
3	STATIC	LC	400.	280.000	570.000	.	.	.	200.	140.000	280.000
4	STATIC	LC	.	.	.	.	.	.	100.	65.000	150.000
5	STATIC	LC	712.	559.000	907.000	.	.	.	678.	542.000	846.000

173

ED_005172C_00000035-00175

CAS # PRINCIPAL USE: ROAD OIL

CLARIFITE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% LIQUID	RAINBOW TROUT		1.20	12.0	7.5	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>18.	.	.		.	.	13.	10.000	17.000

174

ED_005172C_00000035-00176

CAS # 1420-04-8 PRINCIPAL USE: MOLLUSCICIDE

CLONITRALIDE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	93.00% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	21.0	7.4	272
2	99.00% TECHNICAL MATERIAL	CYPRIDOPSIS VIDUA	MATURE	.	21.0	7.4	44
3	99.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	MATURE	.	15.0	7.4	44
4	93.00% TECHNICAL MATERIAL	GAMMARUS PSEUDOLIMNAEUS	MATURE	.	21.0	7.4	272
5	93.00% TECHNICAL MATERIAL	ORCONECTES NAIS	EARLY INSTAR	.	21.0	7.4	272
6	93.00% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	MATURE	.	21.0	7.4	272
7	93.00% TECHNICAL MATERIAL	PTERONARCYS CALIFORNICA	2ND YEAR CLASS	.	21.0	7.1	44
8	93.00% TECHNICAL MATERIAL	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	21.0	7.4	272
9	93.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.40	12.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	190.	140.000	270.000		.	.
2	STATIC	LC		.	.	310.	229.000	419.000		.	.
3	STATIC	LC	2000.	791.000	5058.000		.	.	360.	221.000	586.000
4	STATIC	LC	5600.	4700.000	6700.000		.	.	2400.	1800.000	3100.000
5	STATIC	LC	32000.	23000.000	45000.000		.	.	25000.	19000.000	33000.000
6	STATIC	LC	19000.	12000.000	29000.000	11000.	7800.000	16000.000	10000.	7000.000	15000.000
7	STATIC	LC	2300.	1300.000	3900.000	900.	510.000	1600.000	200.	130.000	300.000
8	STATIC	EC		.	.	1600.	1100.000	2200.000		.	.
9	STATIC	LC	680.	543.000	852.000		.	.	340.	289.000	399.000

ED_005172C_00000035-00177

CAS # 61912-87-6 PRINCIPAL USE: FUNGICIDE

COPPER-COUNT-N
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	8.00 % SPRAY CONCENTRATE	RAINBOW TROUT		1.00	12.0	7.1	44
2	8.00 % SPRAY CONCENTRATE	FATHEAD MINNOW		1.00	17.0	7.1	44
3	8.00 % SPRAY CONCENTRATE	BLUEGILL		1.20	17.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	31.9	22.700	44.600	.	.	.	20.4	14.200	28.200
2	STATIC	LC	55.3	43.300	70.700	.	.	.	35.9	26.300	48.900
3	STATIC	LC	13900.	9450.000	20400.000	.	.	.	3280.	1860.000	5750.000

178

COPPER-COUNT-NS

ED_005172C_00000035-00178

COPPER-COUNT-NS
MEASURED IN UG /L

CAS #

PRINCIPAL USE: FUNGICIDE

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	13.00% SPRAY CONCENTRATE	RAINBOW TROUT		1.00	12.0	7.1	44
2	13.00% SPRAY CONCENTRATE	FATHEAD MINNOW		1.10	17.0	7.1	44
3	13.00% SPRAY CONCENTRATE	BLUEGILL		1.20	17.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	139.	103.000	188.000	.	.	.	121.	90.200	164.000
2	STATIC	LC	484.	271.000	863.000	.	.	.	154.	92.500	255.000
3	STATIC	LC	13900.	9450.000	20400.000	.	.	.	13700.	9360.000	20400.000

177

ED_005172C_00000035-00179

CAS # 1332-40-7 PRINCIPAL USE: FUNGICIDE

COPPER OXYCHLORIDE
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	99.00% TECHNICAL MATERIAL	BLUEGILL		1.00	24.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	1.10	.	.		.	.	0.98	.	.

178

ED_005172C_00000035-00180

CAS # 7758-98-7 PRINCIPAL USE: ALGICIDE

COPPER SULFATE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	98.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.60	13.0	7.1	44
2	98.00% TECHNICAL MATERIAL	GOLDFISH		0.90	18.0	7.4	272
3	98.00% TECHNICAL MATERIAL	FATHEAD MINNOW		1.20	18.0	7.4	272
4	98.00% TECHNICAL MATERIAL	GREEN SUNFISH		1.10	18.0	7.1	44
5	98.00% TECHNICAL MATERIAL	GREEN SUNFISH		1.10	18.0	7.4	272
6	98.00% TECHNICAL MATERIAL	BLUEGILL		1.50	18.0	7.1	44
7	98.00% TECHNICAL MATERIAL	BLUEGILL		1.50	18.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	150.	.	.	.	.	.	135.	.	.
2	STATIC	LC	4490.	3260.000	6200.000	.	.	.	1380.	989.000	1930.000
3	STATIC	LC	1680.	1260.000	2250.000	.	.	.	838.	623.000	1000.000
4	STATIC	LC	4290.	3110.000	5910.000	.	.	.	3510.	2570.000	4800.000
5	STATIC	LC	4050.	2620.000	6260.000	.	.	.	3400.	2600.000	4440.000
6	STATIC	LC	1160.	892.000	1550.000	.	.	.	884.	707.000	1100.000
7	STATIC	LC	7340.	4250.000	12700.000	.	.	.	7340.	4250.000	12700.000

179

ED_005172C_00000035-00181

CORREX

CAS # 52316-55-9 PRINCIPAL USE: FUNGICIDE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	98.10% TECHNICAL MATERIAL	RAINBOW TROUT		0.90	12.0	7.4	44
2	0.70 % LIQUID	RAINBOW TROUT		1.00	12.0	7.4	44
3	98.10% TECHNICAL MATERIAL	FATHEAD MINNOW		0.90	22.0	7.4	44
4	98.10% TECHNICAL MATERIAL	CHANNEL CATFISH		0.80	22.0	7.4	44
5	0.70 % LIQUID	CHANNEL CATFISH		0.80	22.0	7.4	44
6	98.10% TECHNICAL MATERIAL	BLUEGILL		0.90	22.0	7.4	44
7	0.70 % LIQUID	BLUEGILL		0.60	22.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>5600.	.	.	.	.	.	1600.	1240.000	2070.000
2	STATIC	LC	>3200.	.	.	.	.	.	560.	420.000	740.000
3	STATIC	LC	>10000	.	.	.	.	.	>10000	.	.
4	STATIC	LC	>320.	.	.	.	.	.	22.	14.400	33.600
5	STATIC	LC	>100.	.	.	.	.	.	19.	13.000	28.000
6	STATIC	LC	>10000	.	.	.	.	.	>10000	.	.
7	STATIC	LC		.	.	.	.	.	14000.	9350.000	21000.000

180

COUMAPHOS

ED_005172C_00000035-00182

CAS # 56-72-4 PRINCIPAL USE: INSECTICIDE

COUMAPHOS

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	95.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	15.0	7.1	44
2	95.00% TECHNICAL MATERIAL	GAMMARUS LACUSTRIS	MATURE	.	21.0	7.1	44
3	95.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.30	12.0	7.4	44
4	95.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.20	12.0	7.1	44
5	95.00% TECHNICAL MATERIAL	LAKE TROUT		2.10	12.0	7.4	162
6	95.00% TECHNICAL MATERIAL	CHANNEL CATFISH		1.00	18.0	7.4	44
7	95.00% TECHNICAL MATERIAL	BLUEGILL		1.30	18.0	7.1	44
8	95.00% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.90	18.0	7.4	272
9	95.00% TECHNICAL MATERIAL	WALLEYE		0.80	18.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	0.10	.	.	0.074	0.059	0.092
2	STATIC	LC	0.32	0.180	0.580		.	.	862.	645.000	1150.000
3	STATIC	LC	1090.	872.000	1360.000		.	.	890.	.	.
4	STATIC	LC	3000.	.	.		.	.	593.	416.000	846.000
5	STATIC	LC	998.	734.000	1360.000		.	.	840.	620.000	1140.000
6	STATIC	LC	5200.	2980.000	9070.000		.	.	340.	.	.
7	STATIC	LC	1100.	.	.		.	.	1100.	1000.000	1200.000
8	STATIC	LC	1500.	1100.000	2000.000		.	.	780.	645.000	943.000
9	STATIC	LC	1350.	1220.000	1490.000		.	.			

ED_005172C_00000035-00183

CROTOXYPHOS

CAS # 7700-17-16 PRINCIPAL USE: INSECTICIDE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	80.00% TECHNICAL MATERIAL	GAMMARUS LACUSTRIS	MATURE	.	15.0	7.1	44
2	80.00% TECHNICAL MATERIAL	PTEROHARCYS CALIFORNICA	1ST YEAR CLASS	.	15.0	7.1	44
3	80.00% TECHNICAL MATERIAL	CUTTHROAT TROUT		1.00	12.0	7.1	44
4	80.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.00	17.0	7.1	44
5	80.00% TECHNICAL MATERIAL	FATHEAD MINNOW		1.00	17.0	7.1	44
6	80.00% TECHNICAL MATERIAL	CHANNEL CATFISH		1.10	18.0	7.1	44
7	80.00% TECHNICAL MATERIAL	BLUEGILL		1.10	17.0	7.1	44
8	80.00% TECHNICAL MATERIAL	LARGEMOUTH BASS		0.70	18.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	49.0	36.000	67.000	29.0	21.000	41.000		.	.
2	STATIC	LC	6.6	.	.		.	.		.	.
3	STATIC	LC	92.0	49.000	170.000		.	.	51.0	28.000	91.000
4	STATIC	LC	101.	82.400	124.000		.	.	72.4	60.000	87.400
5	STATIC	LC	15500.	12800.000	18700.000		.	.	11900.	9830.000	14400.000
6	STATIC	LC	5000.	4260.000	5870.000		.	.	2600.	2240.000	3020.000
7	STATIC	LC	390.	338.000	450.000		.	.	152.	126.000	183.000
8	STATIC	LC	1800.	.	.		.	.	1100.	.	.

182

ED_005172C_00000035-00184

CRUFOMATE

CAS # 299-86-5 PRINCIPAL USE: INSECTICIDE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	92.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	IMMATURE	.	15.0	7.4	262
2	92.00% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	IMMATURE	.	15.0	7.4	44
3	92.00% TECHNICAL MATERIAL	BLUEGILL		1.00	18.0	7.4	44
4	92.00% TECHNICAL MATERIAL	BLUEGILL		1.00	15.0	7.4	272

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	5600.	4900.000	6400.000	.	.	.	3700.	3400.000	4100.000
2	STATIC	LC	240.	98.000	586.000	.	.	.	>100.	.	.
3	STATIC	LC	3750.	2550.000	5510.000	.	.	.	1800.	1340.000	2420.000
4	STATIC	LC	5100.	3500.000	7400.000	.	.	.	<3200.	.	.

ED_005172C_00000035-00185

CAS # 15096-52-3 PRINCIPAL USE: INSECTICIDE

CRYOLITE

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	96.00% TECHNICAL MATERIAL	DAPHNIA PULEX	1ST INSTAR	.	15.0	7.1	44
2	96.00% TECHNICAL MATERIAL	SIMOCEPHALUS SERRULATUS	1ST INSTAR	.	15.0	7.1	44
3	96.00% TECHNICAL MATERIAL	RAINBOW TROUT		1.80	12.0	7.1	44
4	96.00% TECHNICAL MATERIAL	BLUEGILL		0.80	24.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	10.0	7.600	13.000		.	.
2	STATIC	EC		.	.	5.00	3.600	6.800		.	.
3	STATIC	LC	160.	.	.		.	.	47.	.	.
4	STATIC	LC	>400.	.	.		.	.	>400.	.	.

181

ED_005172C_00000035-00186

CUMYLPHENYL DIPHENYL PHOSPHATE
MEASURED IN MG /L

CAS # 34364-42-6 PRINCIPAL USE: HYDRAULIC FLUID

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% LIQUID	RAINBOW TROUT		0.60	12.0	7.4	40
2	100.0% LIQUID	FATHEAD MINNOW		1.00	22.0	7.3	44
3	100.0% LIQUID	CHANNEL CATFISH		0.23	22.0	7.5	38

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
2	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.
3	STATIC	LC	>100.	.	.	.	.	.	>100.	.	.

185

ED_005172C_00000035-00187

CAS # 66555-33-7 PRINCIPAL USE: ALGICIDE

CUTRINE-PLUS
MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	9.00 % LIQUID	APPLE SNAIL	1-2DAY	.	22.0	7.4	42
2	9.00 % LIQUID	APPLE SNAIL	2-4WK	.	22.0	7.3	40
3	9.00 % LIQUID	APPLE SNAIL	2-4WK	.	27.0	7.5	42
4	9.00 % LIQUID	BLUEGILL		0.70	22.0	7.1	44
5	9.00 % LIQUID	BLUEGILL		0.70	22.0	7.1	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	.062	0.047	0.082	.	.	.	.036	0.030	0.043
2	STATIC	LC	.070	0.053	0.092	.	.	.	.048	0.034	0.068
3	STATIC	LC	.065	0.048	0.088	.	.	.	.048	0.034	0.067
4	STATIC	LC		.	.	.	.	.	2.3	1.910	2.770
5	STATIC	LC	2.8	2.000	3.900	.	.	.	1.32	1.000	1.750

186

ED_005172C_00000035-00188

CAS # 502-39-6 PRINCIPAL USE: FUNGICIDE

CYANO(METHYLMERCURI)GUANIDINE
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	2.20 % WETTABLE POWDER	RAINBOW TROUT		1.00	12.0	7.1	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	128.	96.000	172.000				55.	30.000	101.000

187

ED_005172C_00000035-00189

CYPRAZINE

CAS # 22936-86-3 PRINCIPAL USE: HERBICIDE

MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	12.80% EMULSIFIABLE CONCENTRATE	FATHEAD MINNOW		0.70	17.0	7.4	44
2	98.20% TECHNICAL MATERIAL	FATHEAD MINNOW		0.70	17.0	7.4	44
3	12.80% EMULSIFIABLE CONCENTRATE	BLUEGILL		1.20	17.0	7.4	44
4	98.20% TECHNICAL MATERIAL	BLUEGILL		1.20	17.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	1510.	1120.000	2010.000	.	.	.	942.	703.000	1026.000
2	STATIC	LC	21300.	14100.000	23100.000	.	.	.	13800.	9470.000	20000.000
3	STATIC	LC	1390.	1100.000	1760.000	.	.	.	1280.	1020.000	1600.000
4	STATIC	LC	31600.	28100.000	36000.000	.	.	.	30400.	27100.000	34200.000

CAS # 94-75-7 PRINCIPAL USE: HERBICIDE

2,4-D

MEASURED IN MG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.60	5.0	7.4	44
2	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.40	10.0	7.4	44
3	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.40	15.0	7.4	44
4	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.60	10.0	6.5	44
5	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50	10.0	7.5	44
6	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50	10.0	8.5	44
7	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.80	10.0	6.5	44
8	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.80	10.0	7.5	44
9	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.80	10.0	8.5	44
10	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50	10.0	7.4	44
11	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.80	10.0	7.4	160
12	100.0% TECHNICAL MATERIAL	CUTTHROAT TROUT		0.50	10.0	7.4	300
13	98.70% TECHNICAL MATERIAL	RAINBOW TROUT		0.30	12.0	7.7	40
14	100.0% TECHNICAL MATERIAL	LAKE TROUT		0.70	5.0	7.4	44
15	100.0% TECHNICAL MATERIAL	LAKE TROUT		0.30	10.0	7.4	44
16	100.0% TECHNICAL MATERIAL	LAKE TROUT		0.70	15.0	7.4	44
17	100.0% TECHNICAL MATERIAL	LAKE TROUT		0.70	10.0	6.5	44
18	100.0% TECHNICAL MATERIAL	LAKE TROUT		0.90	10.0	7.5	44
19	100.0% TECHNICAL MATERIAL	LAKE TROUT		0.70	10.0	8.5	44
20	98.70% TECHNICAL MATERIAL	FATHEAD MINNOW		0.90	22.0	7.3	40
21	98.70% TECHNICAL MATERIAL	BLUEGILL		0.50	22.0	7.3	40

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	LC	41.5	32.800	52.500	.	.	.	41.5	32.800	52.500
2	STATIC	LC	64.	57.300	71.500	.	.	.	64.	57.300	71.500
3	STATIC	LC	41.5	33.100	52.000	.	.	.	41.5	33.100	42.000
4	STATIC	LC	67.	57.500	78.000	.	.	.	67.	57.500	78.000
5	STATIC	LC	130.	100.000	170.000	.	.	.	130.	100.000	170.000
6	STATIC	LC	40.	30.700	52.100	.	.	.	40.	30.200	52.100
7	STATIC	LC	43.5	35.000	54.100	.	.	.	43.5	35.000	54.100
8	STATIC	LC	169.	146.000	195.000	.	.	.	169.	146.000	195.000
9	STATIC	LC	185.	165.000	207.000	.	.	.	172.	148.000	199.000
10	STATIC	LC	49.	40.400	59.400	.	.	.	44.	35.700	54.200
11	STATIC	LC	41.	32.400	51.900	.	.	.	37.	27.900	49.000
12	STATIC	LC	32.0	23.700	43.300	.	.	.	24.5	18.400	32.600
13	STATIC	LC		.	.	.	.	.	110.	77.000	157.000
14	STATIC	LC	65.7	47.200	91.400	.	.	.	65.7	47.200	91.400
15	STATIC	LC	44.5	35.300	56.100	.	.	.	44.5	35.300	56.100
16	STATIC	LC	64.	47.800	85.700	.	.	.	64.	47.800	85.700
17	STATIC	LC	63.	59.300	67.000	.	.	.	62.	54.000	71.200
18	STATIC	LC	105.	95.600	115.000	.	.	.	105.	95.600	115.000
19	STATIC	LC	127.5	110.000	149.000	.	.	.	120.	103.000	140.000
20	STATIC	LC		.	.	.	.	.	133.	103.000	171.000
21	STATIC	LC		.	.	.	.	.	180.	.	.

189

ED_005172C_00000035-00191

CAS # 1929-73-3 PRINCIPAL USE: HERBICIDE

2,4-D BUTOXYETHANOL ESTER
MEASURED IN UG /L

OBS	CHEMICAL DESCRIPTION	SPECIES	SIZE	WEIGHT	DEG. CENT.	PH	HARDNESS MG/L
1	62.50% TECHNICAL MATERIAL	DAPHNIA MAGNA	1ST INSTAR	.	21.0	7.4	44
2	60.80% LIQUID	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.4	43
3	29.00% GRANULAR FORMULATION	DAPHNIA MAGNA	1ST INSTAR	.	17.0	7.3	42
4	62.50% TECHNICAL MATERIAL	CYPRIDOPSIS VIDUA	EARLY INSTAR	.	21.0	7.4	272
5	62.50% TECHNICAL MATERIAL	ASELLUS BREVICAUDUS	EARLY INSTAR	.	15.0	7.4	272
6	62.50% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	IMMATURE	.	15.0	7.4	272
7	62.50% TECHNICAL MATERIAL	GAMMARUS FASCIATUS	IMMATURE	.	21.0	7.4	44
8	62.50% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	IMMATURE	.	21.0	7.4	44
9	62.50% TECHNICAL MATERIAL	PALAEMONETES KADIAKENSIS	IMMATURE	.	21.0	7.4	272
10	60.80% LIQUID	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	22.0	7.4	43
11	29.00% GRANULAR FORMULATION	CHIRONOMUS PLUMOSUS	3RD INSTAR	.	22.0	7.2	42
12	62.50% TECHNICAL MATERIAL	FATHEAD MINNOW		0.90	18.0	7.4	44
13	60.80% LIQUID	CHANNEL CATFISH		0.40	22.0	7.3	43
14	29.00% GRANULAR FORMULATION	CHANNEL CATFISH		0.40	22.0	7.3	43
15	62.50% TECHNICAL MATERIAL	BLUEGILL		1.40	18.0	7.4	272
16	60.80% LIQUID	BLUEGILL		0.60	22.0	7.4	42
17	29.00% GRANULAR FORMULATION	BLUEGILL		0.60	22.0	7.4	42
18	62.50% TECHNICAL MATERIAL	FOWLERS TOAD	TADPOLE		15.0	7.4	44

OBS	TEST TYPE		24 HOUR	CONFIDENCE INTERVAL		48 HOUR	CONFIDENCE INTERVAL		96 HOUR	CONFIDENCE INTERVAL	
1	STATIC	EC		.	.	6400.	4490.000	9120.000		.	.
2	STATIC	EC		.	.	1700.	1100.000	3700.000		.	.
3	STATIC	EC		.	.	4300.	3400.000	5500.000		.	.
4	STATIC	EC		.	.	2200.	1460.000	3320.000		.	.
5	STATIC	LC	>3200.	.	.		.	.	2600.	1280.000	5260.000
6	STATIC	LC	10500.	5730.000	19200.000		.	.	6100.	4480.000	8300.000
7	STATIC	LC	1400.	1100.000	1800.000		.	.	440.	310.000	620.000
8	STATIC	LC	>1000.	.	.		.	.		560.000	1000.000
9	STATIC	LC	>10000	.	.		.	.	<1000.	.	.
10	STATIC	EC		.	.	790.	640.000	970.000		.	.
11	STATIC	EC		.	.	390.	250.000	580.000		.	.
12	STATIC	LC	3250.	2540.000	4160.000		.	.	3250.	2540.000	4160.000
13	STATIC	LC	780.	600.000	1010.000		.	.	780.	600.000	1010.000
14	STATIC	LC	1350.	1060.000	1730.000		.	.	1350.	1060.000	1730.000
15	STATIC	LC	1340.	1220.000	1470.000		.	.	1200.	1100.000	1310.000
16	STATIC	LC	760.	590.000	970.000		.	.	760.	590.000	970.000
17	STATIC	LC	760.	600.000	970.000		.	.	760.	600.000	970.000
18	STATIC	LC	>10000	.	.		.	.	>10000	.	.

ED_005172C_00000035-00192