



Neonic Risk Assessments – Final Bee, Aquatics, & Terrestrial

(Clothianidin, Dinotefuran, Imidacloprid, Thiamethoxam)

Presentation for OPP OD

May X, 2019

Purpose

- Provide OPP upper management with an overall summary of neonic risk conclusions
- Preview of final bee risk assessments; recap of risk conclusions for aquatic and non-bee terrestrial taxa
- Provide an update on methodology and RTC for final bee assessment
- Highlight strengths (and potential weaknesses) of our assessments
- To provide detailed background scientific information prior to PRD's briefing on risk mitigation strategy

The goal of this presentation is to provide a summary of the final bee risk assessments

EFED Neonicotinoid Chemical Teams

Chemical	EFED Branch	Eco	Fate
Clothianidin	ERB 6	Michael Wagman	Chuck Peck
Thiamethoxam (combined document)	ERB 1	Kris Garber Ryan Mroz	Chris Koper
Imidacloprid	ERB 5	Keith Sappington Meghann Niesen Hannah Yingling	Mohammed Ruhman
Dinotefuran	ERB 3	Elizabeth Donovan	Rochelle Bohaty
Coordination and supporting roles		Colleen Rossmeisl Frank Farruggia Monica Wait	

Outline of Presentation

- Neonic RR timing
- Final Bee Risk Assessments
  - What's new since preliminary bee RAs [public comments, new data, nectar equivalents (replaces bee bread method), residue bridging strategy]
  - Lines of evidence considered / strength of confidence calls
  - Risk conclusions (low risk calls; weak, moderate, and strong evidence of risk calls)
- Recap of risk conclusions for aquatic taxa & new Guelph data analysis
- Recap of risk conclusions for birds and mammals
- Conclusions
  - Strengths and weaknesses of risk assessment approaches
  - Next steps

Neonic Registration Review Timing

- Preliminary Bee Risk Assessments
  - Imidacloprid published January 2016
  - Clothi / thia, dino published May 2017
- Non-pollinator (Aquatic & Bird, Mammal, Plants) RAs
  - Imidacloprid aquatic May 2017; terrestrial December 2017
  - Clothi, thia, dino published December 2017
- Final Bee RAs, RTC, and Proposed Interim Decisions
  - Signatures by June 30, 2019
  - Dockets to open shortly thereafter

5

Just a quick overview of the registration review timeline and a reminder that although the 4 neonics started registrant review at different times, they have all been aligned during the risk management phase.

MLW - NOT SURE WE NEED THE FOLLOWING
The preliminary bee assessment went out in 2016 and 2017; the non-pollinator assessments went out in 2017 and 2018; the final bee assessments, response to comments, and proposed interim decisions are planned for a March 2019 docket.

Final Bee Risk Assessment

6

Moving into the risk characterization…

Preliminary Bee RAs - Major Public Comments

- Comments that were addressed through modification of the risk assessment
  - Criticism of "bee bread" method and alternative suggestions
  - Lack of non-agricultural use risk assessment
  - Assessments were not adequate due to numerous "Uncertain" calls
- Comments on lack of assessment for less-typical exposure routes
  - Seed dust, soil exposure, drinking water, guttation fluid
- Other substantive comments that did not result in changes to the risk assessment (not exclusive to neonics)
  - Assessments don't consider mixtures, cumulative effects, or synergy
  - Honeybees are not appropriate surrogates
  - Not enough consideration for studies that include sublethal effects or non-apical endpoints (*e.g.,* immunosuppression, foraging ability, biochemical changes)

This slide summarizes the major categories of public comments received on the preliminary bee risk assessments. While these comments are generic to the four neonics being assessed, there are chemical-specific comments that have been incorporated into the risk assessments as appropriate. Major commenters were USDA, registrants, various crop groups, non-profits, and state agencies.

There were several comments that were addressed through modifications to the risk assessment. These included: criticism of the "bee bread" method and alternative suggestions for how to estimate exposure for the Tier II analysis. These comments informed development of the new method for estimating exposure, which we will talk about in a few slide. There were comments that the risk assessments did not consider non-ag uses; these uses are considered in the final assessments (risk calls will be discussed later). And finally, there were comments that the assessments were not adequate because of the numerous "uncertain calls". These calls in the preliminary assessments were due to gaps in the dataset, mainly for the tier II assessment. Since the drafts, we have received new colony feeding studies for the chemicals and residue data that have allowed us to update our higher tiered exposure assessment.
There were also comments on the lack of quantitative assessment for some of the less-typical exposure routes, such as, seed dust, soil exposure, drinking water, and guttation fluid. These routes are discussed qualitatively in the assessments both because the potential exposures are substantially less than those from dietary and contact exposure and because there aren't methods to quantify them. Of all of these less-typical routes, the most relevant is seed dust as there are numerous incidents associated with this type of exposure.
And finally, there are other substantive comments that didn not result in changes to the risk assessments. These are not exclusive to neonics and, ultimately, are not persuasive. Most of them relate to policy decisions and are being addressed with other work (e.g., synergy).

New Data Since Preliminary Bee RAs

- New colony feeding studies (CFS)
  - Sucrose: repeat clothianidin, dinotefuran, repeat thiamethoxam
  - Pollen patties: clothianidin pilot
- Additional pollen and nectar residue data
  - 9 for clothi, 8 for dino, 7 for imi, 11 for thia
- Evaluation of Tier 3 full field studies
  - Imidacloprid for pumpkin and cotton
- New open literature data considered

A considerable amount of data has been submitted since the preliminary risk assessments, including new colony feeding studies, residue data, Tier III data for imidacloprid, and open literature. I'll note that the colony feeding studies and residue data had a substantial influence on the risk assessments, while the Tier III and open literature did not.

New Methodology Since Preliminary Bee RAs

- Tier 2 methodology to combine residues in pollen and nectar (replaces "bee bread" method)
  - Revision is responsive to public comments, more biologically relevant, and consistent with Tier I approach
- The residue bridging strategy was developed
  - Distinguish with confidence between green and red calls
  - Ability to inform potential mitigation options
  - Analyzed data across X number of residue studies from all 4 neonics to determine trends; data rich compared to most other RR cases
- Included non-ag use sites (preliminary bee RA was ag only)

In addition to (and a result of) the new data, we have updated our Tier II methods for estimating colony-level exposures. We'll go into details in the next few slides, but at the high level, we developed a new approach to combining pollen and nectar residue that is responsive to public comments, more biologically relevant, and consistent with the approach used at Tier I. This approach replaces the somewhat controversial "bee bread" method. We also developed a residue bridging strategy that allows us to make the best use of the available residue data to 1) distinguish between green and red calls with some level of confidence, and 2) gives us the ability to inform on potential mitigation options, e.g., a pre-bloom interval that would preclude potential risk. And, as stated previously, we included the non-ag use sites as part of the risk assessment.

New Methodology – Tier 2 Pollen + Nectar

- Honey bee colonies consume more nectar than pollen
  - If concentrations in pollen and nectar are equal, dose from nectar will be greater
- Available information suggest that on a concentration basis, colony level endpoints for nectar should be lower than pollen
- Route of exposure does not appear to influence toxicity
- Three lines of evidence indicate that difference in contribution of colony's dose from pollen is 20x less than that of nectar
- Final equation:

$$C_{total} = C_{nectar} + \frac{C_{pollen}}{20}$$

And now for the detail

The Tier II pollen and nectar method is a way of combining measured concentrations for both matrices into a concentration in total diet that essentially converts residues in pollen to nectar equivalents by the application of a 20x factor. This factor was determined by evaluating three separate lines of evidence that all sort of converged. These are presented on the slide here and include: an evaluation of consumption rates and a comparison of tox endpoints on a concentration and dose basis.

The final equation used to estimate a total dietary concentration at the bottom was used in the risk assessments.

New Methodology – Residue Bridging Strategy

1. Using X number of neonic residue studies, we could develop method(s) to reduce uncertainties in existing neonic residue database for assessing risk to bees.
   a. **Lack of data** (missing chemical/crop/method)
   b. **Data limitations** (sparse temporal, spatial coverage, missing matrix)
2. Improve how residue data are applied to bee risk assessment
   1. Consistent level of protection
   2. Harmonize methodology with other taxa (birds/mammals), EFSA
3. Address residue data for applicable non-agricultural uses
4. There are separate residue bridging strategy documents that explain the methodology in detail that will be provided as attachments to the Final Bee RAs (1 – soil and foliar applications; 2 – seed treatment applications)

11

Moving on to the residues, the goals of the bridging strategy were to 1) develop methods to reduce uncertainties in the existing database due to lack of data or various data limitations; 2) improve how residues are applied to bee risk assessments by instituting a consistent level of protection and attempting to harmonize the methodology, where sufficient data were available, with those employed for other taxa or by other regulatory bodies; 3) and finally, to develop an approach for non-ag uses.

Distinct approaches were developed for seed treatments vs foliar/soil applications.

Seed Treatment Residue Bridging

- Derive Tier I (refined) and Tier II concentrations for crops with available data
  - Do variables related to site and year influence concentrations?
  - Does chemical influence concentrations?
  - Is there a relationship between other tissues and pollen and/or nectar?
- Use available data to set Tier I and II concentrations for crops with no residue data
  - Does crop influence magnitude of residue?
  - Consider crop groups? Use all data for all crops?

For seed treatments, there were two major questions: 1) For crops where you have some kind of data, can you bridge within crop from chemical to chemical? This includes bridging across site and year, chemical, and, in some cases, matrix (i.e., plant tissue); and 2) For crops without any data, can you bridge across crop and chemical.

Seed Treatment Bridging Conclusions

- Crop specific 90th percentile values (from all trials) recommended
  - Separate values for corn, cotton, soybean and canola
    - Data bridged across chemicals
  - Tier I (refined)
  - Tier II (when needed)

- For crops with no data, available data for corn, cotton, canola and soybean used to bridge to other crops
  - 90th percentiles used for all other crops, with adjustments for treatment rate

13

The recommendations based on the seed treatment bridging are: 1) for crops with available data, use the 90th percentile value from all chemicals. There are separate values for corn, cotton, soybean, and canola. These values are used to refine Tier I estimates, and Tier II only when needed (which was not very often). 2) for crops without data, the aggregate 90th percentile across all of the crops (i.e., corn, cotton, canola, and soybean) is used.

[90th percentile is the high end year/site conditions. 90th selected based on policy and level of protection we tend to pick. Consistent with aquatic eecs. Precedent.]

Foliar and Soil Residue Bridging

- Determined the major, quantifiable variables that influence residue levels
  - application method, application timing, and site
- Developed approach for incorporating residue data into risk assessment
  - Available residue data for a crop/crop group are adjusted to the maximum application rate and compared to the colony-level endpoints (NOAEC/LOAEC)
  - For those uses with sufficient data to quantify kinetics of residue declines (*i.e.,* foliar applications for cotton, cucurbit, and berries), Monte Carlo analyses were utilized
    - Probabilistic approach to predict residue concentrations
    - Allowed for calculation of the number of days required for residues to drop below the toxicity endpoint.
    - For orchards (pre-bloom, foliar), data were combined from available studies to derive median estimated values over time

14

Moving on to the foliar and soil applications. The residue bridging analyses identified application method, application timing, and site as the major, quantifiable variables that influence residue levels. I'll note that there are nuances to this and that while these variable consistently came up across the crops/crop groups, others were also identified on a case by case basis. Through this process we also developed an approach for how to incorporate the residue data into risk assessment. At a minimum, the colony level endpoints are compared to the residue data for each crop/crop group adjusted to the appropriate maximum application rate. Then, if there is sufficient data to quantify the kinetics of residue declines, a Monte Carlo analysis was conducted to predict a distribution of daily residue concentrations. This was only possible for foliar applications to cotton, cucurbits, and berries. This allowed for the calculation of the number of days required for residues to drop below the colony level endpoint. For foliar pre-bloom applications to orchards a slightly different approach was taken. In this case the residue data was not sufficient for use in the Monte Carlo analysis, but it was determined that it was possible to combine all of the available data to derive median estimated values over time.

General Trends in Residue Data

- Residues from foliar applications > soil applications > seed treatments
  - Residues from foliar applications = 100 - 1000s ppb
  - Residues from soil applications = 10 - 100s ppb
  - Residues from seed treatments = 1 - 10s ppb
- Although residues from foliar applications are generally higher in magnitude, they decline more rapidly than soil applications (which tend to persist in pollen and nectar for much longer time periods)
- Pre-bloom applications result in residues that are generally much higher than post-bloom applications

15

Based on these analyses we saw some general trends in the data. At the 30,000 foot level, residues from foliar applications are greater than residues from soil applications, which are greater than residues from seed treatments. [Ranges presented here represent the max values normalized to 0.1 lb/a for foliar and soil applications and 1 mg/seed for seed treatments.] I'll note that the range of residues presented for foliar applications is based on samples taken close to application (~2 weeks). After that the second bullet comes into play because residues from foliar applications tend to decline much more rapidly than residues from soil applications, with a steeper slope. Generally there is also a distinction between pre-bloom and post-bloom applications, with the former being greater.

Based on these general trends we decided to separated foliar and soil applications as well as pre-bloom and post-bloom applications. You'll see how this factors into the risk calls in a few slides.

Foliar and Soil Bridging Conclusions

- Based on the bridging analysis, the following crop/crop groups are bridged across chemicals
  - Cotton (THIA only)
  - Cucurbits
  - Orchards (Stone fruit, pome, citrus, tree nuts – DID I MISS ANY?)
    - Residues are also bridged across orchard crop groups
  - Berries and small fruits
    - Orchard data used as additional line of evidence
  - Soybeans
- For herbaceous crops/crop groups with insufficient data
  - Cotton and cucurbit data used as surrogates
  - Relevant crop groups
    - Root and tuber (*e.g.,* sweet potato)
    - Fruiting vegetables (*e.g.,* okra and roselle)
    - Herbs and spices

16

Based on the bridging analysis for foliar and soil applications, cotton, cucurbits, orchards, berries and small fruits, and soybeans are bridged across all chemicals. A few things to note: cotton data is bridged across chemicals for THIA only because there was sufficient data for IMI, CLOTHI, and DINO to use chemical-specific data. Orchards represents an artificial grouping of various tree crop groups (e.g., citrus, stone, pome, tree nut). Residues are bridged across all orchard crop groups.

For herbaceous crops/crop groups with insufficient data, cotton and cucurbit are used as surrogates. These include the honey bee attractive root and tubers and fruiting vegetables. We'll get into more details on this a bit later.

{HED didn't have any issues with approach}

Final Bee Risk Assessment

Risk Characterization

17

Moving into the risk characterization…

Lines of Evidence Considered

- Pollinator Attractiveness
- Agronomic practices (*e.g.,* harvest time)
- Residues above NOAEC and LOAEC
  - Chemical / crop specific residue data
  - Bridging strategy residue data (from other crops or across chemicals)
  - Duration and frequency of LOC exceedance
  - Magnitude of exceedance
    - Ratio of max residue value to NOAEC/LOAEC
    - % of diet from the treated field needed to reach the NOAEC/LOAEC
  - Geographical scale and spatial distribution of crop group / use pattern; usage
- Incidents

Several lines of evidence were considered in making risk calls. At Tier I, we considered whether the crop was honey bee attractive, as well as any agronomic practices that may limit exposure, e.g., harvest time or flower tenting (as in the case of mandarin oranges). At Tier I we also considered whether predicted or measured residues exceed the individual effect level endpoints, which they mostly did. At the Tier II we considered whether residues exceeded the colony level endpoints. This was the major basis for a risk call; however, we provided several additional pieces of information to better characterize the potential for risk, including whether the exceedances were based on chemical specific or bridged residue data; the duration, frequency, and magnitude of the exceedance; and the geographic scale and spatial distribution of the use. And then there are the incidents. The teams are working on incorporating this additional characterization into a discussion of the strength of the risk call for each crop/crop group within a chemical.

Seed treatments: use and usage

- Registered for variety of crops on imidacloprid, clothianidin and thiamethoxam
  - Not registered on dinotefuran
- Majority of pounds applied and acres treated
  - Corn and soybeans represent majority of usage





Starting with use and usage. Seed treatments are registered for a variety of crops on IMI, CLOTHI, and THIA. They are not registered for DINO. The three pie graphs show the average acres treated (calculated by multiplying acres grown by average PCT from SLUA)of seed treatments versus all other uses registered for IMI, CLOTHI, and THIA. These highlight just how much of the use is seed treatment, which, spoiler alert, is mostly considered low risk. However, it is important to note that numerous bee kill incidents have been associated with dust-off for each chemical: IMI has 5 reported incidents (canola, corn, soybean) from 2006 to 2016; THIA has 2 reported incidents that we can associate with seed treatment in the US (IN and MN- the magnitude of these incidents is on the order of thousands of hives) in 2012 with additional international incidents. It is possible that some of the other reports were associated with seed treatment, but we cannot confirm due to lack of details in the reports. CLOTHI has 18 incidents (corn or general ag areas) from 2010 to 2016 with numerous additional international incidents.

These incidents highlight the potential for effects and the large spatial scale of these uses.

To get a sense for the specific crops with the greatest usage we have the figure on the right, which shows the lbs applied for each chemical by crop. Corn and soybean pop out as representing the majority of the usage.

Usage for foliar/soil applications



20

Moving on to the foliar and soil applications, this figure shows the average acres treated (calculated the same was as for seed treatments) for all of the low risk crops/crop groups. For foliar/soil aps, the most usage is on soybeans, cereal grains. Take note of the y-axis before we transition to the next slide.



This shows the average acres treated for the risk crops. These data are for uses that have at least one red risk call so there are some nuances that don't translate, but in general, the crops with the highest acres treated are cotton, berries, and citrus. If you recall the scale from the previous slide you can see that the acreage for the risk crops is substantially less than the acreage for the low risk crops.

[For some of the orchard crop data, it is unknown whether usage was pre- or post- bloom. So, some of these acres treated may be green. E.g., clothi use on pome stone and tree nuts.]

Low Risk Calls

Foliar and Soil Applications

Crop Group or Crop	IMI		CLOTHI		THIA		DINO	
	Foliar	Soil	Foliar	Soil	Foliar	Soil	Foliar	Soil
Bulb Vegetables								
Leafy Vegetables								
Brassica Vegetables								
Legumes								
Cereal Grains								
Cucurbits								
Citrus Fruits	**	**	Post-	Post-	Post-	Post-		
Pome Fruits			Post-		Post-			
Stone Fruits			Post-		Post-		Post-	Pre-/Post-
Tree Nuts	Post-		Post-		Post-			
Tropical Fruits			Post-		Post-			
Berries/Small Fruits	Post-	Post-	Post-	Post-	Post-	Post-	Post-	Post-
Root/Tubers*								
Fruiting Veg*								

Seed Treatments

Crop Group or Crop	IMI	CLOTHI	THIA
Bulb Vegetables			
Leafy Vegetables			
Brassica Vegetables			
Legumes			
Cereal Grains			
Oilseed			
Cucurbit Vegetables			
Root/Tuber Vegetables*			

* Denotes call is for non-attractive crops
** Crop tented during bloom

22

This table summarizes the low risk calls for foliar and soil applications, represented by green cells. The gray cells indicate either the chemical is not registered for a particular use or there was a risk call (we'll get to those soon). A few things to note: the calls for root and tubers and fruiting vegetables are for non-attractive crops within the groups, and the call for IMI citrus is only for mandarin oranges… all other citrus are high for both foliar and soil applications.

The table on the right summarizes the low risk calls for seed treatments, which as we discussed on the previous slide, account for the large majority of usage for imi, clothi, and dino. So things like soybean, corn, which are major uses for these chemicals, were identified as low risk (not accounting for dust-off)

MLW – also make the point that the additional data generated for these 3 chemicals resulted in some previously "uncertain risk" groups were moved to the low risk category

Low Risk Calls

- Harvested prior to bloom
  - Bulb, leafy and brassica vegetables; artichoke and tobacco
- Not attractive to honey bees
  - Root and tuber, fruiting vegetables
- Residues below the colony-level effects endpoint
  - All seed treatments:
    - Except CLOTHI turmeric and IMI peanut and bean
  - Foliar applications:
    - Legumes
    - post-bloom applications
      - Orchards (except IMI stone and pome)
      - Berries and small fruits (except CLOTHI soil)
  - Soil applications:
    - Dino cucurbits

Residues in orchard following post-bloom foliar applications vs. the dinotefuran endpoint



23

Crops/crop groups were considered low risk because they were harvested prior to bloom (see list), not considered attractive to honey bees (i.e., certain crops within the root and tuber and fruiting vegetables crop groups), or had measured residues below the colony-level effects endpoints. The figure on the left is an example for foliar post-bloom applications to orchard crops where the residues are substantially lower than the dinotefuran colony level NOAEC.

Risk Calls

- Cotton
  - Foliar: IMI; CLOTHI; THIA; DINO
  - Soil: IMI
- Cucurbits
  - Foliar: CLOTHI; THIA; DINO
  - Soil: IMI; CLOTHI, THIA
- Orchards
  - Foliar Pre-bloom: IMI; CLOTHI; THIA; DINO
  - Foliar Post-bloom: IMI (except tree nuts)
  - Soil Pre-, Post-bloom: IMI; CLOTHI; THIA
- Berries and Small Fruits
  - Foliar Pre-bloom: IMI; CLOTHI; THIA; DINO
  - Soil Pre-bloom: IMI; THIA; DINO
- Other Herbaceous Crops
  - Root and Tuber
  - Fruiting Vegetables
  - Herbs and Spices
- Seed Treatments
  - Turmeric: CLOTHI
  - Bean and Peanut: IMI
- Ornamentals and Forestry
  - Foliar/Soil: IMI; CLOTHI; THIA; DINO
  - Trunk Injection: IMI; DINO
- Turf

24

Moving to the risk calls… We'll walk through the details in the following slides, but risk calls were made for all of the uses shown here. As discussed previously, risk calls are based on residue values exceeding the NOAEC. However, all of the lines of evidence were used to characterize the strength of the risk call.

Evidence of Risk

- Strong Evidence of Risk
  - Residues exceed colony-level endpoint(s) by a high magnitude, frequency, or duration
  - May be supported by modeled exposures or ecological incidents
- Moderate Evidence of Risk
  - Residues exceed colony-level endpoint(s) but magnitude, frequency, or duration are limited
- Weak Evidence of Risk
  - Residues exceed colony-level endpoint(s) but there are uncertainties in surrogacy

Risk Calls – Strong Evidence of Risk

Imidacloprid	Clothianidin	Thiamethoxam	Dinotefuran
			cotton (foliar) **stone** (foliar, pre-bloom) **berries** (foliar, pre-bloom), **attractive fruiting vegetables** (foliar), and **attractive ornamentals and forest trees** (foliar, soil, trunk injection)

Risk Calls – Moderate Evidence of Risk

Imidacloprid	Clothianidin	Thiamethoxam	Dinotefuran
			cucurbits (foliar), **berries** (soil, pre-bloom), and **turf** (residential lawns with bee-attractive blooming weeds).

Risk Calls – Strong Evidence of Risk

Imidacloprid	Clothianidin	Thiamethoxam	Dinotefuran
			attractive root/tubers (foliar, soil), **attractive fruiting vegetables** (soil), and **stone** (soil, pre-bloom).

Risk Calls- Cotton Foliar

Lines of Evidence	Imidacloprid		Clothianidin		Thiamethoxam		Dinotefuran
Residue Data	CS		CS		B		CS
Endpoint	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC*
Frequency (# of daily mean residues > endpoint)	2 (FN) 0 (XFN)	0 (FN) 0 (XFN)	8 (FN) 28 (XFN)	4 (FN) 26 (XFN)	0 (FN) 4 (XFN)	0 (FN) 2 (XFN)	2 (FN) 11 (XFN)
Magnitude (Ratio of max residue to endpoint)	**1.1**X (FN) **1.5**X (XFN)	0.5X (FN) 0.7X (XFN)	**7.7**X (FN) **412** (XFN)	**4.0**X (FN) **220**X (XFN)	0.24X (FN) **3.4**X (XFN)	0.12X (FN) **1.7**X (XFN)	**9.7**X (FN) **177**X (XFN)
Duration (# of days > endpoint)	20 (FN) 22 (XFN)	8 (FN) 7 (XFN)	9 (FN) 49 (XFN)	7 (FN) 49 (XFN)	8 (FN) 20 (XFN)	6 (FN) 17 (XFN)	10 (FN) 35 (XFN)
Incidents	None		3 honey bee kills		None		None

Additional considerations for cotton:

- Considered attractive for floral nectar, potentially attractive for extrafloral nectar, and not attractive for pollen
- Has an indeterminant bloom period
- Does not require managed pollinators but used for honey production by some commercial beekeepers

23

CS = Chemical Specific Data, B=Bridged Data

The next few slides have a similar format so I will take some time to orient folks. These tables summarize the information included in the lines of evidence tables presented in the Tier II risk characterizations. The rows are the lines of evidence, including, whether risk conclusions are based on chemical specific residues, bridged residues, or both; the endpoint under consideration (NOAEC or LOAEC); the frequency of exceedances based on the number of daily mean residues that are greater than the endpoint [note, we are presenting absolute number of residues exceeding, but it is important to note that some have a lot of data and some don't]; the magnitude of the exceedances as presented by the ratio of the max residue to the endpoint; the duration of exceedances; and the incidents. The columns are the individual chemicals. I know they don't look it, but these tables are actually pretty high level and roll up a lot of information. The nuances are described in the documents.

A few considerations: Given the importance of the THIA degradate (i.e., CLOTHI), residue data are typically compared to both the thia and clothi endpoints (you'll see these presented in parentheses in the following slides). However, for cotton we are not considering the clothi endpoints due to low percent of clothi present in studies with thia. A definitive LOAEC is not available for DINO, but based on transient effects observed at the NOAEC for all of the neonics, the LOAEC is likely with a factor of 5X the NOAEC.

Ok so moving into the risk call for cotton. pollen is not considered honeybee attractive so it is not considered in the residues (i.e., we did not calculate a total concentration in diet based on the method described previously). Somewhat unique to cotton, is that plants produce both floral nectar and extrafloral nectar. Extrafloral nectar is basically nectar produced by the plant from nectaries that occur on stems rather than flowers. It is produced even when the plant is not flowering. While floral nectar is considered honey bee attractive, the attractiveness of extrafloral nectar is uncertain. However, based on similarities in composition [and a few open literature sources], it is reasonable to assume bees are attracted to extrafloral nectar, though the extent and duration of its use is uncertain. For the purposes of the risk assessment, we present risks for both floral and extrafloral nectar.

As mentioned previously. There was sufficient data to conduct monte carlo analyses based on chemical-specific data for IMI, CLOTHI, and DINO, while THIA relies on bridged data from the other three chemicals. In general, the frequency, duration, and magnitude of exceedances are greater for extrafloral nectar than floral nectar. Cotton does not require managed pollinators but commercial beekeepers do utilize cotton fields for their hives so there is potential for exposure. And there are confirmed incidents for CLOTHI that indicate this is a relevant route of exposure.

There is strong evidence of risk based on magnitude of exceedances (up to 412 for extrafloral nectar) and duration of exceedances (from weeks to months).

Risk Calls- Cucurbit Foliar

Lines of Evidence	Clothianidin		Thiamethoxam*		Dinotefuran
Residue Data	CS, B		CS, B		B
Endpoint	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC*
Frequency (# of daily mean residues > endpoint)	15	12	8 (14)	5 (10)	3
Magnitude (Ratio of max residue to endpoint)	**15.6X**	**8.3X**	**5X (11X)**	**2.5X (6.3X)**	**3.2X**
Duration (# of days > endpoint)	26	17	16 (22)	6 (13)	15
Incidents	None		None		None

Additional considerations for cucurbits:

- Considered highly attractive for nectar and pollen
- Has an indeterminant bloom period
- Requires managed pollinators

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

30

Here is the lines of evidence table for foliar applications to cucurbits. Note that imi is not registered for foliar applications which is why it is not in the table and we are presenting results for thia based on both the thia and clothi endpoints.

Cucurbits are pollinator attractive and do use managed pollinators. There are no reported incidents.

There is strong evidence of risk for clothi and thia based on exceedance of chemical-specific and bridged residues, the magnitude of those exceedances (up to 15x), and the number of days above the endpoint on the order of weeks. For DINO the residues exceed the bridged residue only and the magnitude of exceedance is lower, so the evidence of risk isn't as strong. [Dino-specific data are not suitable for use in risk assessment because the nectar and sometimes pollen were collected from in-hive sources].

Risk Calls- Cucurbit Soil

Lines of Evidence	Imidacloprid		Clothianidin		Thiamethoxam*	
Residue Data	CS, B		CS, B		CS, B	
Endpoint	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC	LOAEC
Frequency (# of daily mean residues > endpoint)	25	9	13	7	1 (9)	0 (1)
Magnitude (Ratio of max residue to endpoint)	**5.5**X	**2.7**X	**3.6**X	1.1X	0.69X (**1.5**X)	0.57X (**1.4**X)
Duration (# of days > endpoint)	67	57	57	47	34 (47)	0 (34)
Incidents	None		None		None	

Additional considerations for cucurbits:

- Considered highly attractive for nectar and pollen
- Has an indeterminant bloom period
- Requires managed pollinators

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

31

Moving on the soil applications to cucurbits. Note dino soil applications are considered “low risk”, which is why its not in the table.

Compared to the foliar applications, the magnitude of exceedances are lower, but the duration of those exceedances is much longer (months vs. weeks) This will be a common theme for the crop groups with both foliar and soil applications.

Risk Calls- Orchards Foliar

Lines of Evidence	Imidacloprid				Thiamethoxam*		Dinotefuran
Application Timing	Pre-bloom (Citrus)		Post-bloom (Stone/Pome)		Pre-bloom		Pre-bloom
Residue Data	CS, B		CS, B		CS, B		B
Endpoint	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC*
Frequency (# of daily mean residues > endpoint)	24	22	4	0	20 (23)	11 (21)	8
Magnitude (Ratio of max residue to endpoint)	**166**X	**80**X	**1.5**X	0.7X	**33**X (**76**X)	**19**X (**47**X)	**16.6**X
Duration (# of days > endpoint)	46	34	200+	0	35 (35)	18 (27)	14
Incidents	2 bee kill incidents on orchard crops				13 bee kill incidents on orchard crops		None

Additional considerations for orchards:

- Considered attractive or highly attractive for nectar and pollen
- Bloom duration varies by crop and variety
- Requires managed pollinators and used for honey production by some commercial beekeepers

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

32

Here is the risk table for foliar applications to orchards. Not that pre-bloom and post-bloom applications are included for imi, while post-bloom applications for thia and dino are considered low risk. Also note that Clothi is not registered for pre-bloom foliar applications.

Orchards are considered attractive to highly attractive, require managed pollinators and may be used for honey production by some commercial beekeepers. There is strong evidence of risk for imi and thia pre-bloom application based on exceedance of chemical-specific and bridged residues, the magnitude of those exceedances (up to 166x for NOAEC and 80x for LOAEC), and the number of days above the endpoint on the order of months. There are also reported incidents for both imi and thia that confirm the route of exposure. While the incidents are not tied to a particular application method or timing, based on the measured residues it is reasonable to infer they are the result of foliar applications.

Risk Calls- Orchards Soil

Lines of Evidence	Imidacloprid		Clothianidin		Thiamethoxam*	
Application Timing	Soil: Pre-/Post-bloom		Soil: Pre-/Post-bloom		Soil: Pre-/Post-bloom	
Residue Data	CS, B		CS, B		CS, B	
Endpoint	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC	LOAEC
Frequency (# of daily mean residues > endpoint)	63	24	23	12	5 (16)	1 (8)
Magnitude (Ratio of max residue to endpoint)	**3.6-19X**	**1.7-8.9X**	**9.1X**	**4.8X**	**2.7X (6.5X)**	**1.5X (3.5X)**
Duration (# of days > endpoint)	186	179	180	156	60 (156)	20 (156)
Incidents	2 bee kill incidents on orchard crops		None		13 bee kill incidents on orchard crops	

Additional considerations for orchards:

- Considered attractive or highly attractive for nectar and pollen
- Bloom duration varies by crop and variety
- Requires managed pollinators and used for honey production by some commercial beekeepers

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

33

Now for soil pre and post bloom soil applications to orchards. Dino soil applications are considered LOW risk so they are not presented here.

As we've seen with other soil applications, the magnitude of exceedances are lower than foliar (up to 19x for NOAEC and 8.9x for LOAEC), but the duration of those exceedances is much longer (up to 6 months in some cases).

Risk Calls- Berries and Small Fruits Foliar (Pre-bloom)

Lines of Evidence	Imidacloprid		Clothianidin		Thiamethoxam*		Dinotefuran
Application Timing	Pre-bloom		Pre-bloom		Pre-bloom		Pre-bloom
Residue Data	B		CS		CS, B		CS, B
Endpoint	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC*
Frequency (# of daily mean residues > endpoint)	42	39	4	3	32 (37)	23 (33)	36
Magnitude (Ratio of max residue to endpoint)	**63X**	**30X**	**3.4X**	**1.8X**	**20X (50X)**	**12X (28X)**	**25X**
Duration (# of days > endpoint)	57	45	37	37	23 (31)	19 (24)	41
Incidents	None		None		None		None

Additional considerations for berries and small fruits:

- Considered attractive or highly attractive for nectar and pollen
- Bloom duration varies by crop and variety
- Requires managed pollinators and used for honey production by some commercial beekeepers

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

34

Here are the lines of evidence for pre-bloom foliar applications to berries and small fruits. Note that CLOTHI is only registered for pre-bloom applications to grapes. All post-bloom foliar applications are considered low risk.

Berries and small fruits is a diverse group, but for the most part crops are considered attractive to highly attractive for nectar and pollen, require managed pollinators, and may be used for honey production. Bloom duration varies but is at least a couple of weeks. There is strong evidence of risk for all chemicals. Duration of exceedances on the order of months.

Risk Calls- Berries and Small Fruits Soil (Pre-bloom)

Lines of Evidence	Imidacloprid		Thiamethoxam		Dinotefuran
Application Timing	Pre-bloom		Pre-bloom		Pre-bloom
Residue Data	B		CS, B		B
Endpoint	NOAEC	LOAEC	NOAEC	LOAEC	NOAEC*
Frequency (# of daily mean residues > endpoint)	15	14	5 (11)	2 (8)	4
Magnitude (Ratio of max residue to endpoint)	**21X**	**9.9X**	**3.2X (7.7X)**	**2.7X (6.1X)**	**2.4X**
Duration (# of days > endpoint)	83	83	83	83	60
Incidents	None		None		None

Additional considerations for berries and small fruits:

- Considered attractive or highly attractive for nectar and pollen
- Bloom duration varies by crop and variety
- Requires managed pollinators and used for honey production by some commercial beekeepers

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

33

Moving to pre-bloom soil applications. Again, clothi is only registered for pre-bloom applications to grape and is considered LOW risk based on chemical-specific residue data. For the other chemicals, risk is indicated. Again, the magnitude of exceedances are lower than foliar, but those exceedances occur for about twice as long.

Risk Calls- Berries and Small Fruits Soil (Post-bloom)

- Risk indicated for CLOTHI and THIA
- Conflicting lines of evidence
  - Residues of IMI in blueberry below colony-level endpoints
  - Residues bridged from orchards exceed colony-level endpoints

36

Risk was also indicated for post-bloom soil applications for clothi and thia only. However, this is based on conflicting lines of evidence and would be an example of weak evidence of risk.

Risk Calls – Other Herbaceous Crops

- Members from Fruiting Vegetable[1], Root/Tuber[1], Herbs and Spices crop groups producing honeybee attractive pollen and nectar
  - Honeybee attractive crops include: **sweet potato, okra, herbs and spices, roselle**
  - Potential for Risk for all chemicals is bridged from all herbaceous plant data including, cotton and cucurbits
    - Based on plant relationships (*e.g.* at the family level)
    - High risk is indicated for both cucurbit (foliar and soil) and oilseed crops (foliar)
    - For pre-bloom neonic applications residues exceed colony level endpoints
- Members from Fruiting Vegetable[1] crop group producing honeybee attractive pollen only
  - Honeybee attractive crops include: **Chili peppers**
  - Potential for Risk for all chemicals is bridged from fruiting vegetable data including, tomato, bell and chili pepper
    - Based on plant relationships (*e.g.* at the genus level)
    - Foliar – Pollen residues in tomato for thia and dino exceed colony level NOAECs
    - Soil – Pollen residues in bell pepper, chili pepper, and tomato exceed colony level NOAECs
    - Supported by exceedances in other crop groups

37

[1] the majority of crops in these groups are harvested prior to bloom or are not attractive to honeybees. These are generally bumblebee attractive.

In addition to the crop groups already discussed, there are a variety of other herbaceous crops that are considered honeybee attractive, including sweet potato, okra, roselle, chili peppers, and herbs and spices. Given the lack of crop-specific residues, data are bridged to other crops/crop groups based on taxonomic relationships (e.g., family or genus level similarities). Risk is indicated based on surrogate residue values exceeding the colony level endpoints.

These crops do not use managed pollinators.

Risk Calls – Seed Treatment Uses

- Turmeric (CLOTHI); bean and peanut (IMI)
  - Residues reflect a high-end estimate (e.g., 90th percentile);
  - Estimate derived from multiple crops and neonicotinoids, and
  - IMI- residues are between the colony-level NOAEC and LOAEC
  - CLOTHI- uncertain how representative residue data are for treated root/rhizome

38

There are also a few seed treatment uses that exceed the risk threshold; however there are uncertainties associated with those calls, including residue values falling between the colony-level NOAEC and LOAEC for IMI uses on bean and peanut, and how representative available residue data are for treated root/rhizome as in the case for CLOTHI use on turmeric.

Risk Calls – Non-Ag Uses

- Ornamentals and forestry
  - Residue data available for thiamethoxam, imidacloprid, and dinotefuran
    - Residues for ornamental species used for forestry uses
  - Risk indicated for ornamentals (all chemicals) and forestry (imi, dino)
  - Incidents for IMI, CLOTHI, and DINO
  - Uncertainty considerations:
    - Very limited data set for a diverse set of plants
      - Unable to refine exceedances based on time
    - Application rates and scaling to lb/A for a standard evaluation
- Turfgrass (residential): risk is indicated for all chemicals
  - Residues from open lit. study with IMI and CLOTHI
  - Based on the assumption of flowering weeds on residential lawns

33

For non-agricultural ornamentals and forestry uses, residue data are available for IMI (soil), THIA (foliar and soil), and DINO (foliar and trunk injection). While ornamental plants are of known pollinator attractiveness just from everyone observing bees in our gardens, IR-4 has developed a website that contains a list to help gardeners find ornamental plants they can use in their gardens to serve as a resource for foraging pollinators. In addition, various tree species are considered bee attractive, e.g., maple, serviceberry, crapemyrtle, black tupelo, sourwood, black locust, and linden. In addition, an article by Hill and Webster (1995) discusses the potential economic benefits of combining apiculture and forestry operations as many of the commercially valuable trees produce nectar and pollen that are available during the spring, when other bee resources are limited.

The assessment for residential turf assumes that bee attractive weeds are present and flowering during application.

While there are notable uncertainties, for these uses residue levels are in the PPM range. These large exceedances are likely much greater than any chemical specific influence we would see (based on foliar/soil residue). Therefore, risk is indicated for ornamental, forestry, and turf uses.

Tier I Risk Conclusions

- Clothi poultry house use high risk based on Tier I
  - No residue data available
  - Incorporates proposed mitigation scenarios from the registrant
- Spray drift- no change from preliminary

In addition to the Tier II risk conclusions there Tier I conclusions that have not changed from the preliminary assessments, including conclusions for the clothi poultry and risks from spray drift.

Summary of risk conclusions for foliar applications

Crop Group or Crop	Imidacloprid		Clothianidin		Thiamethoxam		Dinotefuran	
Bulb Vegetables								
Leafy Vegetables								
Brassica Vegetables								
Legumes								
Cereal Grains								
Cotton								
Cucurbit Vegetables								
Citrus Fruits	Pre-	Post-			Pre-	Post-		
Pome Fruits	Pre-	Post-	Pre-	Post-	Pre-	Post-		
Stone Fruits	Pre-	Post-	Pre-	Post-	Pre-	Post-	Pre-	Post-
Tree Nuts	Pre-	Post	Pre-	Post-	Pre-	Post-		
Tropical Fruits	Pre-	Post-	Pre-	Post-	Pre-	Post-		
Berries/Small Fruits	Pre-	Post-	Pre-	Post-	Pre-	Post-	Pre-	Post-
Root/Tubers Vegetables*								
Fruiting Vegetables*								
Herbs/Spices								

41

* denotes call is for honeybee attractive crops within the crop group

The next few tables summarize the risk calls for agricultural crops. This table summarizes the risk conclusions for foliar applications. Red cells are risk, green cells are low risk, and gray cells are not registered. For orchards and berries and small fruits, risk calls are distinguished for pre-bloom vs. post-bloom applications. Note that most of these calls were yellow in the preliminary assessments due to gaps in the residue database. Bridging really allowed us to make them all green or red.

Summary of risk conclusions for soil applications

Crop Group or Crop	Imidacloprid		Clothianidin		Thiamethoxam		Dinotefuran	
Bulb Vegetables								
Leafy Vegetables								
Brassica Vegetables								
Legumes								
Cereal Grains								
Cotton								
Cucurbit Vegetables								
Citrus Fruits	Pre-	Post-	Pre-	Post-	Pre-	Post-		
Pome Fruits	Pre-	Post-						
Stone Fruits	Pre-	Post-					Pre-	Post-
Tree Nuts	Pre-	Post-						
Tropical Fruits	Pre-	Post-						
Berries/Small Fruits	Pre-	Post-	Pre-	Post-	Pre-	Post-	Pre-	Post-
Root/Tubers Vegetables*								
Fruiting Vegetables*								
Herbs/Spices								

4.2

* denotes call is for honeybee attractive crops within the crop group

Here is the table summarizing risk conclusions for soil applications.

Summary of risk conclusions for seed treatment applications

Crop Group or Crop	Imidacloprid	Clothianidin	Thiamethoxam
Bulb Vegetables			
Leafy Vegetables			
Brassica Vegetables			
Legumes			
Cereal Grains			
Oilseed			
Cucurbit Vegetables			
Citrus Fruits			
Pome Fruits			
Stone Fruits			
Tree Nuts			
Tropical Fruits			
Berries/Small Fruits			
Root/Tubers Vegetables*			
Fruiting Vegetables*			
Herbs/Spices			

43

* denotes call is for honeybee attractive crops within the crop group

Here is the table summarizing the risk conclusions for seed treatment uses. Yellow cells indicate potential risk.

Non-*Apis* Risk Conclusions

- Comparison of tox data and previous analysis of exposure indicate that honeybees are an appropriate surrogate
  - Red calls for honey bees extend to non-Apis
  - Fruiting vegetable green calls for honey bees are red for non-Apis

Surrogate both at individual and colony level

Summary of Major Changes and Risk Conclusions

- Numerous potential risk calls in draft bee assessments due to gaps in dataset.
- Final bee assessments incorporate:
  - New CFS data
  - New residue data
  - Updated higher tiered exposure assessment
    - Tier 2 methodology to combine residues in pollen and nectar
    - Residue bridging strategy
- General risk conclusions
  - Foliar and soil applications to bee attractive plants = risk

45

As discussed through the presentation, additional data received since the preliminary assessments allowed us to refine the yellow risk calls. The general conclusion is that if foliar and soil applications are made to bee attractive plants there is risk.

Aquatic Taxa Risk Conclusions
& New Guelph Data Analysis

46

Moving into the risk characterization…

New Data Set – Guelph (Raby et al) Aquatic Invert Toxicity Data

- Large acute and chronic datasets across all 4 neonics and acetamiprid
- Became publicly available on x date
- X species tested for acute, and X species test for chronic; including a range of species' sensitivities
- Allowed for apples-to-apples comparison of toxicity data across the 4 neonics accounting for lab and study conduct variability

Big Picture Risk Conclusions

Aquatic invertebrates

- *Foliar, Soil, Seed Treatment:*
  - Acute and chronic risk concerns for all 4 chemicals (some ➔ most uses)
  - Note: risk concerns for dinotefuran based lines of evidence other than RQs

Fish and Aquatic Plants

All uses:

- Low risk

Aquatic Invertebrate Risks

- Acute/chronic LOC exceedances
  - Chronic EECs > Freshwater LOAEC for most uses
- Open lit toxicity data supports potential for mortality/sublethal effects
- 1 crayfish incident (Imi)
- Frequency of exceedance analysis (thia/clothi)
- Monitoring data (largely non-targeted) exceed chronic endpoints and approach acute endpoints
  - When detected, monitored concentrations aligned with modelled EECs

- Imi had wider range of aquatic invertebrate species tested
  - Data from University of Guelph (CAN) expected by time of final assessment (Spring 2018)
  - Dino had quantitative data only for crustaceans, several orders of magnitude less sensitive

43

Mayfly sensitivity, quant data exist only for IMI

1-in-10 vs entire time series (30 yr); persistent yet don't see buildup based on PWC modelling
Exceedances every year for foliar uses (thia); less frequently for seed treatment uses



FW Inverts:
Foliar, Soil & Seed: Chronic risk indicated for clothi, Imi, thia;
Dino: no chronic risk (Daphnia)

SW Inverts:
Foliar, Soil & Seed: chronic risk indicated for clothi & imi;
Dino: chronic risk for foliar (rice, watercress)



FW Inverts (Non-listed):
Foliar: acute risk for clothi (rice only), imi (all uses)
Soil: acute risk for clothi (fruit/nut trees, poultry only), imi (many uses)
Seed: acute risk for clothi (rice only), imi, thia (rice only)

SW Inverts (Non-listed):
Foliar & Seed: acute risk for clothi (rice only)

Aquatic Monitoring Data

- Sourced primarily from Water Quality Portal (multiple databases within)
  - Generally non-targeted in nature
  - Some targeted open literature data available for imidacloprid
- For imi, clothi, and thia:
  - Some – most detections = chronic risk
  - Few – many detections – acute risk

Chemical	# Samples	% Detection Frequency	Highest concentration (µg a.i/L)
Imidacloprid			
Clothianidin	2,051	14%	1.34
Thiamethoxam	3,005	9%	4.37
Dinotefuran	1316	30%	11.7

Bird & Mammal Risk Conclusions

53

Moving into the risk characterization…

Terrestrial Big Picture Risk Conclusions

Birds and mammals

- *Foliar and soil uses:*
  - Acute and chronic risk concerns for imidacloprid and clothianidin

- *Seed treatments:*
  - Acute and chronic risk concerns for imi, clothi and Thia
  - Dino not currently registered for seed treatments

Terrestrial Plants

All uses:

- Low risk

Birds and Mammals (non-listed only)

Birds:

- **Acute Risk:**
  - ***Foliar/Soil:*** Acute (dose-based) risk indicated for **Clothi/Imi** (certain uses only)**;** low acute risk on subacute, dietary basis (all uses)
  - ***Seed:*** Acute (dose-based) risk for **all 3 chemicals**; area-based risk for **Clothi/Imi**

- **Chronic Risk:**
  - ***Foliar/Soil:*** No chronic risk indicated (**all 4 chemicals**)
  - ***Seed***: Chronic (diet) risk indicated for **all 3 chemicals**;

Mammals:

- **Acute Risk:**
  - ***Foliar/Soil:*** Acute risk indicated for **Clothi**
  - ***Seed:*** Dose-based risk for **all 3 chemicals**; area-based risk for **Clothi/Imi**

- **Chronic Risk:**
  - ***Foliar/Soil:*** Chronic risk indicated for **Clothi/Imi**
  - ***Seed:*** Chronic risk indicated for **all 3 chemicals**

Two things to note:

Acute risk conclusions for non-listed species only, and as discussed previously, dino has no registered seed treatments

Differences in acute dose-based risk driven by more sensitive imi and clothi endpoints

Big picture, marginal exceedances for folair/soil, seed treatment driving risk concerns



Application method breakdown, increase figure size

Do not highlight listed results, remove dotted lines

Big picture, marginal exceedances for foliar/soil, seed treatment driving risk concerns



Same story as acute

Refinements for Seed Treatments Acute Risk

- RQs assume seeds = 100% of avian/mammalian diet and palatable
- Refinements: (1) seed size (passerines only), (2) % of diet = seeds @ LOC

(1) **Seed Size: Corn, soybean, potato**: seed considered too large for consumption by small & medium passerine birds; **Cotton** too large for small passerines

(2) **% of Daily Diet = Seeds @ LOC:**
- **Large birds:** ≥ 99% for **cotton**, **soybean**, **corn** (clothi), **potato** (imi);
- **Large mammals:** >96% for **soybean** (clothi & Imi); **cotton, sorghum, wheat, potato** (imi)
- Small birds and mammals: small veg seed: few seeds and low % of diet (10%)

Group use patterns in "higher risk" category

1. Seed size driving some conclusions:
Lettuce, sugarbeet, (only few needed, possible to be ingested)

2. Few seeds needed, but seed size too big (small/ed passerines)
Corn, soybean, cotton (small only)

3. Use Patterns and size class of lower concern
Larger percentage of diet, more seeds to consume

Tie back to SLUA, larger percentage of diet to reach LOC for major uses (corn, soybean, cotton)

Highlight relative ease of mitigating on small vegetable seeds (e.g. lettuce) by recommending bittering agent on seed coating

Uncertainties:
Terrestrial Risk: Assumes seed is palatable available for consumption
Aquatic Risk: > 2 cm assume no runoff
Actual seeding depth likely to be variable

Conclusions & Next Steps

Moving into the risk characterization…

Other Regulatory Assessments

- PMRA has released final pollinator assessment for IMI
  - Targets specific application methods and timing of certain use patterns
  - CLOTHI and THIA are yet to be published but preliminary indication is that many of the conclusions are in line with EPA's

- EFSA has released seed treatment assessments
  - IMI and CLOTHI are low risk from pollen and nectar residues from oral route
  - THIA inconclusive from oral route
  - Dust-off is high risk for IMI, CLOTHI and THIA

Potential Weaknesses of RA Approach

Deliberative Process / Ex. 5

Strengths of RA Approach

Deliberative Process / Ex. 5

Next Steps and Ongoing Work

- Briefings for OD and beyond?
- Support PRD with risk management / mitigation options development
- Finalize bee risk assessments
- Complete work on RA supporting attachments (residue bridging strategy, etc.)
- Complete response to comments, addendums, and data reviews

63

While the draft documents are with PRD, the teams are busy updating attachments, going through comments, and adding characterization of the strength of the evidence for risk conclusions.

A future step is to synthesize the "lessons learned" from the bridging strategy and the pollen weighting factor and consider implications for risk assessments for other chemicals. this will also include revisiting BeeREX.

Back-pocket Slides

Use and Usage Information

- Registered on wide variety of agricultural and non-agricultural use patterns
- Applied as seed treatment, soil, or foliar, or as combination of methods
- Max annual rates up to **0.4** (clothi), **0.27** (thia), **0.54** (dino), and **0.5** (imi) lbs a.i/A

Chemical	Estimated annual usage (lbs/year)	Major uses (lbs/year)
Clothianidin	1,500,000	**Corn** (seed treatment; 1,400,000)
Imidacloprid	1,120,000	**Soybean** (seed treatment, 430,000) **Cotton, Potato, Wheat** (all app. methods, 100,000 ea.)
Thiamethoxam	919,000	**Corn** (seed treatment; 300,000) **Cotton** (foliar, soil, seed; 160,000) **Soybean** (seed treatment; 300,000)
Dinotefuran	22,500	**Cantaloupes** (5,000) **Rice** (4,000)

Data Bridging Needs vs. Available Data--Foliar

Crop Group	Chemical (Foliar Application)			
	Imidacloprid	Clothianidin	Thiamethoxam	Dinotefuran
Root/Tuber Vegetables		Potato		
Legumes	Soybean		Soybean	
Cucurbits	Watermelon	Pumpkin	Pumpkin, Cucumber	Pumpkin, Cucumber
Citrus Fruits	Orange		Orange	
Pome Fruits	Apple	Apple	Apple	
Stone Fruits	Cherry, Peach, Plum, Apricot	Peach	Cherry, Peach, Plum	Cherry, Peach
Berries/Small Fruits		Grape	Strawberry, Blueberry, Cranberry	Blueberry, Cranberry
Cereal Grains			**	***
Tree nuts	*	Almonds		
Oilseed	Cotton	Cotton	Cotton	Cotton
Fruiting Vegetables	Tomato		Tomato	Tomato

* Except almond for IMI; ** registered for barley only (not bee attractive); *** registered for rice only (not bee attractive)

Data Bridging Needs vs. Available Data--Soil

Crop Group	Chemical (Soil Application)			
	Imidacloprid	Clothianidin	Thiamethoxam	Dinotefuran
Root/Tuber Vegetables		Potato		Potato
Legumes				
Cucurbits	Melon, Watermelon	Melon, Pumpkin, Cucumber, Squash	Melon, Pumpkin, Cucumber, Squash	Melon, Pumpkin, Cucumber, Squash
Citrus Fruits	Orange, Mandarin, Grapefruit	Orange, Lemon	Orange	
Pome Fruits	Apple			
Stone Fruits	Cherry, Peach, Plum, Apricot			
Berries/Small Fruits	Strawberry, Blueberry	Grapes	Strawberry	
Cereal Grains		Corn**		
Tree nuts	*			
Oilseed	Cotton			
Fruiting Vegetable	Tomato		Pepper, Tomato	Pepper

* Except almond for IMI; ** Experimental Use permit for in-furrow soil application for corn.

Data Bridging Needs vs. Available Data—Seed and Trunk Injection

Crop Group	Application Method	Chemical			
		Imidacloprid	Clothianidin	Thiamethoxam	Dinotefuran
Root/Tuber Vegetables	**Seed**				
Legumes	**Seed**	Soybean	Soybean	Soybean	
Cucurbits	**Seed**	Melon*	Melon*		
Cereal Grains	**Seed**	Corn	Corn	Corn	
Forage, fodder, straw, hay (alfalfa)	**Seed**				
Peanut	**Seed**				
Oilseed	**Seed**	Sunflower, Canola	Cotton, Sunflower*, Canola	Cotton, Sunflower*, Canola	
Stone Fruit	**Trunk Injection**				Cherry

*only studies available are for European data

Seed Treatment Bridging Results

- Influence of site
  - For the same crop, residues are similar from site to site
- Influence of year
  - For some sites, residues are different from year to year
    - confidence intervals do not overlap
- Influence of chemical
  - Chemical does not appear to impact residue levels
  - Data can be bridged across chemicals
- Consideration of other plant tissues
  - Pollen and anthers – similar for corn
  - Pollen/nectar and flowers – same order of magnitude, but higher for soybean, cotton, canola
  - Pollen/nectar and leaves – order of magnitude higher for corn, soybean, cotton
- Influence of crop on concentrations
  - Residues in canola were higher than other crops

Foliar and Soil Residue Bridging

1. Estimate mean & variance of 2 parameters:
   A. **Initial (peak) Concentration:** (C_initial)
   B. **Dissipation rate:** (k)
2. Calculate dissipation curves associated with randomly sampled C_peak and k using Monte Carlo analysis (e.g., analogous to simulating 1000 fields). Select various "percentiles"
3. Use residue dissipation curve(s) for subsequent Tier 2 risk assessment

Monte Carlo simulation of residue dissipation curves



- Conceptually similar to EPA TREX approach (arthropod residues) & EFSA SHVAL tool for bee risk assessment, and HED tolerance analysis

Simulation Components

1. Select “k” mean, standard deviation (SD), Distribution type
2. Select C_initial mean, SD, distribution type
3. Set bounds around “k” and C_initial based on min and max of data
4. Based on distribution parameters of k and C_initial, randomly select values of k and C for each day of analysis (1000 times)
5. Calculate nectar equivalents based on new pollen method, # days to reach NOAEC, LOAEC
6. Plot risk curves for each matrix type along side of measured data.

Example Simulation: Berries foliar applications



Example Simulation: Berries foliar applications



Risk Calls – Cotton Foliar

Line of evidence	Imidacloprid		Clothianidin		Thiamethoxam*		Dinotefuran	
	Floral Nectar	Extrafloral Nectar	Floral Nectar	Extrafloral Nectar	Floral Nectar	Extrafloral Nectar	Floral Nectar	Extrafloral Nectar
Residue data	CS		CS		B		CS	
FREQUENCY: Number of residues exceeding NOAEC	2	0	8	28	0 (0)	4 (10)	2	11
FREQUENCY: Number of residues exceeding LOAEC	0	0	4	26	0 (0)	2 (3)	-	-
DURATION: Window of residue exceedances (days after last application)	20	NA	7	28	NA (NA)	12 (14)	9	25
# of days above NOAEC (90th %-tile estimates)	15	22	9	49	8 (11)	20 (24)	10	17
# of days above LOAEC (90th %-tile estimates)	8	0	7	44	6 (8)	17 (21)	-	-
# of days above NOAEC (median estimates)	7	2	2	25	0 (3)	7 (10)	7	33
# of days above LOAEC (median estimates)	8	0	0	22	0 (0)	5 (8)	-	-
% of diet from treated area required to reach NOEC	90%	>100%	13%	<1%	>100% (>100%)	29% (12%)	52%	1%
Incidents	None		3 honey bee kills following foliar applications		None		None	

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

The next few slides have a similar format so I will take some time to orient you to the tables. These are slightly abridged versions of the lines of evidence tables presented in the Tier II risk characterizations. The rows are the lines of evidence, including, residues considered (whether chemical specific or bridged); the duration and frequency of exceedances based on measured data [for the number of residues exceeding: we are presenting absolute number of residues exceeding, but it is important to note that some have a lot of data and some don't]; the duration of exceedances based on predicted data; the magnitude of residue exceedances as presented by the % of diet from the treated area required to reach the NOAEC [When the % of diet needed to reach the NOAEC is >100%, more than 100% of a colony's diet would need to come from a treated field to elicit an effects. This essentially equates to unlikely.]; and the incidents. The columns are the individual chemicals. Given the importance of the THIA degradate (i.e., CLOTHI), residue data are compared to both the thia and clothi endpoints (presented in parentheses).

For cotton, pollen is not considered honeybee attractive so it is not considered in the residues (i.e., we did not calculate a total concentration in diet based on the method described previously). Somewhat unique to cotton, is that plants produce both floral nectar and extrafloral nectar. Extrafloral nectar is basically nectar produced by the plant from nectaries that occur on stems rather than flowers. While floral nectar is considered honey bee attractive, the attractiveness of extrafloral nectar is uncertain. However, based on similarities in composition, it is reasonable to assume bees are attracted to extrafloral nectar, though the extent and duration of use are uncertain. For the purposes of the risk assessment, we present risks for both floral and extrafloral nectar. Cotton does not require managed pollinators but commercial beekeepers do utilize cotton fields for their hives so there is potential for exposure. And there are confirmed incidents for IMI and CLOTHI that indicate this is a relevant route of exposure.

There was sufficient data to conduct monte carlo analyses based on chemical-specific data for IMI, CLOTHI, and DINO, while THIA relies on bridged data from the other three chemicals. In general, the frequency, duration, and magnitude of exceedances are greater for extrafloral nectar than floral nectar when considering measured and predicted concentrations. IMI is an exception to this.

Risk Calls- Cucurbits

Line of evidence	Imidacloprid	Clothianidin		Thiamethoxam*		Dinotefuran
	Soil	Foliar	Soil	Foliar	Soil	Foliar
Chemical specific residue data	**CS, B**	**CS, B**	**CS, B**	**CS, B**	**CS, B**	**CS, B**
Number of residues exceeding NOAEC	25	15	13	8 (14)	1 (9)	5
Number of residues exceeding LOAEC	8	12	7	5 (10)	0 (1)	-
Window of residue exceedances (days after last application)	65	19	57	16	34[1]	15
# of days above NOAEC (90th %-tile estimates)	NA	26	NA	11 (22)	NA	7
# of days above LOAEC (90th %-tile estimates)	NA	17	NA	6 (13)	NA	-
# of days above NOAEC (median estimates)	NA	13	NA	4 (11)	NA	0
# of days above LOAEC (median estimates)	NA	9	NA	0 (6)	NA	-
% of diet from treated area required to reach NOEC	18%	6.4%	28%	20% (9%)	>100% (87%)	30%
Incidents	None	None		None		None

*() Based on CLOTHI endpoints
[1] Single day exceedance only

76

CS = Chemical Specific Data, B=Bridged Data

Here is the lines of evidence table for cucurbits. Risks are presented for the different application methods. You'll see more "NAs" in the table, that is because the Monte Carlo analysis could not be conducted for soil applications so days above NOAEC/LOAEC could not be estimated. Note that imi is not registered for foliar applications and dino soil applications are considered "low", which is why you don't see them in the table.

The general conclusions are that residues from foliar applications are higher that those from soil applications, as evidenced by the lower percent of diet necessary to reach the NOAEC, however, the duration of the exceedances are much longer for the soil applications (months vs. weeks). This will be a common theme to the crop groups with both foliar and soil applications. I'll note that this discussion is not currently in the risk assessments, but can be inferred.

Cucurbits are pollinator attractive and do use managed pollinators. No incidents.

Risk Calls – Orchards

Line of evidence	Imidacloprid			Clothianidin	Thiamethoxam*		Dinotefuran
	Foliar: Pre-bloom (Citrus)	Foliar: Post-bloom (Stone/Pome)	Soil: Pre-/Post-bloom	Soil: Pre-/Post-bloom	Foliar: Pre-bloom	Soil: Pre-/Post-bloom (Citrus Only)	Foliar: Pre-bloom
Chemical: specific residue data	**CS, B**	**CS, B**	**CS, B**	**CS, B**	**CS, B**	**CS, B**	**B**
Number of residues exceeding NOAEC	24	2	63	23	11-20 (20-23)	5 (16)	8
Number of residues exceeding LOAEC	22	0	24	12	8-11 (14-21)	1 (8)	-
Window of residue exceedances (days after last application)	35	6+ months	6 months	6 months	35	5 months	12
# of days above NOAEC (90th %-tile estimates)	NA	NA	NA	NA	NA	NA	NA
# of days above LOAEC (90th %-tile estimates)	NA	NA	NA	NA	NA	NA	-
# of days above NOAEC (median estimates)	46	NA	NA	NA	18-25 (26-35)	NA	14
# of days above LOAEC (median estimates)	34	NA	NA	NA	12-18 (20-27)	NA	-
% of diet from treated area required to reach NOEC	0.7%	67%	5%	11%	3-5%	36%	6%
Incidents	2 bee kill incidents on orchard crops			None	13 bee kill incidents on orchard crops		None

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

77

Here is the lines of evidence table for the orchards. As discussed previously, based on the residue bridging strategy it was determined that foliar and soil applications should be separated, as well as pre-bloom and post- bloom applications. You can see these distinctions in the column headings. You'll also note that some of the calls indicate risks for particular crop groups within the broader "orchards" grouping. This is because of differences in application restrictions for particular crops (i.e., no pre-bloom applications) as well as application rates.

The major take-aways from this table are 1) foliar applications result in a greater magnitude of exposure (i.e., relatively low % of diet necessary to reach the NOAEC), but residues remain above the endpoints for a relatively short duration; and 2) foil applications result in a lower magnitude of exposure but residues remain above the endpoints for an extended period of time (up to 6 months after application).

Orchards are pollinator attractive (some are highly attractive) and require managed pollinators. IMI and THIA have reported incidents of bee kills from applications to orchards crops.

Risk Calls- Berries and Small Fruits

Line of evidence	Imidacloprid			Clothianidin		Thiamethoxam*			Dinotefuran	
	Foliar: Pre-bloom	Soil: Pre-bloom	Soil: Post-bloom	Foliar: Pre-bloom	Soil: Post-bloom	Foliar: Pre-bloom	Soil: Pre-bloom	Soil: Post-bloom	Foliar: Pre-bloom	Soil: Pre-bloom
Chemical- specific residue data	**B**	**B**		**CS**		**CS, B**	**CS, B**		**CS, B**	**B**
Number of residues exceeding NOAEC	42	15		4		32 (37)	5 (11)		37	4
Number of residues exceeding LOAEC	39	11		3		23 (33)	2 (8)		-	-
Window of residue exceedances (days after last application)	22	83		37		23	83		5-22	23-60
# of days above NOAEC (90th %-tile estimates)	57	NA		NA		23 (31)	NA		40	NA
# of days above LOAEC (90th %-tile estimates)	45	NA		NA		19 (24)	NA		-	NA
# of days above NOAEC (median estimates)	28	NA		NA		14 (20)	NA		19	NA
# of days above LOAEC (median estimates)	22	NA		NA		10 (16)	NA		-	NA
% of diet from treated area required to reach NOEC	1%	24%		1		5% (2%)	31% (13%)		4%	41%
Incidents	None			None		None			None	

*() Based on CLOTHI endpoints

CS = Chemical Specific Data, B=Bridged Data

This is the lines of evidence table for the berries and small fruits crop group. Note that risk is identified for pre-bloom applications only, while all post-bloom applications are low risk. CLOTHI only registered on grape for pre-bloom applications

Berries and small fruits are pollinator attractive and some require managed pollinators.

Similar conclusions as orchards: 1) foliar applications result in a greater magnitude of exposure (i.e., relatively low % of diet necessary to reach the NOAEC), but residues remain above the endpoints for a relatively short duration; and 2) foil applications result in a lower magnitude of exposure but residues remain above the endpoints for an extended period of time (up to 6 months after application).

Post-bloom soil applications: berries and small fruits, except grape (CLOTHI and THIA)
Based on residues bridged from orchards
Residues of IMI in blueberry below colony-level endpoints