UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

MEMORANDUM

SUBJECT: EPA's Response to Comments Received on the May 31, 2016, Notice for a Pesticide Product with a New Active Ingredient 89668-U, *Wolbachia pipientis,* ZAP strain male *Aedes albopictus* (Asian tiger mosquitoes) - FIFRA (Docket ID Number: EPA-HQ-OPP-2016-0205; FRL- 9945-49)

FROM: Shannon Borges, Senior Scientist (Acting)
Microbial Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)

Milutin S. Djurickovic, Biologist
Microbial Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)

Clara Fuentes, Biologist
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)

John L. Kough, Ph.D., Senior Scientist
Microbial Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)

Jeannette Martinez, Ph.D., Senior Scientist
Microbial Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)

Wiebke Tapken, Ph.D., Regulatory Action Leader
Microbial Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)

Gail Tomimatsu, Ph.D., Plant Pathologist
Microbial Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)

Chris A. Wozniak, Ph.D., Biotechnology Special Assistant
Biopesticides and Pollution Prevention Division (7511P)

THRU: ShaRon Carlisle, Associate Chief
Microbial Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)

TO: Robert McNally, Director
Biopesticides and Pollution Prevention Division (7511P)

BACKGROUND

On July 26, 2013 the U.S. Environmental Protection Agency issued experimental use permit 89668-EUP-1 to MosquitoMate, Inc. for the new microbial pesticide *Wolbachia pipientis* ZAP strain (also referred to as *w*Pip strain) for use in the tiger mosquito *Aedes albopictus* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2013-0254"]; 78 FR 56227). The *Wolbachia* ZAP strain does not naturally occur in wild *Aedes albopictus* populations. When ZAP-infected male mosquitoes mate with ZAP-free wild females they produce non-viable offspring, leading to a reduction in the *Aedes albopictus* population upon recurrent release of ZAP-infected males.

Under permit 89668-EUP-1, MosquitoMate Inc. was able to release and monitor 100,000 male *Aedes albopictus* mosquitoes, containing a total of 38.4 mg of the pesticidal active ingredient *Wolbachia* ZAP strain. Tests were carried out in KY, CA, FL, and NY from 2013 to 2015, covering a total of 4,118 acres across these four states.

89668-EUP-1 was subsequently extended and amended in 2014 (79 FR 24427; 79 FR 49775) to allow for weekly releases of 100,000 male *Aedes albopictus* at the original test sites and six additional sites in CA. Tests were conducted on a total of 15,213 acres, using 249.6 milligrams (mg) of the microbial pesticide over a 26-week period. Under this EUP an extension was granted for testing in FL until October 31, 2015 and until September 30, 2016 for the sites in NY, KY and CA.

89668-EUP-1 allowed MosquitoMate Inc. to collect information (e.g., product performance data) necessary to support a pesticide registration application under section 3 of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

On February [redacted], 2016 EPA received application 89668-U from MosquitoMate Inc. for the registration of the microbial pesticide *Wolbachia pipientis* ZAP strain for its use in *Aedes albopictus* under section 3 of FIFRA. In accordance with [redacted], EPA published a Notice of Receipt (NOR) for the registration application in the *Federal Register* of April 28, 2016 (81 FR 25401). In response to this publication, EPA received 10 public comments that consist of a mix of negative, neutral, and positive comments from private citizens, a company (Oxitec, Ltd.), and a non-governmental organization (Center for Food Safety). EPA appreciates all of the comments received, although its primary focus below is to reiterate and respond to the negative and neutral comments.

PUBLIC COMMENTS AND EPA'S RESPONSE

Because some of the comments highlighted similar issues, EPA grouped similar comments and generated one response to each grouping. When grouping text from multiple sources, EPA provides specific details as to where the text originates. A substantial number of comments were previously addressed by EPA in the context of the issuance of an experimental use permit for a different strain of *Wolbachia* and its use in *Aedes aegypti* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; 81 FR 24605; U.S. EPA, 2015). EPA will direct to these responses whenever appropriate.

ED_006741_00009824-00002

I. PRODUCT DESCRIPTION

Comments

From EPA-HQ-OPP-2016-0205-0002 (Anonymous) - […] Second, dispursing Genetically modified mosquitos has never been proven safe or effective.

From EPA-HQ-OPP-2016-0205-0005 (Anonymous) - I'm 100% against this plan to release GMO mosquitos in FL. There's other methods that should be tried first like spraying, getting rid of standing water, and releasing mosquito eating fish.

EPA's response

The subject of this registration application is the ZAP strain (also referred to as *w*Pip) of the bacterium *Wolbachia pipientis*. This particular strain has been used to infect the tiger mosquito *Aedes albopictus*. Neither the bacterium, nor the mosquito have been genetically engineered in the process. The *Wolbachia pipientis* ZAP strain is one of at least four distinct approaches currently in development that work through the release of live mosquitoes into the environment. These can be grouped based on their effect on the progeny that result from mating with the wild mosquito population: (1) the offspring is not viable, which reduces the number of individual mosquitoes in the second generation after the initial release (population suppression), (2) the released individuals lower the ability of their progeny to transmit human diseases, while the population size remains unaffected. The first approach relies on the continuous release of mosquitoes into the environment to maintain the desired effect, while the second approach strives to establish *Wolbachia* in the wild mosquito population. Below EPA provides a short description of the approach employed by the subject of the current registration application *Wolbachia pipientis* ZAP strain. EPA directs to the response to comments to the experimental use permit 88877-EUP-2 ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]) for a more detailed discussion of the other three approaches.

(1) Subject of current registration application
Wolbachia pipientis-infected mosquitoes intended to suppress populations of *Aedes albopictus* mosquitoes (registration application 89668-U).

Purpose:	Population suppression
Mechanism:	Mosquito eggs do not hatch due to cytoplasmic incompatibility (CI)
Mosquito species:	*Aedes albopictus*
Bacterium:	*Wolbachia pipientis,* ZAP (*w*PIP) strain
Released individuals:	Males, non-biting

The *Wolbachia pipientis* ZAP-infected mosquitoes are intended to suppress the populations of *Aedes albopictus* (tiger mosquito). The *Wolbachia* bacterium is naturally occurring in the mosquito *Culex pipiens* from which it is extracted and subsequently microinjected into lab colony-reared embryos of *Aedes albopictus*. These newly infected mosquitoes are then reared in

ED_006741_00009824-00003

isolated containment facilities that follow Arthropod Containment Level-1 procedures (Tabachnick, 2006). The presence of the ZAP strain in the mosquitoes is confirmed by polymerase chain reaction (PCR). Non-biting *Aedes albopictus* males are separated from females by size differentiation and shipped to the site of application where they can mate with wild-type females that do not carry this particular strain of *Wolbachia*. Through cytoplasmic incompatibility (CI), the embryos from these matings die and the eggs do not hatch (Weeks, 2015). These mosquitoes and the bacteria within them are not genetically modified or genetically engineered. The *Wolbachia pipientis* ZAP strain in these mosquitoes is regulated as a microbial pesticide by EPA under FIFRA (U.S. EPA, 2013). EPA issued, amended, and/or extended other experimental use permits using this same technology in *Aedes aegypti* [1] (see below) and *Aedes polynesiensis* in 2012 [2].

(2) Experimental use permit issued by EPA
Wolbachia pipientis-infected mosquitoes intended to suppress populations of *Aedes aegypti* mosquitoes (88877-EUP-2; [HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]).

Purpose:	Population suppression
Mechanism:	Mosquito eggs do not hatch due to cytoplasmic incompatibility
Mosquito species:	*Aedes aegypti*
Bacterium:	*Wolbachia pipientis, w*AlbB strain
Released individuals:	Males, non-biting

(3) The Eliminate Dengue Program [3]
Wolbachia pipientis-infected mosquitoes intended to reduce the capacity of mosquitoes to harbor and transmit arboviruses (e.g., dengue and Zika viruses)3.

Purpose:	Reduce capacity of mosquitoes to harbor and transmit arboviruses
Mechanism:	Bacterial presence interferes with mosquitoes' ability to transmit diseases
Mosquito species:	*Aedes aegypti*
Bacterium:	*Wolbachia pipientis,* e.g., *w*Mel and *w*MelPop strains
Mosquitoes released:	males, non-biting and females, biting

(4) Oxitec Inc.'s Release of Insects with Dominant Lethality technology
Genetically engineered mosquitoes intended to suppress populations of *Aedes aegypti* [4].

Purpose:	Population suppression
Mechanism:	Heritable gene that produces a lethal protein, genetically engineered into mosquitoes
Mosquito species:	*Aedes aegypti*
Bacterium:	Not utilized for this method

---

[1] See docket number EPA-HQ-OPP-2015-0374 at [HYPERLINK "http://www.regulations.gov"]
[2] See docket number EPA-HQ-OPP-2012-0181 at [HYPERLINK "http://www.regulations.gov"]
[3] See [HYPERLINK "http://www.eliminatedengue.com/program"] for more information on these mosquitoes.
[4] See docket number FDA-2014-N-2235 at [HYPERLINK "http://www.regulations.gov"] or see U.S. FDA (2016) for more information on these mosquitoes.

ED_006741_00009824-00004

Mosquitoes released: Males, non-biting

II. REGULATORY OVERSIGHT

Comments

From EPA-HQ-OPP-2016-0205-0011 (Oxitec, Ltd.) - FIFRA requires that EPA can only register a pesticide if, *inter alia*, it determines that the pesticide (1) will perform its intended function without unreasonable adverse effects on the environment, and (2) when used in accordance with widespread and commonly recognized practice, will not generally cause unreasonable adverse effects on the environment. (7 U.S.C. § 136a(c)(5)) FIFRA defines "unreasonable adverse effects on the environment," in pertinent part, as "any unreasonable risk to man or the environment, taking into account the economic, social, and environmental costs and benefits of the use of any pesticide". (7 U.S.C. § 136(bb)) As is discussed in detail below, granting a full FIFRA Section 3 registration for a pesticide product consisting of *Wolbachia pipientis*, ZAP strain present at 100% in male *Aedes albopictus*, with uncontained release to the environment of unlimited numbers of modified mosquitoes carrying *Wolbachia pipientis* may entail significant undefined risk to human health and the environment. EPA is obligated, pursuant to FIFRA, to have a rational basis for any determination that the risks entailed with the intended function and use of this pesticide, if registered, will not cause unreasonable adverse effects on humans or the environment. Thus, to meet its obligations under FIFRA, prior to granting a registration for the new pesticide active ingredient *Wolbachia pipientis*, ZAP strain, which is proposed to be used at 100% in male *Aedes albopictus*, EPA must conduct a sufficiently rigorous assessment of the potential adverse impacts to human health and the environment that the uncontained release of these mosquitoes may entail. Oxitec herein discusses some of the serious and significant human health and environmental risks that must be addressed prior to the granting of a registration for *Wolbachia pipientis Aedes albopictus* mosquitoes.

At the outset, we note that Oxitec has developed a different genetic insect control technology that has been demonstrated to be efficacious in significantly reducing the population of disease-carrying mosquitoes (>90% in the Cayman Islands, Brazil, and Panama). Because Oxitec's insect control technology utilises genetic engineering of the insect genome, it has been determined that it is to be regulated by the U.S. Food and Drug Administration as a new animal drug. It is the oft-stated policy of the U.S. government that it regulates the products of biotechnology on the basis of a "risk-based, scientifically sound approach . . . that focuses on the characteristics of the biotechnology product and the environment into which it is being introduced, not the process by which the product is created." Exercise of Federal Oversight Within Scope of Statutory Authority; Planned Introductions of Biotechnology Products Into the Environment, 57 Fed. Reg. 6753 (Feb. 27, 1992) (this seminal statement is repeated throughout the policy statement, see, e.g., id. at 6754-55, 6755, 6756, 6757, and 6760). Notwithstanding this consistently stated position, the reality is that Oxitec's self-limiting mosquitoes have been subjected to a mandatory pre-market approval regulatory process at FDA that has been much more onerous than the regulatory requirements faced by the *Wolbachia*-containing mosquitoes at EPA. The distinction between the EPA review process for *Wolbachia* and that faced by Oxitec's self-limiting mosquitoes has even been noted by Nature [Waltz, 2016].

As is detailed below, horizontal gene transfer could result in *Wolbachia* effectively introducing over one thousand new genes into the recipient organism. Notwithstanding this potential for

indiscriminate gene transfer, the *Wolbachia* IIT (Incompatible Insect Technique) vector control method is subjected to a substantially less rigorous regulatory review process than is a targeted genetic engineering methodology. This is directly contrary to the intent of the Coordinated Framework and to a scientifically valid risk-based regulatory process.

Such a disparate regulatory approach to two products intent on achieving similar public health ends is inconsistent with the stated Federal regulatory policy, and, as a matter of risk-based regulatory process, is without reason and justification. Oxitec's self-limiting technology and the *Wolbachia* IIT approach both have the intended purpose of end point reductions in the population of mosquitoes and involve releases of substantial numbers of non-wild type mosquitoes to the environment. Importantly, however, Oxitec's self-limiting genetic engineering is well-defined and includes only two well-studied genes, which were purposefully added and reviewed by the regulatory agencies, whereas the potential genetic modification that may result from use of *Wolbachia* pipientis, ZAP strain bacterium is wholly undefined. Several studies have shown that horizontal gene transfer between *Wolbachia* and their insect hosts may result in gene transfers ranging from nearly the entire *Wolbachia* genome (>1 megabase) to short insertions (<500 base pairs) into various hosts [Kondo *et al*., 2002; Fenn *et al*., 2006; Dunning Hotopp *et al*., 2007; Hou *et al.,* 2014]. Effectively the *Wolbachia* IIT approach could introduce over one thousand new genes into the target mosquito with unknown consequences, and if they provide a positive selection to the mosquito in the environment this could result in novel strains pervading and spreading through the population. Therefore, as a matter of sound regulatory action, it is incongruous for the regulatory burden placed on the *Wolbachia* IIT technology to be significantly less onerous and burdensome than the requirements imposed on Oxitec's self-limiting targeted genetic engineering technology.

As an example of the disparate regulatory treatment that Oxitec's self-limiting mosquitoes have faced, FDA's Center for Veterinary Medicine (CVM) established an Animal Biotechnology Interdisciplinary Group (ABIG) to evaluate Oxitec's technology. This ABIG included experts from FDA/CVM, CDC, and EPA. It is not clear from the record if EPA intends to consult other regulatory agencies during its consideration of the *Wolbachia pipientis Aedes albopictus* Section 3 application, but Oxitec believes that the FDA review process for Oxitec's self-limiting technology raised questions and considerations that are particularly relevant to MosquitoMate application 89668-U.

Granting a full Section 3 registration for the *Wolbachia pipientis Aedes albopictus* pesticide entails significantly greater potential risk of unreasonable adverse effects than the limited *Wolbachia* EUPs that have been granted previously. The regulatory requirements for a Section 3 registration are substantially higher than are the requirements for a limited release experimental use permit. As Oxitec has detailed in these comments, there are significant and serious risk considerations that must be addressed before EPA can determine that the proponent of registration has met the FIFRA Section 3 burden of demonstrating that the *Wolbachia pipientis Aedes albopictus* will not result in unreasonable adverse effects on the environment. In its review of the *Wolbachia pipientis Aedes albopictus* Section 3 application, EPA must, at a minimum, require the proponent of registration to adequately address these serious risk considerations. If the applicant's initial submission does not address each of these issues, EPA must require

ED_006741_00009824-00007

submission of all additional data and information that is necessary to ensure that all relevant risk issues are fully addressed.

EPA's response

The *Aedes aegypti* OX513A Release of Insects with Dominant Lethality (RIDL) technology is unique in its approach to mosquito population suppression and relies on a novel antibiotic sensitive, transcriptional activator mechanism genetically engineered into *Aedes aegypti* in order to effect lethality of insects following release into the environment and the absence of tetracycline (as present under laboratory conditions). *Wolbachia* is naturally present in many arthropod species, including all orders of insects and many other arthrpods and nematodes, where it can affect the reproductive outcome of the infected individual through cytoplasmic incompatibility. The ZAP strain of *Wolbachia* is a symbiont of the mosquito *Culex pipiens* and it is not associated with wild *Aedes albopictus*. When ZAP-infected male *Aedes albopictus* mosquitoes mate with ZAP-free females, their progeny does not survive, reducing the mosquito population in the subsequent generation. This self-limiting mode of action is a) the basis for the bacterium's insecticidal activity and b) reduces the likelihood of the *Wolbachia* ZAP strain to establish in wild populations.

With regards to EPA's regulatory authority to regulate *Wolbachia*, EPA originally clarified its jurisdiction in the May 20, 2013 response to comments regarding the experimental use permit for the release of *Aedes polyniensis* mosquitoes infected with a strain of *Wolbachia pipientis* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2012-0181"]; U.S. EPA, 2013). This response was later discussed in the context of the Coordinated Framework for the Regulation of Biotechnology, and cited in the April 26, 2016 response to comments to the experimental use permit for *Wolbachia pipientis w*AlbB strain for use in *Aedes aegypti* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; U.S. EPA, 2016):

> "EPA has jurisdiction over the *Wolbachia pipientis* bacteria that are the subject of this EUP because such bacteria constitute a substance intended for preventing, destroying or mitigating a pest, and therefore meet the definition of "pesticide" under Section 2(u) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA). … In contrast, because the recombinant DNA (rDNA) construct in the genetically engineered mosquitoes developed by Oxitec using RIDL technology is being regulated by the U.S. Food and Drug Administration (FDA) as a "new animal drug" under the Federal Food, Drug, and Cosmetic Act (FFDCA), such technology does not meet the definition of "pesticide" under section 2(u) of FIFRA, which states, in part, that "… the term 'pesticide' shall not include any article that is a 'new animal drug' within the meaning of section 321(w) of Title 21….
> The type or quantity of information requested by FDA in its analysis of the OX513A product as a "new animal drug" under FFDCA is a matter for FDA, not EPA […]."

FIFRA section 3(5) (7 U.S.C. § 136a(c)(5)) specifies four conditions that have to be met prior to granting the registration of a new pesticide. Two of these conditions were highlighted by one of

ED_006741_00009824-00008

the commenters, which are "[…] (C) [the pesticide] will perform its intended function without unreasonable adverse effects on the environment; and (D) when used in accordance with widespread and commonly recognized practice it will not generally cause unreasonable adverse effects on the environment." Unreasonable adverse effects on the environment is defined in FIFRA section 2(bb) (7 U.S.C. § 136(bb)) as "(1) any unreasonable risk to man or the environment, taking into account the economic, social, and environmental costs and benefits of the use of any pesticide, or (2) a human dietary risk from residues that result from a use of a pesticide in or on any food inconsistent with the standard under section 408 of the Federal Food, Drug, and Cosmetic Act (21 U.S.C. 346a)." MosquitoMate Inc. submitted the data required under [redacted] that support a section 3 FIFRA registration for its *Wolbachia* ZAP strain and its use in *Aedes albopictus*. EPA has completed the risk assessment of the product at this time and determined that the provided information adequately addresses human health and environmental concerns pertaining to the release of the *Wolbachia* ZAP strain into the environment as will be discussed below. With the conclusion of the risk assessment, EPA is now making these data available for review by the public (see EPA's response to III.; [HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2016-0205"]).

In addition to ensuring that no unreasonable harm to humans and the environment is posed by the registration of a pesticide, FIFRA section 2(bb) (7 U.S.C. § 136(bb)) requires the Administrator in making a decision on whether to register a pesticide to "[…] weigh any risks of the pesticide against the health risks such as the diseases transmitted by the vector to be controlled by the pesticide." In the case of the *Wolbachia* bacterium, the majority of insects, and as a consequence the environment with which they interact, are naturally exposed to the bacterium and no negative effects have been documented as a result. In contrast, *Aedes albopictus* mosquitoes are a known carrier of viruses that are of concern to human health, including the West Nile, chikungunya, equine encephalitis, dengue and yellow fever viruses, and ~~possibly~~ the Zika virus. Suppressing the population of *Aedes albopictus* through the release of ZAP-infected male mosquitoes has the potential to benefit human health by decreasing the chances for disease transmission, while at the same time posing minimal risk to human health and the environment. Moreover, MosquitoMate Inc. is obligated to inform EPA of any "[…] additional factual information regarding unreasonable adverse effects on the environment of the pesticide" noted during use of the pesticide product after a registration has been granted pursuant to section 6(a)(2) of FIFRA.

EPA has addressed and serious and other risk considerations in its review of tis *Wolbachia*-based product and not found any reasonable or plausible concern with respect to its release into the environment. The *Aedes albopictus* males infected with Wolbachia pipientis are highly managed during production and quality assurance protocols are adequate to ensure its safety to man and the environment within the standards associated with pesticides under FIFRA.

III. AVAILABILITY OF PRODUCT INFORMATION

Comments

From EPA-HQ-OPP-2016-0205-0010 (Nina Fedoroff) - […] As such, I recently reviewed an application to release male mosquitoes genetically modified by the company Oxitec (now a subsidiary of Intrexon) to pass a lethal gene to offspring as a mosquito control measure. The

application had been submitted to the FDA, which has undertaken to regulate such mosquitoes as a "new animal drug." As part of the regulatory process, Oxitec was required to submit an Environmental Impact Assessment, which was available in its entirety for public comment, as was the application for experimental release of mosquitoes.

By contrast, application 89668-U from the MosquitoMate, Inc to register ZAP Males (*Aedes albopictus* males infected with *Wolbachia pipientis*) as a pesticide is not available for examination, although public comment on the application is requested. I therefore base my comments on examination of the relevant literature.
From EPA-HQ-OPP-2016-0205-0011 (Oxitec, Ltd.) - Based on the available record, it is not clear what level of environmental assessment has been conducted to this point. Because *Wolbachia* has been demonstrated to affect insects in the environment – changing their behaviour, disease transmission status, gene expression and biology [Marshall, 2007; Werren, et al., 2008; Endersby and Hoffmann, 2013].

If, after its initial review, EPA determines to grant the *Wolbachia pipientis Aedes albopictus* Section 3 registration, then, as part of its FIFRA public transparency policy, EPA must make available a full record of the data and information considered in the context of application 89668-U, and must provide a full and comprehensive explanation of the Agency's assessment of the significant concerns raised in these comments.

EPA's response

In accordance with 40 CFR §152.102, EPA published a Notice of Receipt (NOR) in the *Federal Register* of April 28, 2013 for the registration application for the new active ingredient *Wolbachia pipientis* ZAP strain in *Aedes albopictus* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; 81 FR 25401). As is typical for NORs published for section 3 applications, the information made available to the public was not extensive because EPA's review of the materials submitted by MosquitoMate Inc. was still in progress.
At this time, EPA has completed the risk assessments and drafted a proposed decision on the action. In the current Notice, and in accordance with agency policy for the application for registration of new biopesticide active ingredients, EPA is making available the full risk assessment, which includes the review of the product characterization, manufacturing process, human health data, ecological risk assessment, efficacy data, and modeling analyses for population dynamics, as well as a draft of the decision document, and provides a public comment period of [redacted] where the public will have an opportunity to comment on the full record ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]).

IV. MOSQUITOES AS DISEASE VECTORS

Comments

1. Presence of viruses

From EPA-HQ-OPP-2016-0205-0002 (Anonymous) - These mosquitos have not been tested to be disease free of Zika or other viruses and until the EPA can determine that these mosquitos are safe, I opposed this application. […]

From EPA-HQ-OPP-2016-0205-0010 (Nina Fedoroff) - […] Essential as well are data on the ability of the *Wolbachia*-infected population to acquire and transmit relevant diseases, including dengue, West Nile Virus, and particularly the Zika virus.

From EPA-HQ-OPP-2016-0205-0011 (Oxitec, Ltd.) - *Aedes albopictus* mosquitoes can naturally be superinfected with two *Wolbachia* strains (wAlbA, wAlbB), yet are still able to transmit chikungunya and dengue virus [Sinkins *et al*., 1995; Zhou *et al*., 1998].

The Sinkins et al. publication does not explore the potential for *Ae. albopictus* to transmit chikungunya (CHIKV) or the dengue (DENV) viruses. The Sinkins et al. article does discuss the potential for superinfection in *Drosophila simulans*, but the evidence for superinfection in *Aedes albopictus* is equivocal at best with complications in the interpretation of RFLPs in the digestion of the 16S RNA operon, it should be noted that *E. coli* contains several polymorphisms in 16S RNA forms within a single isolate or strain (Suardana 2014).

Formatted: Font: Italic

Aedes aegypti artificially modified with *Wolbachia* show a reduction in dengue virus replication but virus is still found in the saliva of these engineered mosquitoes which therefore have the capacity, even if reduced, to transmit disease [Ye *et al*., 2015].

Given that interactions between specific *Wolbachia* strains and mosquito species combinations are variable, and that *Wolbachia* has been shown to increase the vectorial capacity of some diseases in a given mosquito species, a full analysis of the capacity should be checked for all diseases

Research has shown *Wolbachia* enhances West Nile virus infection in the mosquito *Culex tarsalis*. This introduces the possibility that the *Wolbachia* infection could spread to *Culex* populations in areas where West Nile virus is a concern [Dodson *et al*., 2014].

Research has shown *Wolbachia* can enhance malaria parasite infection in two genera of mosquitoes [Hughes *et al*., 2012; Hughes *et al*., 2014a; Zele *et al*., 2014].

Temperature impacts *Wolbachia*-malaria interaction in mosquitoes suggesting impact of transfection might vary across diverse environments [Murdock *et al*., 2014].

Moreover, it is not unreasonable to surmise that the pathogen may evolve in response to *Wolbachia* infection in mosquitoes, with potential adverse results. Oxitec believes that additional information and analysis is required regarding interactions of host insect, the pathogen, and *Wolbachia*, to ensure that *Wolbachia* does not ultimately select for a more dangerous pathogen [Brelsfoard and Dobson, 2009].

2. Accidental female release

From EPA-HQ-OPP-2016-0205-0010 (Nina Fedoroff) - […] The use of *Wolbachia*-infected mosquitoes to control mosquito populations is in its experimental infancy. The basic observation is that a *Wolbachia* infection spreads and takes over an insect population by a variety of mechanisms, but particularly by cytoplasmic incompatibility, such that eggs produced in an incompatible cross fail to hatch. Suppression of a local insect population is based on the release of *Wolbachia* infected males only. However, the females are not genetically sterile, hence sorting is most likely done by size and visual inspection, neither of which is 100% effective.

From EPA-HQ-OPP-2016-0205-0011 (Oxitec, Ltd.) - Given that the release of females carrying the *Wolbachia* strain in the cytoplasm is to be avoided, sex sorting should approach 100%. Yet, studies have shown that mechanical sex sorting with IIT is only 90% effective in removing females. Thus, the reported inefficiency of sex sorting of *Aedes* mosquitoes infected with *Wolbachia* presents a risk that must be evaluated [O'Connor *et al*., 2012]. This is particularly true in the context of a FIFRA Section 3 registration, which may result in potentially unlimited numbers of modified mosquitoes may be released.
Research has shown the mosquito's microbiome can impede vertical transmission of *Wolbachia* [Hughes *et al*., 2014b].

The release of any females as a result of mis-sorting for any reason would allow the *Wolbachia* strain to invade the wild populations of mosquitoes in the release area, thus rendering this control approach ineffective and with the potential to spread a new strain of *Wolbachia* into the environment with unknown outcomes and consequences.

Further work is needed to define the underlying molecular mechanisms of *Wolbachia* induced reproductive modifications, particularly cytoplasmic incompatibility [Brelsfoard and Dobson, 2009].

EPA's response

1. Presence of viruses
With regard to questions on the potential for ZAP-infected *Aedes albopictus* to be vectors of diseases, it is correct that wild *Aedes albopictus* can carry and transmit certain viruses that are of concern to human health (see response to comments IV. and below). The most common approach to combat mosquito-borne diseases is to reduce the number of disease vectors, e.g. through the use of larvicides and adulticides. While the presence of the *Wolbachia* ZAP strain does not change the competency of *Aedes albopictus* to transmit viral agents, the release of male ZAP-infected mosquitoes serves a function analogous to these classes of pesticides, by reducing the mosquito population, albeit through a species-specific, non-toxic mode of action.

ZAP-infected *Aedes albopictus* are lab-reared in MosquitoMate Inc.'s facilities, which minimizes their exposure to viruses that are of concern to human health. The individuals reared in the facility are several generations removed from the original parent lines, which were collected from wild populations at a time when the Zika virus had not reached the western hemisphere, in an area where the Zika virus is currently not present. As part of the EUP requirements (89668-EUP-1) MosquitoMate, Inc. submitted an annual report to EPA in 2013 ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]) in which the company

addressed the concern that the released ZAP-infected *Aedes albopictus* individuals could themselves be carriers of human disease-causing viruses. The company tested a total of 96 mosquitoes in its rearing facility for the presence of four pathogenic arboviruses, dengue, eastern equine and St. Louis encephalitis, and West Nile. None of the 96 mosquitoes tested positive for the presence of these viruses. MosquitoMate Inc. continually screens for the presence of these arboviruses as part of their quality assurance protocols.

Several comments were made that the presence of the *Wolbachia* ZAP strain might increase the ability of *Aedes albopictus* to vector human diseases, with the implication that they could potentially pose a higher risk to human health. The risk of a mosquito to transmit any virus, irrespective of its vectorial capacity, is limited by the physical transmission of the virus from the infected mosquito to humans. In the mosquito family, this can only occur through females, as males do not bite. Only lab-reared male ZAP-infected *Aedes albopictus* will be released into the environment and, as will be discussed in 2), the expected frequency of an accidental female release is low, i.e., 1 female per 250,000 males. Given the expected reduction of the wild *Aedes albopictus* population upon release of the ZAP-infected males, the potential for the wild mosquitoes to vector human diseases is likely to decrease.

With regard to the observation that *Wolbachia* enhances West Nile virus infection in *Culex tarsalis*, it is unclear through which mechanism the ZAP strain could spread from *Aedes albopictus* to *Culex* species, as they do not interbreed. In fact, the ZAP strain is naturally occurring in a different *Culex* species, *Culex pipiens*, and the transmission to *Aedes albopictus* involves a multi-step process in which the bacterium is extracted from *Culex pipiens* and subsequently injected into *Aedes albopictus* eggs that were cleared of their endogenous *Wolbachia w*AlbA and *w*AlbA strains. The increase in WNV titer in *w*AlbB-infected *Culex tarsalis* described in Dodson *et al.*, 2014, resulted from a somatic *Wolbachia* infection, and cannot be considered the same as a stably inherited *Wolbachia* infection through vertical transmission.

In response to the comment regarding the publication by Brelsfoard and Dobson, 2009, the *Wolbachia* strain *w*Mel has been shown to be able to influence the susceptibility of the fly *Drosophila melanogaster* to RNA-type viruses (Brelsfoard and Dobson, 2009). However, there is currently no direct evidence that the same effect is elicited in mosquitoes. Furthermore, any significant changes in viral abundance in the lab-reared *Aedes albopictus* would likely be noted as part of the quality assurance protocols set in place by MosquitoMate Inc., as discussed in V. and in response to comments to the experimental use permit for *Wolbachia pipientis w*AlbB strain for use in *Aedes aegypti* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; U.S. EPA, 2016).

2. Accidental release of females

Males are separated from females through mechanical sorting by size. MosquitoMate Inc. continuously monitors for the presence of females and their accidental release as part of their quality control process. During the EUP MosquitoMate Inc. monitored the presence of female ZAP-infected *Aedes albopictus* in NY, CA, and KY during the 2013 and 2015 field seasons. Of the 1,200 ZAP-infected individuals tested, none of them were found to be female. Based on these total numbers of released individuals, EPA calculated that in 95% of release events the predicted

rate for unintended releases is 1 female per 250,000 males. Furthermore, u
nder FIFRA section (6)(a)(2) a pesticide registrant shall also submit to the Administrator "additional factual information regarding unreasonable adverse effects on the environment of the pesticide […].", which would include reports of accidental releases of ZAP-infected female *Aedes albopictus*.

As stated in the response to comments to the issuance of the experimental use permit of *w*AlbB in *Aedes aegypti* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; U.S. EPA, 2016), in the case of an accidental ZAP-infected female release, females could bite a person, but this is not expected to have negative consequences because mosquitoes that are naturally infected with *Wolbachia* bite humans and no negative effects have been reported. Additionally, no antibodies are detected in relation to *Wolbachia* when *Wolbachia*-infected mosquitoes bite humans; *Wolbachia pipientis* has no known history of infecting mammals or other vertebrates (Werren *et al*., 2008). As will be discussed in V., EPA does not expect that the low rate of accidental female releases will result in the establishment of the ZAP strain in the wild *Aedes albopictus* population.

V. ENVIRONMENTAL EFFECTS

Comment

From EPA-HQ-OPP-2016-0205-0004 (Anonymous) - […] Although there is a risk of unintended impacts to species that feed on mosquitoes through a population reduction of the food source, I believe that with appropriate and already necessary monitoring this risk will be minimal. Other mosquito species that do not transmit the Zika Virus to people will in a relatively short order fill the vacancy in the mosquito population created through this mitigation strategy, and thus any impact to the feeding species should be transient as well as minimal.

From EPA-HQ-OPP-0205-0008 (Anonymous) - The larvae are a significant food source for fish. If the fish population takes a hit, that travels up the food chain and you have top level predators starved and expanding their hunting range. […]

From EPA-HQ-OPP-2016-0205-0010 (Nina Fedoroff) - The possibility remains that the introduced strain will simply replace the native strain through amplification of the offspring of matings between *Wolbachia*-infected males and females, however infrequent. Suppression will likely be short-term and simple takeover of the population by *Wolbachia*-infected *A. albopictus* will be the long-term outcome.

Yet it appears that the EPA is about to approve unrestricted release of large numbers of infected insects without either an EIA, studies on its safety, or a back-up plan should the population be fully converted to the released *Wolbachia*-infected type or should other unexpected adverse effects surface.

Hence both an environmental impact assessment that addresses the probability of population replacement based on actual data is essential, a concern arising from the results of recent studies (Maurizio *et al.; incomplete citation not found*).

ED_006741_00009824-00014

From EPA-HQ-OPP-2016-0205-0011 (Oxitec, Ltd.) - What is the likelihood that the *Wolbachia* mosquitoes will survive and disperse once released into the environment?

What are the potential impacts of the *Wolbachia* mosquitoes in the environment, including on humans?

What are the likely consequences for the surrounding environment, should the *Wolbachia* mosquitoes survive and establish in the environment?
Any unexpected effects of *Wolbachia* could persist in the wild if any females are released, with little possibility of recall [Alphey, 2009].

In addition, for EPA to conduct an adequate scientific review of the *Wolbachia pipientis Aedes albopictus* pesticide application, EPA must address the following ecological risk assessment considerations raised in the public literature. In these comments, we highlight specific questions, risk considerations, and scientific data and information relevant to the ecological and human health risk assessment of release of *Wolbachia* mosquitoes that must be addressed prior to EPA determining that the proponent of registration of *Wolbachia pipientis Aedes albopictus* has met their burden to demonstrate that use of the pesticide will not result in unreasonable adverse effects to the environment.

Wolbachia is a bacterium residing within the cells of insects, and is passed through vertical transmission from mother to offspring. Even a single *Wolbachia* infected female could lay hundreds of eggs that would invade the wild population, thereby potentially rendering the Incompatible Insect Technique ineffective and spreading a new strain of *Wolbachia* into the environment. Modelling has shown that conditions of lower competition can favour infected females [Endersby and Hoffmann, 2013; Hancock. *et al*., 2011; Jansen *et al*., 2008]. In other words, as a mosquito population is reduced, or if a population is already low, the chances of *Wolbachia* invading the wild population are increased.

What is the likelihood that *Wolbachia* mosquitoes can reproduce and establish in the environment into which they are released?

Oxitec believes that approval of a Section 3 registration, with the subsequent likelihood of widespread release of the modified organism, poses significant ecological risks. Therefore, to meet its FIFRA obligations, EPA must conduct a comprehensive ecological risk assessment to ensure that such a registration would not entail unreasonable adverse effects.

EPA's response

The subject of the current application for registration received a substantial number of comments that are in some cases identical to ones previously addressed in response to the Notice of Receipt of the extension and amendment of *Wolbachia pipientis w*AlbB strain in the mosquito *Aedes aegypti* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; U.S. EPA, 2016). Given the substantial overlap in the environmental considerations of these two microbial pesticides, such as their mode of action and demonstrated natural abundance in the

environment, EPA's focus is on addressing new comments, those that are unique to the biology of the *Wolbachia* ZAP strain, or pertain specifically to the ecological risk assessment under section 3 of FIFRA. EPA will refer to the above-cited response to comments document whenever appropriate.

With regards to the comments that the reduction of a mosquito population might negatively affect their natural predators, it is true that certain fish and bat species eat mosquitoes. However, there is currently no evidence that *Aedes albopictus*, or any other mosquito species, constitutes a significant part of their diet. This might to some extent be due to the relatively recent arrival of *Aedes albopictus* in the continental U.S., where its presence was first reported in the mid-1980s [Moore and Mitchell, 1997] and the fact that bats and birds prefer larger prey species such as lepidopterans and coleopterans since mosquitoes offer little in the way of protein for the predator. Additionally, it must be remembered that *Aedes aegypti* and *Ae. albopictus* are invasive species which are non-native to the United States. Prior to their introduction, bats and birds established populations and thrived in the absence of these introduced species. Not unlike conventional insecticides, the effect of the *Wolbachia* ZAP strain is transient and needs to be sustained through recurrent releases of ZAP-infected male mosquitoes. The spatial effect of the ZAP-infected *Aedes albopictus* is furthermore limited by the range of the mosquito, which is about 100 m (or 109 yds). However, unlike broad-spectrum insecticides, the mode of action is species specific and thus, as one commenter pointed out, other mosquito species might take their place.

With regards to the comments on the establishment of the ZAP strain in the environment, due to bidirectional cytoplasmic incompatibility ZAP-infected *Aedes albopictus* mosquitoes will not reproduce when mating with wild populations in which the ZAP strain is not present. This effect is the basis for the bacterium's insecticidal activity. In the response to comments to the EUP of the *Wolbachia w*AlbB strain, EPA discussed results achieved by the Eliminate Dengue Program that show that a new *Wolbachia* strain can successfully establish in wild *Aedes aegypti* mosquito populations. However, this effect is dependent on the substantial, repeated releases of high numbers of both *Wolbachia*-infected males and females, and the attempts were not always successful ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]). Data submitted by MosquitoMate Inc. as part of the EUP also suggest that there is a reproductive cost associated with carrying the ZAP strain, resulting in lower reproductive success of two ZAP-infected *Aedes albopictus* compared to two ZAP-free individuals. Given the reasons immediately above, the low rate of accidental female releases i.e., 1 female per 250,000 males, and other quality control measures set in place by MosquitoMate Inc. (discussed in EPA's response to IV.), the probability for the *Wolbachia* ZAP strain to establish in wild *Aedes albopictus* populations is expected to be low. As mentioned previously, under FIFRA section (6)(a)(2) a pesticide registrant shall also submit to the Administrator "additional factual information regarding unreasonable adverse effects on the environment of the pesticide […].", which would include reports of accidental releases of ZAP-infected female *Aedes albopictus*.

As noted in the response to comments for *Wolbachia w*AlbB strain ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; U.S. EPA, 2016), the comment referring to an "environmental impact assessment" and "EIA" appears to intend to refer to either an "Environmental Assessment" (EA) or an "Environmental Impact Statement"

(EIS) under the National Environmental Policy Act (NEPA). EPA is not required to perform a NEPA analysis when issuing registrations under FIFRA. *Merrell v. Thomas*, 807 F.2d 776 (9th Cir., 1986). However, EPA does conduct an environmental risk assessment for pesticide actions under FIFRA, which the Court in *Merrell* found to be equivalent to an analysis under NEPA. In the present case, EPA conducted a comprehensive environmental risk assessment for the release of *Wolbachia* ZAP strain-infected male *Aedes albopictus* mosquitoes as it does with all pesticidal regulatory actions. The *Wolbachia* ZAP strain is a new microbial active ingredient and thus requires a risk assessment for nontarget organism impacts s and endangered species. MosquitoMate Inc. submitted data fulfilling the requirements under 40 CFR § 158.2150 for determining the potential risks for nontarget organisms. EPA conducted a comprehensive ecological risk assessment for the *Wolbachia* ZAP strain and its use in male *Aedes albopictus*.

With regards to the comment on the potential widespread release of ZAP-infected *Aedes albopictus* upon a section 3 registration, EPA did a comprehensive ecological assessment which is now available for public comment in that concludes that no adverse effects for nontarget organisms and endangered species are expected as a result of a registration of the *Wolbachia* ZAP strain in male *Aedes albopictus*. It is true that the exposure to a registered pesticide is generally expected to be higher than through the use of a pesticide under an experimental use permit. However, based on the reasons mentioned above, EPA believes that the risks posed by the release of male ZAP-infected *Aedes albopictus* is low and thus the risk to humans and the environment remains low under a section 3 registration.

As summarized in the April 26, 2016 response to comments to the experimental use permit for the *w*AlBb strain of *Wolbachia pipientis* and its use in *Aedes aegypti* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; U.S. EPA, 2016):

> "[…] EPA anticipates beneficial impact of *Wolbachia*-infected mosquitoes on nontarget vertebrate organisms, including humans, as these pesticides are intended for suppression of mosquito vectors of viral diseases that threaten public health. Adverse impacts are not expected for humans or other vertebrates because strains of *Wolbachia pipientis* are known to establish endosymbiotic relationships only with invertebrates. Because these strains of *Wolbachia pipientis* are naturally occurring, there is already a significant history of exposure to these microorganisms, and information from the open peer-reviewed scientific literature indicates that *Wolbachia* establishes endosymbioses (some mutualistic) with approximately 60% of the insect species, as well as other invertebrates."
>
> EPA anticipates minimal or transient impact on nontarget invertebrate organisms, including nontarget mosquitoes, some of which may be beneficial, owing to the complex endosymbioses and the commonality of this bacterium in invertebrate populations. Adverse impacts are expected to the majority of targeted mosquito populations of *Aedes aegypti*, if naturally occurring females of these species successfully breed with male mosquitoes that have been infected with *Wolbachia pipientis* Strain *w*AlbB."

VI. EFFECTS OF ZAP-STRAIN ON *AEDES ALBOPICTUS*

ED_006741_00009824-00017

Comments

From EPA-HQ-OPP-2016-0205-0011 (Oxitec, Ltd.) - Since *Wolbachia* is an endosymbiont living in the cytoplasm of the cell that has been shown to transfer genes into the nuclear genome of host insects, there is the potential that over one thousand bacterial genes could be transferred into the mosquito's nuclear genome through random integration events with unknown consequences.

The whole genome of *Wolbachia* can transfer to a host genome, meaning a host mosquito could be transformed with over one thousand new genes with unpredictable results [Kondo *et al.*, 2002; Fenn *et al.,* 2006*;* Dunning Hotopp *et al.*, 2007; Hou *et al.,* 2014].

Aedes albopictus is not naturally infected with Wolbachia ZAP strain, therefore this is an artificial infection in this species. However, *Aedes albopictus* mosquitoes can be naturally infected with certain *Wolbachia* strains (wAlbA, wAlbB), and even "super-infected" with both of these strains simultaneously. This is concerning as:

First, these naturally infected *Aedes albopictus* with wAlbA and wAlbB *Wolbachia* strains are still able to transmit chikungunya and dengue virus despite the presence of *Wolbachia* [Sinkins *et al.*, 1995; Zhou *et al.*, 1998].

Second, the consequences of any inadvertent release of female *Aedes albopictus* mosquitoes infected with the ZAP strain of *Wolbachia* mating with the naturally wAlbA/wAlbB infected *Aedes albopictus* is entirely unknown. What impact would this have on their offspring? Indeed, are there other natural *Wolbachia* strains in wild *Aedes albopictus* mosquitoes, and are all the possible interactions known? Has this been explored and addressed? Also, are there available data and information and analysis related to potential adverse effects on the environment?

It has already been shown that horizontal gene transfer (HGT) can transfer genes between *Wolbachia* and its host, in certain mosquito genera including *Aedes* [Klasson *et al.*, 2009; Woolfit *et al.*, 2009]. Therefore, *Wolbachia* can genetically transform its host with functional genes with currently unknown consequences.

Widespread recombination occurs throughout the *Wolbachia* genome [Baldo *et al.*, 2006], increasing the likelihood of genes changing as the *Wolbachia* evolves. In addition *Wolbachia* has been shown to change its phenotypic effects on the host insect as it evolves [Carrington *et al.*, 2010]. This could potentially change how *Wolbachia* responds to a number of factors, including how it influences host immune response and vectorial status. Therefore, potentially the vectorial capacity of *Aedes albopictus* infected with *Wolbachia* could change over time and should be continually assessed.

Importantly, however, Oxitec's self-limiting genetic engineering is well-defined and includes only two well-studied genes, which were purposefully added and reviewed by the regulatory agencies, whereas the potential genetic modification that may result from use of *Wolbachia* pipientis, ZAP strain bacterium is wholly undefined. Several studies have shown that horizontal

gene transfer between *Wolbachia* and their insect hosts may result in gene transfers ranging from nearly the entire *Wolbachia* genome (>1 megabase) to short insertions (<500 base pairs) into various hosts [Kondo *et al*. 2002; Fenn *et al.,* 2006*;* Dunning Hotopp *et al.*, 2007; Hou *et al.,* 2014]. Effectively the *Wolbachia* IIT approach could introduce over one thousand new genes into the target mosquito with unknown consequences, and if they provide a positive selection to the mosquito in the environment this could result in novel strains pervading and spreading through the population.

Introduction of *Wolbachia* into a mosquito provides the possibility to introduce over time over one thousand new genes, yet *Wolbachia* mosquitoes have not been subjected to the rigorous regulatory scrutiny that appears to be the norm for recombinant genetic modification (notwithstanding that typical genetic modification is a 'rifle shot' approach, involving very few, and fully characterised, introduced genes). As noted above, this seems entirely incongruent with the stated policy of the U.S. government as set forth in the Coordinated Framework. EPA must address the possibility of the introduction of genes of unknown function into the *Aedes albopictus* genome and the potential ecological effects that may result from such transfer.

All genes, including those introduced through genetic engineering in the case of the Oxitec OX513A product, are capable of being horizontally transmitted to other organisms, including those species which are unrelated, taxonomically speaking. The question of relevance to risk assessors is the probability of such occurrences and the potential impacts should such a transfer, integration and heritable expression occur.

Moreover, the possible persistence of *Wolbachia* mosquitoes themselves is a significant concern. For the reasons set forth below, each new strain of mosquito, or indeed any artificially *Wolbachia* infected insect needs to be treated as a new strain and thoroughly tested in the laboratory before any field releases.

Horizontal transmission between unrelated host species is a proven phenomenon in *Wolbachia* [Werren *et al*., 2008]. Studies have demonstrated that genetic sequences, ranging in size from single genes to entire bacterial genomes, have been transferred from *Wolbachia* to many of their insect hosts including *Aedes mosquitoes* [Kondo *et al*. 2002; Fenn *et al.,* 2006*;* Dunning Hotopp, *et al*., 2007; Klasson *et al*., 2009; Woolfit *et al*., 2009; Hou *et al.,* 2014], and its effect on disease transmission is variable and potentially dangerous.

There is evidence that male age and overcrowding during development (i.e., under mass rearing conditions required to produce enough males for IIT to be effective) can reduce the cytoplasmic incompatibility effect in certain insects, rendering the males fertile [Yamada *et al*., 2007] and able to spread the *Wolbachia* infection through surviving females. Thus, it is imperative that this physiological effect be investigated for *Aedes albopictus* to determine if it is a possibility in this species. If so, EPA has to impose terms and conditions on the registration to address the potential adverse impact of this phenomenon. Specifically, as part of its Section 3 application review, EPA must require the applicant to submit data and information on age effects and rearing conditions on the cytoplasmic incompatibility penetrance in *Aedes albopictus*.

ED_006741_00009824-00019

Further work is needed to define the underlying molecular mechanisms of *Wolbachia* induced reproductive modifications, particularly cytoplasmic incompatibility [Brelsfoard and Dobson, 2009].

EPA's response

Many of the issues raised above were addressed in response to the Notice of Receipt of the extension and amendment of *Wolbachia pipientis w*AlbB strain in the mosquito *Aedes aegypti* ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]; U.S. EPA, 2016). EPA's focus will be on new comments and comments that address specific issues regarding the *Wolbachia* ZAP strain and its effect on *Aedes albopictus*. EPA will refer to the previous responses whenever appropriate.

Several comments were made concerning the potential for the integration of *Wolbachia* genes into the *Aedes albopictus* genome, which might subsequently genetically alter the existing wild *Aedes albopictus* populations. Gene and genome integrations occur on a multi-generational or evolutionary timescale and the retention of new genes is determined by selection pressure, which eliminates those genes that do not confer a selective advantage to their new host. Thus, a prerequisite for the persistence of new genetic combinations arising from the release of ZAP-infected *Aedes albopictus* is their ability to pass on their genes to the next generation. As male ZAP-infected mosquitoes are dead-end hosts, and for other reasons discussed in II., IV., and V., this is an unlikely scenario.

It is correct to suggest that potential gene integrations through horizontal or lateral gene transfer from one organism to another unrelated organism(s) can occur over evolutionary time, as is the case with all genes in all organisms. While greater than 1 million species, conservatively speaking, are infected with *Wolbachia pipientis*, common evidence for transfer, integration and expression of bacterial genes in eukaryotic organisms is a rarity, although documented in the literature in specific cases. Some of these 'examples' are implied in that other competing mechanisms for significant homology amongst genes in a complex genome are pertinent, such as derivation of pseudogenes and convergent evolution.

It should be noted that lateral gene transfer from a prokaryote to a eukaryote is significantly more complex and less likely phenomenon than prokaryotic to prokaryotic transfer and integration. This is partly due to the presence of both outer cell membranes and nuclear membranes, which often preclude large, negatively charges molecules such as DNA from transiting freely. Additionally, prokaryotic genes are usually present absent introns and have features (e.g., promoters, ribosome binding sites, etc) which are specific to prokaryotic expression systems, but typically fail to function in a eukaryotic genomic background. This, in large part, explains why the presence of prokaryotic genes in eukaryotes is a rare find. Absence of positive selection of such transferred prokaryotic genes is another strong force precluding heritable transmission of such genes in a eukaryotic genome over time.

Male *Aedes albopictus* mosquitoes intended for release are evaluated for the presence of arboviruses, including DENV and ZIKV, such that they do not carry these infectious agents nor the ability to transmit them to humans or animals. Male mosquitoes do not consume human or

animal blood and therefore do not have a plausible probability of obtaining such viruses for subsequent transmission.

While *Wolbachia* can cause male killing, feminization of males, parthenogenesis, and cytoplasmic incompatibility, only the cytoplasmic incompatibility phenomenon has been associated with the ZAP strain-infected *Aedes albopictus*. As for the effects on cytoplasmic incompatibility, Calvitti et al. (2015) showed that *w*AlbA *Aedes albopictus* in dense rearing conditions did not decrease the cytoplasmic incompatibility (CI) effect. Islam and Dobson (2006) also showed that rearing *Aedes albopictus* with *Wolbachia* under crowded, low food conditions did not impact the CI effect. Yamada et al. (2007) cited by Oxitec, Ltd. refers to the effect in *Drosophila*, not mosquitoes. [NEW: address age effects and rearing conditions on CI]

Maintenance and propagation of *Aedes albopictus*, as used in production of *Wolbachia*-infected males for environmental release by MosquitoMate, does not allow for low food or overcrowded conditions as part of the manufacturing process approved by EPA.

VII. EXTENSION OF COMMENT PERIOD

Comment

From EPA-HQ-OPP-2016-0205-0007 (Center for Food Safety) - On behalf of the Center for Food Safety (CFS) and our 750,000 members, I am writing to request a 30-day extension of the comment period to provide input on MosquitoMate, Inc.'s application to register *Wolbachia pipientis*, ZAP Strain for use in non-biting, male *Aedes albopictus* (Asian tiger mosquito) to be released to mate with indigenous/wild female Asian tiger mosquitoes in order to control this specific species of mosquito through population suppression by prevention of egg hatch.
EPA has received an application to register a pesticide product containing an active ingredient not included in any currently registered pesticide products. Pursuant to the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), EPA is required to provide notice of receipt and an opportunity to comment on this application. *Wolbachia pipientis* (ZAP Strain) will be used in male *Aedes albopictus* mosquitoes and released to mate with indigenous female mosquitoes, making this a unique pesticide.
CFS does not believe that it has sufficient time to meaningfully review, analyze, and provide comments on a potential action of this magnitude. An extension to this unique field trial encompasses an array of complicated issues and requires careful review by the public, scientists, and other experts.
Therefore, CFS urges EPA to extend the comment period by at least 30 days.

EPA's response

On May 1, 2013 EPA published in the *Federal Register* the initial notice of receipt (NOR) with a request for comment for application 89668-EUP-R. In the application, MosquitoMate Inc. requested an experimental use permit for *Wolbachia pipientis* ZAP strain in male *Aedes albopictus* (78 FR 25436). One comment was received, expressing support for the issuance of the EUP. In accordance with 40 CFR § 152.102, EPA published the NOR for the registration application for the new active ingredient *Wolbachia* ZAP strain in *Aedes albopictus* in the April 28, 2016 *Federal Register*. As required by section 3(c)(4) of FIFRA, a 30-day public participation period was provided.
Subsequent to the issuance of a NOR, EPA's "Public Participation Process for Registration Actions" policy (*see* [HYPERLINK "https://www.epa.gov/pesticide-registration/public-participation-process-registration-actions"]) allows EPA to be more transparent for certain registration actions (e.g., new active ingredients and first food uses), under FIFRA section 3 by providing the public a chance to comment at a time when comprehensive information and analyses are available. With the conclusion of the risk assessment of 89668-U, this information is provided at this time, and a concurrent public comment period of [redacted] is provided ([HYPERLINK "https://www.regulations.gov/docket?D=EPA-HQ-OPP-2015-0374"]).

For the reasons discussed immediately above, as well as in consideration of encouraging the development of pesticides that may be helpful in combating mosquitoes that vector diseases (e.g., Zika, dengue, and chikungunya viruses) detrimental to human health, EPA decided not to extend the comment period on the NOR for the present EUP action or to open another comment period as is done for certain registration actions under the Public Participation Process for Registration Actions policy.

REFERENCES

Alphey, L., Natural and engineered mosquito immunity. Journal of Biology, 2009. 8: p. 40.

Baldo L, Bordenstein S, Wernegreen JJ, Werren JH. 2006. Widespread recombination throughout *Wolbachia* genomes. Molecular Biology and Evolution. 23:437-449. Available from [HYPERLINK "http://mbe.oxfordjournals.org/content/23/2/437.long"].

Baldo L, Ayoub NA, Hayashi CY, Russell JA, Stahlhut JK, Werren JH. 2008. Insight into the routes of *Wolbachia* invasion: High levels of horizontal transfer in the spider genus *Agelenopsis* revealed by *Wolbachia* strain and mitochondrial DNA diversity. Molecular Ecology 17:557-569. Available from [HYPERLINK "http://onlinelibrary.wiley.com/wol1/doi/10.1111/j.1365-294X.2007.03608.x/abstract"].

Brelsfoard CL, Dobson SL. 2009. *Wolbachia*-based strategies to control insect pests and disease vectors. Asia-Pacific Journal of Molecular Biology and Biotechnology 17:55-63. Available from [HYPERLINK "http://www.msmbb.org.my/apjmbb/html173/173a.pdf"].

Calvitti M, Marini F, Desiderio A, Puggioli A, Moretti R. 2015. *Wolbachia* density and cytoplasmic incompatibility in *Aedes albopictus*: Concerns with using artificial *Wolbachia* infection as a vector suppression tool. PLoS One 10:e0121813. Available from [HYPERLINK "http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0121813"].

Carrington LB, Hoffmann AA, Weeks AR. 2010. Monitoring long-term evolutionary changes following *Wolbachia* introduction into a novel host: The *Wolbachia* popcorn infection in *Drosophila simulans*. Proceedings of the Royal Society B 277:2059-2068. Available from [HYPERLINK "http://rspb.royalsocietypublishing.org/content/277/1690/2059.long"].

Dodson BL, Hughes GL, Paul O, Matacchiero AC, Kramer LD, Rasgon JL. 2014. *Wolbachia* enhances West Nile virus (WNV) infection in the mosquito *Culex tarsalis*. PLoS Neglected Tropical Diseases 8:e2965. Available from [HYPERLINK "http://journals.plos.org/plosntds/article?id=10.1371/journal.pntd.0002965"].

Dunning Hotopp JC, Clark ME, Oliveira DC, Foster JM, Fischer P, Muñoz Torres MC, Giebel JD, Kumar N, Ishmael N, Wang S, Ingram J, Nene RV, Shepard J, Tomkins J, Richards S, Spiro DJ, Ghedin E, Slatko BE, Tettelin H, Werren JH. 2007. Widespread lateral gene transfer from intracellular bacteria to multicellular eukaryotes. Science 317:1753-1756. Available from [HYPERLINK "http://science.sciencemag.org/content/317/5845/1753.full"].

Endersby NM, Hoffmann AA. 2013. Effect of *Wolbachia* on insecticide susceptibility in lines of *Aedes aegypti*. Bulletin of Entomological Research 103:269-277. Available from [HYPERLINK "http://www.eliminatedengue.com/library/publication/document/scientifix_paper/endersbyand_hoffmann_2012.pdf"].

Fenn K, Conlon C, Jones M, Quail MA, Holroyd NE, Parkhill J, Blaxter M. 2006. Phylogenetic relationships of the *Wolbachia* of nematodes and arthropods. PLoS Pathogens 2:e94. Available from [HYPERLINK "http://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.0020094"].

Hancock PA, Sinkins SP, Godfray HC. 2011. Population dynamic models of the spread of *Wolbachia*. The American Naturalist 177:323-333. Available from [HYPERLINK "http://www.journals.uchicago.edu/doi/10.1086/658121"].

Hou Q, He J, Yu J, Ye Y, Zhou D, Sun Y, Zhang D, Ma L, Shen B, Zhu C. 2014. A case of horizontal gene transfer from *Wolbachia* to *Aedes albopictus* C6/36 cell line. Mobile Genetic Elements 4:e28914. Available from [HYPERLINK "http://www.tandfonline.com/doi/full/10.4161/mge.28914"].

ED_006741_00009824-00023

Hughes GL, Vega-Rodriguez J, Xue P, Rasgon JL. 2012. *Wolbachia* strain wAlbB enhances infection by the rodent malaria parasite *Plasmodium berghei* in *Anopheles gambiae* mosquitoes. Applied and Environmental Microbiology 78:1491-1495. Available from [HYPERLINK "http://aem.asm.org/content/78/5/1491.long"].

Hughes GL, Rivero A, Rasgon JL. 2014a. *Wolbachia* can enhance *Plasmodium* infection in mosquitoes: Implications for malaria control? PLoS Pathogens 10:e1004182. Available from [HYPERLINK "http://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1004182"].

Hughes GL, Dodson BL, Johnson RM, Murdock CC, Tsujimoto H, Suzuki Y, Patt AA, Cui L, Nossa CW, Barry RM, Sakamoto JM, Hornett EA, Rasgon JL. 2014b. Native microbiome impedes vertical transmission of *Wolbachia* in *Anopheles* mosquitoes. Proceedings of the National Academy of Sciences of the United States of America 111:12498-12503. Available from [HYPERLINK "http://www.pnas.org/content/111/34/12498.long"].

Jansen VA, Turelli M, Godfray HC. 2008. Stochastic spread of *Wolbachia*. Proceedings of the Royal Society B 275:2769-2776. Available from [HYPERLINK "http://www.eliminatedengue.com/library/publication/document/jansen2008.pdf"].

Klasson L, Kambris Z, Cook PE, Walker T, Sinkins SP. 2009. Horizontal gene transfer between *Wolbachia* and the mosquito *Aedes aegypti*. BMC Genomics 10:33. Available from [HYPERLINK "http://bmcgenomics.biomedcentral.com/articles/10.1186/1471-2164-10-33"].

Kondo N, Nikoh H, Ijichi N, Shimada M, Fukatsu T. 2002. Genome fragment of *Wolbachia* endosymbiont transferred to X chromosome of host insect. Proceedings of the National Academy of Sciences of the United States of America 99:14280-14285. Available from [HYPERLINK "http://www.pnas.org/content/99/22/14280.long"].

Marshall JL. 2007. Rapid evolution of spermathecal duct length in the *Allonemobius socius* complex of crickets: Species, population and *Wolbachia* effects. PLoS One 2:e720. Available from [HYPERLINK "http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0000720"].

Maurizio et al. (N. Fedoroff)

Moore CG and Mitchell CJ. 1997. *Aedes albopictus* in the United States: ten-year presence and public health implications. Emerging infectious diseases 3.3: 329. Available from [HYPERLINK "https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2627635/"]

Murdock CC, Blanford S, Hughes GL, Rasgon JL, Thomas MB. 2014. Temperature alters *Plasmodium* blocking by *Wolbachia*. Scientific Reports 4:3932. Available from [HYPERLINK "http://www.nature.com/articles/srep03932"].

ED_006741_00009824-00024

O'Connor L, Plichart C, Sang AC, Brelsfoard CL, Bossin HC, Dobson SL. 2012. Open release of male mosquitoes infected with a *Wolbachia* biopesticide: Field performance and infection containment. PLoS Neglected Tropical Diseases 6:e1797. Available from [HYPERLINK "http://journals.plos.org/plosntds/article?id=10.1371/journal.pntd.0001797"].

Sinkins, S.P., H.R. Braig, and S.L. O'Neill, Wolbachia superinfections and the expression of cytoplasmic incompatibility. Proc Biol Sci, 1995. 261: p. 325-30. Available from [HYPERLINK "http://rspb.royalsocietypublishing.org/content/261/1362/325.long"]

U.S. Environmental Protection Agency (U.S. EPA). 2013. Memorandum from S. Bacchus through J. Kough, Ph.D. and K. Nesci to K.A. Matthews. Subject: Response to Comments – EPA-HQ-OPP-2012-0181/*Wolbachia pipientis* Experimental Use Permit – EPA Reg. No. 88877-EUP-1, dated May 20, 2013. Available from [HYPERLINK "https://www.regulations.gov/document?D=EPA-HQ-OPP-2012-0181-0008"].

Waltz E. 2016. US reviews plan to infect mosquitoes with bacteria to stop disease. Nature 533:450-451. Available from [HYPERLINK "http://www.nature.com/news/us-reviews-plan-to-infect-mosquitoes-with-bacteria-to-stop-disease-1.19967"].

Weeks E. 2015. *Wolbachia* Bacteria Can Control Mosquitoes with Fewer Chemicals (Article from Entomology Today Dated March 23, 2015). Available from [HYPERLINK "https://entomologytoday.org/2015/03/23/wolbachia-bacteria-can-control-mosquitoes-with-fewer-chemicals/"].

Werren JH, Baldo L, Clark ME. 2008. *Wolbachia*: Master manipulators of invertebrate biology. Nature Reviews Microbiology 6:741-751. Available from [HYPERLINK "http://www.nature.com/nrmicro/journal/v6/n10/abs/nrmicro1969.html"].

Woolfit M, Iturbe-Ormaetxe I, McGraw EA, O'Neill SL. 2009. An ancient horizontal gene transfer between mosquito and the endosymbiotic bacterium *Wolbachia pipientis*. Molecular Biology and Evolution 26:367-374. Available from [HYPERLINK "http://mbe.oxfordjournals.org/content/26/2/367.long"].

Yamada R, Floate KD, Riegler M, O'Neill SL. 2007. Male development time influences the strength of *Wolbachia*-induced cytoplasmic incompatibility expression in *Drosophila melanogaster*. Genetics 177:801-808. Available from [HYPERLINK "http://www.genetics.org/content/177/2/801.long"].

Ye YH, Carrasco AM, Frentiu FD, Chenoweth SF, Beebe NW, van den Hurk AF, Simmons CP, O'Neill SL, McGraw EA. 2015. *Wolbachia* reduces the transmission potential of dengue-infected *Aedes aegypti*. PLoS Neglected Tropical Diseases 9:e0003894. Available from [HYPERLINK "http://journals.plos.org/plosntds/article?id=10.1371/journal.pntd.0003894"].

ED_006741_00009824-00025

Zélé F, Nicot A, Berthomieu A, Weill M, Duron O, Rivero A. 2014. *Wolbachia* increases susceptibility to *Plasmodium* infection in a natural system. Proceedings of the Royal Society B. 281:20132837. Available from [HYPERLINK "http://rspb.royalsocietypublishing.org/content/281/1779/20132837.long"].

Zhou, W, Rousset F, and O'Neil S. 1998. Phylogeny and PCR-based classification of *Wolbachia* strains using *wsp* gene sequences. Proc Biol Sci 265: p. 509-15. Available from [HYPERLINK "http://rspb.royalsocietypublishing.org/content/265/1395/509.short"]

ED_006741_00009824-00026