From:	Danielle DeVoney
To:	Diana-M Wong
Cc:	Bob Benson; Krista Christensen; Charles Ris
Subject:	Reply to remaining SAB information requests
Date:	02/09/2012 05:02 PM

Diana -

Hi -

I believe we have identified the remaining information requests. This includes a few items from the 3 Feb e-mail which we could not provide immediately, additional information on our 6 Feb memo, and new request from the mtg.

Can you coordinate this with Dr. Sheppard and make sure we have captured everything? I have attached a description of what we are getting together below.

Thank You,
Danielle

Danielle DeVoney, PhD, DABT, PE
National Center for Environmental Assessment
USEPA Office of Research and Development
1200 Pennsylvania Ave., NW (8623P)
Washington, DC 20460
703.347.8558
FAX: 703.347.8692

Our memorandum (6 Feb 20112) provided additional information on the exposure-response modeling presented in chapter 5 and Appendix E of the External Review Draft of the Toxicological Review of Libby Amphibole Asbestos currently under review, in response to your request [e-mail, 3 Feb2012, 4:25 PM]. As noted in that memo, we were not able to fulfill the third request at the time. Attachment 1 of this memo provides the model parameters, standard errors and p-values for the full cohort models shown in Table E-6 of the draft document, thus fulfilling this last request.

During discussion at the SAB meeting (6-8 Feb, 2012) it became apparent that you needed additional information regarding the Tables 1 and 2 attached to our memorandum (6 Feb 2012). Table 1 provided LPT prevalence by deciles of exposure for both the full and subcohorts analyzed from the OM Scott worker cohort. As requested we have added a column for mean TSFE for each decile of exposure to this table. This new table is provided as attachment 2 below.

Table 2 in our 6 Feb memo provided the prevalence of localized pleural thickening (LPT) by quintiles of exposure, and by quintiles of time since first exposure (TSFE). We now understand that you would like to this information cross-tabulated. Attachment 3 of this memo provides this cross-tabulation of the prevalence of LPT by quintiles of exposure, further stratified by quintiles of TSFE. It is important to note that this is not the information plotted in Figure E-2 discussed during the

meeting.

The five exposure groups in Figure E-2 of the draft document are not based on quintiles of exposure, but rather a priori selected categories of exposure (See page E-16 of the Draft document.) Per discussion during the SAB panel meeting we are also providing the data plotted in Figure E-2 both by exposure and TSFE categories used in this Figure. This information is provided in Attachment 4 below.

Attachment 1: Supplemental modeling parameters for model-fitting for the full cohort analysis of the O.M. Scott workers cohort, Marysville, OH. These parameters augment the information presented in Table E-6 of the External Review Draft of the Toxicological Review of Libby Amphibole Asbestos.

Attachment 2: Prevalence of localized pleural thickening (LPT) by deciles of exposure for both the full and subcohorts analyzed from the OM Scott worker cohort.

Attachment 3: Prevalence of localized pleural thickening (LPT) by quintiles of exposure, and further stratified by quintiles of time since first exposure (TSFE).

Attachment 4: Prevalence of localized pleural thickening (LPT) for the exposure categories and TSFE categories plotted in Figure E-2 of the External Review Draft of the Toxicological Review of Libby Amphibole Asbestos.