TOTAL.....

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

See top panel for additional precautionary statements.

Does not Match current EPA stamped Label.

EPA Reg. No. 10324-159-46597
EPA Est. No. 46597-GA-1

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
IF ON SKIN OR CLOTHING: Take off contaminated clothing, Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

NET CONTENTS: 2.5 GALLONS (9.46 LITERS)

G12

LS-10P

SOLD BY: **CHEMSTAR CORP.**
INTERSTATE WEST PARKWAY, SUITE 100
LITHIA SPRINGS, GEORGIA
800-327-0777

Spout

Phy ✓
Doc___ Sample# 031517BWB0103

This label/labeling is the same (word for word) as that attached to the sampled container that has been packaged and released for shipment

BIN LABEL - Does not Match current EPA stamped label.
230 3/15/17
DATE COLLECTED 3 / 15 / 17

(Continued)
ENTRYWAY SANITIZING SYSTEMS: To prevent cross contamination from treated area to treated area, set the system to deliver 1.25 – 3.75 ounces per 2.5 gallons of water (390-1171 ppm active) of sanitizing solution. The spray/foam must cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor.

Do not mix other foam additives to the sanitizing solution.

FOAMING DISINFECTANT: At 1 ounce per gallon, this product is an effective foaming disinfectant. Using a foam generator or similar device shoot foam onto surface. Set equipment according to manufacturers' instructions to deliver proper dilution of 1:128. Apply foam onto surface. Allow foam to sit on surface for 10 minutes, then let air dry or rinse. Do not mix other foam additives to the disinfecting solution.

FOAMING SANITIZER FOR FLOOR DRAINS

Foam carries active into the drain pipes.
Concentrated liquid foamer for effective sanitizing. This product utilizes a unique foaming surfactant system which is synergistically formulated to give you long lasting foam for contact on all surfaces. May be used in all types of foaming equipment that can deliver a dilution of 1:256 product to water. It is widely accepted that it is easier to maintain drains than to unstop clogged ones. No additional foam booster required. Apply foam to all surfaces of the drain.
For Drains and Trough Drainage Systems. The foam generated from this product is an effective non-food contact sanitizer as evaluated using the Method for Efficacy of Sanitizers recommended for Inanimate Non-Food Contact Surfaces at 2 ounces per 4 gallons of water solution in the presence of 50% organic soil with a contact time of 5 minutes against the following:
Enterobacter aerogenes, Listeria monocytogenes, and Staphylococcus aureus

FLOOR DRAIN AND DRAINAGE SYSTEMS FOAMING SANITIZER

At 2 ounces per 4 gallons (390 ppm) this product is an effective drain foaming sanitizer. Thoroughly preclean drain with brush applicator prior to sanitization. Sanitization will occur only on precleaned surfaces of the drain. Using a foam generator or similar device, shoot foam into drain or drainage trough. Set equipment according to manufacturers' instructions to deliver proper dilution of 1:256. Hold foaming nozzle to drain or drainage trough. Foam for at least 10 seconds or longer depending upon length of drain pipe to sanitize. Check with foam equipment manufacturer for exact time to fill length of drain pipe requiring sanitizing. Foam will completely fill pipe and push water through the J trap or other such catch device. Allow the drain to sit for a minimum of 5 minutes before use. Drain can then be returned to service with or without rinsing. It is recommended that a small amount of water be flushed down the drain to refill the J trap. Repeat use at least weekly or as may be needed to control organism growth. Do not mix other foam additives to the sanitizing solution.

CHEMSTAR FOAM SAN

Disinfectant - Sanitizer Non-food Contact Surfaces, For Dairy, Food Handling and Process Areas, Federally Inspected Meat and Poultry Plants. Foam Entryway Sanitizer. Small Fly Ovicidal Treatment.

ACTIVE INGREDIENTS:	
Octyl Decyl Dimethyl Ammonium Chloride	3.0%
Didecyl Dimethyl Ammonium Chloride	1.5%
Dioctyl Dimethyl Ammonium Chloride	1.5%
Alkyl (C14, 50%; C12, 40%; C16, 10%) dimethyl benzyl ammonium chloride	4.0%
INERT INGREDIENTS:	90.0%
TOTAL	100.0%

Does not match current EPA stamped label!

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

See top panel for additional precautionary statements.

EPA Reg. No. 10324-159-46597
EPA Est. No. 46597-GA-1

?

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call a poison control center

Rinse skin immediately with plenty of water for 15-20 minutes.

CHEMSTAR FOAM SAN

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Apply this product with a cloth, mop, mechanical spray device or foaming device. When applied with a mechanical spray device, surfaces must be sprayed until thoroughly wetted. For disinfection, treated surfaces must remain wet for 10 minutes. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. For foaming applications, foam must be applied at a rate of 1:128. See device manufacturer for operating instructions. Allow to air dry. Do not breathe spray. Rinse with potable water after use on surfaces that come in contact with food.

NOTE: With spray applications or foaming device, cover or remove all food products. Prepare a fresh solution at least daily or when use solution becomes visibly dirty. For heavily soiled areas, a preliminary cleaning is required.

Before use in federally inspected meat and poultry food processing plants and dairies, egg processing plants, meat/poultry producing establishments, food products and packaging materials must be removed from the room or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water. Do not use this product to clean or disinfect utensils, glassware, dishes or interior surfaces of appliances.

DISINFECTION: This product is bactericidal according to the AOAC Use Dilution Test method on hard, inanimate surfaces, modified in the presence of 5% serum against: Salmonella enterica, Staphylococcus aureus
Preparation of Use Solution: Add 1 ounce per gallon of water to disinfect hard, non-porous surfaces. This product is not for use to disinfect or clean eating utensils, glassware and dishes.

FOR CONTROL OF DROSOPHILA spp. AND THE PHORIDAE FAMILY OF FLIES ON NON-FOOD CONTACT SURFACES:
For control of small flies: Drosophila spp. and the Phoridae family. To control flies on non-food contact surfaces such as floors, walls, countertops, metal surfaces, painted surfaces, glazed porcelain, glazed tile, glass, chrome, rubber, and plastic in restaurants, kitchens, dishwashing areas, and bar and wait station areas. Remove food and food packaging prior to use. Cover exposed food-handling surfaces. After removing gross filth, apply a solution of 3 ounces of this product per 5 gallons of water (or equivalent use dilution) (450 ppm active quat) to surfaces and locations where flies may breed. Spray surfaces thoroughly or apply by pouring, mopping or sponging onto the surface. Repeat application 1-2 times per week or as needed. Do not contaminate food and food packaging.

FOR CONTROL OF SMALL FLIES IN DRAINS:
For control of small flies: Drosophila spp. and the Phoridae family. Spray or pour solution of 3 ounces per 5 gallons of water (or equivalent use dilution) (450 ppm active quat) into drain during time of lowest level of drain use. Add 14 ounces of solution daily to each drain to maintain fly control. Apply around edge of the drain and coat all sides of inside of drain.

SANITIZER DIRECTIONS FOR NON-FOOD CONTACT SURFACES
This product is an effective non-food contact sanitizer as evaluated using the Method for Efficacy of Sanitizers recommended for Inanimate non-food Contact Surfaces at 200 ppm, 1 ounce per 4 gallons of water solution (or equivalent use solution) in the presence of 5% organic soil with a contact time of 1 minute against the following:

Escherichia coli O157:H7	*Salmonella enteritidis*	*Klebsiella pneumoniae*	*Shigella dysenteriae*	*Listeria monocytogenes*
Staphylococcus aureus	*Salmonella enterica*	*Yersinia enterocolitica*	*Campylobacter jejuni*	

Preparation of Sanitizer Use Solution: For floors, walls, tables, etc. At 1 ounce per 4 gallons of water use-level, this product is an effective sanitizer on hard, non-porous environmental surfaces.
Apply sanitizer use solution to precleaned, hard non-porous surfaces with a cloth, mop, sponge, sprayer or foaming device or by immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-9 inches from surface. Rub with a brush, sponge, or cloth. Do not breathe spray.
NOTE: With spray applications cover or remove all food products.
Treated surfaces must remain wet for 1 minute. Then wipe with a sponge, mop or cloth or allow to air dry.

SHOE BATH SANITIZER DIRECTIONS: To prevent cross contamination from treated area to treated area in animal areas, and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made sanitizing solution must be placed at all entrances to buildings, hatcheries, and at all the entrances to the production and packaging rooms. Scrape waterproof shoes and place in a 1/4 ounce of this product per gallon of water (or equivalent use dilution) (200 ppm active) solution for 60 seconds prior to entering area. Change the sanitizing solution in the bath at least daily or sooner if solution appears dirty.

SHOE FOAM DIRECTIONS: To prevent cross contamination from treated area to treated area in animal areas, and the packaging and storage areas of food plants, apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 1 to 1.5 ounces per gallon of water (or equivalent use dilution) (781 to 1171 ppm active) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacturer of the foam generator/aerator. Scrape waterproof shoes. Stand and/or walk through foamed area for 60 seconds prior to entering area. Foam area should be washed and replaced at least daily or when it appears dirty.

SOLD BY: **CHEMSTAR**
CORP.
120 INTERSTATE WEST PARKWAY, SUITE 100
LITHIA SPRINGS, GEORGIA
800-327-0777

LS-10P

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS
DANGER. KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Do not get into eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS
Do not mix with oxidizers, anionic soaps and detergents

ENVIRONMENTAL HAZARD

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS

DANGER. KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Do not get into eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix with oxidizers, anionic soaps and detergents

ENVIRONMENTAL HAZARD

This product is toxic to fish.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Open dumping is prohibited. Store only in original container. Do not reuse empty container. If a leaky container must be contained within another, mark the outer container to identify the contents. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse this container to hold materials other than pesticides or rinsate. Offer for recycling if available or puncture and dispose in a sanitary landfill, or by other procedures approved by state and local authorities. Follow Pesticide Disposal instructions.

RESIDUE REMOVAL INSTRUCTIONS: Triple rinse container promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water and recap. Shake for 10 seconds. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

This product was specifically formulated to provide a high foam layer of protection for foaming doorway sanitizers.

When used as directed, this product will prevent cross contamination from treated area to treated area in packaging and storage areas of food plants.

This product will lay down a foam blanket of protection to protect against cross contamination of organisms between treated area in food processing plants.

This product combines the proven efficacy of a quat with a high foaming specialty surfactant to provide you with an system specifically designed for entryway foam sanitizers.

This product may be applied through foaming apparatus and low-pressure sprayers.