EPA Reg. Jacket 69361-11 vol 2

Notebook

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Madhu Mandava
Vice President
Mandava Associates, LLC
6860 N. Dallas Parkway, Suite 200
Plano, TX 75024

Subject:	Revised label – update application rates and reapplication intervals
Product Name:	Tebucon 3.6F Fungicide
EPA Reg. No.:	69361-11
Your submission:	Amendment dated May 3, 2013
OPP Decision Number:	474152

Dear Mr. Mandava:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable.

One copy of the label stamped "Accepted" is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. Products released for shipment after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,

Hope Johnson
Acting Product Manager (21)
Fungicide Branch
Registration Division (7504P)

Enclosure

Westin, Robert

From: Madhu Mandava [Madhu@Mandava.com]
Sent: Friday, May 03, 2013 10:36 AM
To: Westin, Robert
Subject: RE: Revised tebuconazole labels: EPA Reg 69361-11 amd 69361-27
Attachments: 069361-00011-20130315c.ReparCorp..pdf; 069361.00027.20130315c.reparcorp.lblamend.pdf

Hi Bob, 5934860

I thought I had those to you and was expecting approval soon since I completed them on March 15, 2013. I am attaching the labels for further review.

Best Regards,

Madhu Mandava
Vice President
Mandava Associates, LLC
6860 N. Dallas Parkway
Suite 200
Plano, TX 75024
Tel: (972) 265-7924
Fax: (202) 223-0141
www.mandava.com

From: Westin, Robert [mailto:Westin.Robert@epa.gov]
Sent: Thursday, May 02, 2013 2:19 PM
To: Madhu Mandava
Subject: Revised tebuconazole labels: EPA Reg 69361-11 amd 69361-27

Hi Madhu:

On March 14 EPA sent you emails that specified the label changes that you had to make to make the labels acceptable. Are you intending to resubmit these labels or should I close out our files on these two amendments that you submitted January 11?

Robert Westin

USEPA/OPP/RD/FB/7504P
phone: 703-305-5721
fax: 703-305-6920
westin.robert@epa.gov

Westin, Robert

From: Westin, Robert
Sent: Thursday, March 14, 2013 2:27 PM
To: Madhu Mandava
Subject: Required label changes: Tebucon 3.6F Fungicice

EPA Reg. No.	69361-11
Your submission:	Amendment dated February 15, 2013
OPP Decision No.	474152

Dear Mr. Mandava:

The proposed label is not acceptable. Please make the changes that I specified for 69361-27 in my emails dated March 13 and 14.

Please email me a .pdf file of the revised label.

Sincerely,

Robert Westin

USEPA/OPP/RD/FB/7504P
703-305-5721
westin.robert@epa.gov

Westin, Robert

From: Westin, Robert
Sent: Thursday, March 14, 2013 2:14 PM
To: Madhu Mandava
Subject: More label change requirements: Tebuconazole 3.6F Fungicide

EPA Reg. No.	69361-27
Your submission:	Amendment dated February 15, 2013
OPP Decision No.	474153

Dear Mr. Mandava:

The following label changes are required in addition to those that I specified in my email yesterday:

1. On page 8 in the first line under the header **APPLICATION HEIGHT** change "Applications should not be made at a height greater than...." to "Do not apply at a height greater than...."

2. On page 8 in the third line under the header **WIND** change "Application should be avoided below 2 mph due to...." to "Do not apply when the wind speed is below 2 mph due to...."

Please let me know when I can expect to receive the corrected label.

Sincerely,

Robert Westin

USEPA/OPP/RD/FB/7504P
703-305-5721
westin.robert@epa.gov

Westin, Robert

From: Westin, Robert
Sent: Wednesday, March 13, 2013 3:07 PM
To: Madhu Mandava
Subject: Required label changes: Tebuconazole 3.6F Fungicide

EPA Reg. No.	69361-27
Your submission:	Amendment dated February 15, 2013
OPP Decision No.	474153

Repar Corporation
% Madhu Mandava
Vice-President
Mandava Associates, LLC
6860 N. Dallas Parkway, Suite 200
Plano, TX 75024

Dear Mr. Mandava:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is not acceptable. The following changes are required:

1. In the second line of the **STORAGE AND DISPOSAL** boxes on pages 4 and 28 change "...storage or disposal." to "...storage and disposal."

2. Reformat the **APPLICATION DIRECTIONS** for all crops in Sub Label A. Do not use the subheaders **Notes** and **General Comments**. Use the subheaders **Application Directions**, **Restrictions**, and **Resistance Management**. For instance, the instructions for Asparagus should be:

"Application Directions:
Apply Tebuconazole 3.6F Fungicide as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl. oz. of Tebuconazole 3.6F Fungicide per acre (0.11 lb ai – 0.17 lb ai per acre) in alteration with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebuconazole 3.6F Fungicide. Repeat applications on a 14-day interval as necessary to maintain control of rust.

Restrictions:
- Do not apply to harvestable spears.
- Applications may be made using ground or aerial application equipment.
- A 50 foot spray drift buffer zone is required for all aerial applications.
- Do not apply within 100 days of harvest in California and 180 days in all other states.
- Do not make more than three foliar applications per season (18 fl. oz./acre or 0.51 lb ai/acre).
- Restricted-entry interval (REI) = 12 hours.

Resistance Management:

Tebuconazole 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3) Alternating Tebuconazole 3.6F Fungicide with other DMI fungicides may lead to resistance."

3. On page 15 in the list of crops under Green onion change "Welch onion" to "Welsh onion"

4. On page 23 in the box change the subheader **USE PRECAUTION** to **USE RESTRICTION**. Draw a horizontal line above this subheader so that the **SEED LABELING** instructions are in their own box.

5. On page 23 in the header change "**OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING....**" to "**OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING....**"

6. On page 26 in the **PERSONAL PROTECTIVE EQUIPMENT** box change "If no such instructions for washables, use detergent...." to "If no such instructions for washables exist, use detergent...."

7. On page 27 in the **AGRICULTURAL USE REQUIREMENTS** box delete the sentence "The REI for each crop is listed in the application directions associated with each crop."

8. On page 29 change "**OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING....**" to "**OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING....**"

9. On page 30 change the header "**Recommended Mixing Procedure:**" to "**Mixing Procedure:**"

10. On page 31 under **TURF USE RESTRICTIONS AND PRECAUTIONS** change "Not for homeowner use." to "Not for residential use."

11. On page 31 in the next to last paragraph revise the second sentence to read "Use a volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1000 square feet.)"

12. On page 31 in the last paragraph revise the first sentence to read "Depending on the disease, water Tebuconazole 3.6F Fungicide into the crown and active root zone for best results."

13. In the bottom box on page 34 change the first sentence from "...in 10 to 30 gallons of water per 1000 sq ft." to "...in 1.5 to 3.0 gallons of water per 1000 sq. ft."

14. On page 35 change the header "**ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS**" to "**ORNAMENTAL USE RESTRICTIONS**"

15. On page 35 under **ORNAMENTAL USE RESTRICTIONS** change "Not for homeowner use" to "Intended only for use by professional applicators."

Please email me a .pdf file of the corrected label.

Sincerely,

Robert Westin

EPA/OPP/RD/FB
westin.robert@epa.gov
703-305-5721

Reply Reply All Forward Chat

RE: FW: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

Madhu Mandava [Madhu@Mandava.com]

Friday, February 15, 2013 8:17 PM

S 931299

To: Westin, Robert

Attachments: (2) Download all attachments

Tebucon 3.6F Master Label+~1.doc (918 KB) [Open in Browser]; 069361-00011-20130215b.Rep~1.pdf (357 KB) [Open in Browser]

Bob,

My apologies for sending the wrong label. I have attached a fresh version for the label amendment with the necessary revisions.

Please note that we had also submitted a Notification for Alternate Brand Name for Revolution Fungicide which was the label I had accidently e-mailed to you. Both the these actions were submitted simultaneously.

The EPA label to reference is Torque Fungicide, EPA Reg. No. 1001-87 accepted on May 15, 2012.

Best Regards,
Madhu

Madhu Mandava
Vice-President
Mandava Associates, LLC
6860 N Dallas Parkway
Suite 200
Plano, TX 75024
Tel: (972) 265-7924
Fax: (972) 265-7942
www.mandava.com

Notice: This E-mail (including attachments) is covered by the Electronic Communications Privacy Act, 18 U.S.C. §§2510-2521, is confidential and may be legally privileged. If you are not the intended recipient, you are hereby notified that any retention, dissemination, distribution, or copying of this communication is strictly prohibited. Please reply to the sender that you have received the message in error and then delete it. Thank you.

From: Westin.Robert@epamail.epa.gov [mailto:Westin.Robert@epamail.epa.gov]
Sent: Tuesday, January 29, 2013 2:17 PM
To: Madhu Mandava
Subject: RE: FW: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

re: EPA Reg No 69361-11
OPP Decision Number 474152

Madhu:

Master Label

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**
*** Alternate Brand Name: Revolution Fungicide**

Sub label A: Agricultural Uses

A. Vegetable Crops including Asparagus, Beans (fresh and dry except succulent shelled), Cucurbit Vegetable Group, Dry Bulb Onion, Garlic, Great Headed (Elephant) Garlic, Welsh Onion, Shallot, Green Onion, Leek, Spring Onion, Scallion, Japanese Bunching Onion, Green Shallots, Green Eschalots, Garden Beet, Leafy Brassica Greens Group, Okra, Turnip.
B. Field Crops including Barley, Corn (sweet corn, field corn, field corn grown for seed and popcorn), Cotton, Grasses Grown for Seed, Peanuts, Soybeans, Sunflower, Wheat and Seed Treatment (sweet corn, field corn, field corn grown for seed and popcorn).
C. Fruit and Nut Crops including Lychee, and Pecan.
D. Miscellaneous Crops: Hops

Sub label B: Turf and Ornamental Uses

A. Disease Control in Golf Course Turf
B. Disease Control in Field, Nursery and Container Ornamentals and Commercial and Residential Landscapes including Roses, Flowers, Ornamental Crabapples, Dogwoods, and Other Landscape Trees, Azales, Camellias, Rhododendrons and Other Landscape Ornamental Shrubs, Ground Covers, Vines

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

Manufactured for:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11	NET CONTENTS: 2.5 GALLONS	EPA Est. No.

Not accepted 3-14-13

Sub label A

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on various agricultural crops

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.
Cleaning the container before final disposal is the responsibility of the person disposing of this container.
Cleaning before refilling is the responsibility of the refiller.
To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.
Nonrefillable containers of 5 gallons or larger:
Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.
Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Spray Volume: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Chemigation: Apply Tebucon® 3.6F Fungicide through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.
Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add labeled amount of Tebucon® 3.6F Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Tebucon® 3.6F Fungicide should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

Compatibility Test for Mix Components:
Before mixing components, always perform a compatibility jar test. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water in a clear, clean, mixing jar. For other spray volumes, adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 2 teaspoons for each pound of dry product or 1 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions.
When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order:

1) **Water**. Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2) **Agitation**. Maintain constant agitation throughout mixing and application.
3) **Inductor**. If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA bags**. Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water-dispersible products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC), or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrate when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable).
9) **Remaining quantity of water.**

Maintain constant agitation during application.

SPRAY AND DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE Standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Additional requirements for aerial applications:
The boom length must not exceed 75% of the wingspan or rotor diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Applications greater than 10 feet above the canopy should be avoided.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications during temperature inversions.

Additional requirements for ground boom application:
Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

AERIAL DRIFT REDUCTION ADVISORY

This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply medium to large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

WIND

Do not apply when wind velocity exceeds 15 mph. Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HIMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Do not make applications during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to

remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions may be identified by temperatures that rise with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Resistance Management Statement

Tebucon® 3.6F Fungicide is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Tebucon® 3.6F Fungicide and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Repar Corporation encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Asparagus	Rust (*Puccinia* spp.)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Tebucon® 3.6F Fungicide per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).	
General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance. Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Barley	Rusts (*Puccinia* spp.) Head blight *(Fusarium* spp.*)* – Suppression	4 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Tebucon® 3.6F Fungicide. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Beans (fresh & dry except succulent shelled)	Rust (*Uromyces appendiculatus*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Beans, dry: Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (*Puccinia* spp.) Northern leaf blight (*Helminthosporium turcicum*) Southern leaf blight (*Helminthosporium maydis*) Northern leaf spot (*Helminthosporium carbonum*) Gray leaf spot (*Cercospora zeae-aydis*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Sweet corn: Tebucon® 3.6F Fungicide may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: Tebucon® 3.6F Fungicide may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) for sweet corn = 19 days. Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cotton	Southwestern cotton rust (*Puccinia cacabata*)	6 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Tebucon® 3.6F Fungicide may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew (*Sphaerotheca fuliginea* / *Podosphaera xanthii*) (*Erysiphe cichoracearum*)	4 to 6 fl. oz. per acre
	Gummy stem blight – suppression (*Didymella bryonae*) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Dry bulb onion Garlic Great-headed (elephant) garlic Shallot	White rot *(Sclerotium cepivorum)*	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii*)	4 to 6 fl. oz. per acre
	White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Tebucon® 3.6F Fungicide per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre. Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Tebucon® 3.6F Fungicide per acre per application. Repeat at an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 32.5 fl. oz. Tebucon® 3.6F Fungicide per acre per season if an in-furrow treatment is made. If Tebucon® 3.6F Fungicide is not applied as an in-furrow treatment then do not apply more than 12 fl oz. Tebucon® 3.6F Fungicide per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Grasses Grown For Seed	Rusts *(Puccinia* spp.*)*	4 to 8 fl. oz. per acre
	Apply the specified rate of Tebucon® 3.6F Fungicide as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl. oz. per acre
	Apply specified rate of Tebucon® 3.6F Fungicide when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	

General Comments: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.
For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon® 3.6F Fungicide.
A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Tebucon® 3.6F Fungicide may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots Welsh onion	White rot (*Sclerotium cepivorum*) suppression only Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii)*	4 to 6 fl. oz. per acre
	For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 24 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).
Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Hops	Powdery mildew (*Sphaerotheca humuli/ Spharerotheca macularis)*	4 to 8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (*Cercospora brassicicola*) Powdery mildew (*Erysiphe cruciferarum*) Alternaria leaf spot (*Alternaria brassicicola*)	3 to 4 fl. oz. per acre
	Notes: Do not apply more than 16 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). Do not apply more often than once every 10 days.	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restriction: Application to turnip greens is limited to East of the Rockies.
Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Garden beet roots and tops (leaves)	Cercospora leaf spot (*Cercospora beticola*)	3 to 7.2 fl. oz. per acre
	Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Lychee	Anthracnose (*Colletotrichum gloesporioides)*	4 to 6 fl. oz. per acre
	Notes: Begin first application of Tebucon® 3.6F Fungicide as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Tebucon® 3.6F Fungicide per acre per season. Tebucon® 3.6F Fungicide can be applied up to and including the day of harvest (PHI = 0 days).	
General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 2 days.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Okra	Cercospora leaf spot *(Cercospora* spp.*)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Peanut	**SOILBORNE:** Sclerotium stem and pod rot (white mold, southern blight, southern stern rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) **FOLIAR:** Early leaf spot Late Leaf spot Leaf rust Web blotch (*Phoma*) Pepper spot (*Leptoshaerulina*)	7.2 fl. oz. per acre
	FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Tebucon® 3.6F Fungicide to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Tebucon® 3.6F Fungicide in the first advisory spray in July and continue Tebucon® 3.6F Fungicide applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.	

General Comments: For optimum control of the specified soilborne diseases, four consecutive applications of Tebucon® 3.6F Fungicide must be made at 14-day intervals.

A maximum of 28.8 fluid ounces of Tebucon® 3.6F Fungicide may be applied per crop season. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.

Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Tebucon® 3.6F Fungicide as a leaf spot resistance management strategy. A spray surfactant is not necessary when Tebucon® 3.6F Fungicide is tank mixed with chlorothalonil. Mixing or alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance.

Tebucon® 3.6F Fungicide must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Tebucon® 3.6F Fungicide against the root and pod rots.

Use Tebucon® 3.6F Fungicide in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.

Restricted-entry interval (REI) = 12 hours.

Timing of Tebucon® 3.6F Fungicide Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	TEBUCON® 3.6F Fungicide Application No.	Chlorothalonil Application No.
7 Applications	3,4,5 and 6	1,2 and 7

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCON® 3.6F Fungicide
Pecan	Brown leaf spot (*Sirosporium diffusium*) Downy spot *(Mycosphaerella caryigena)* Liver spot *(Gnomonia caryae)* Scab *(Cladosporium caryigenum)* Vein spot *(Gnomonia nerviseda)* Zonate leaf spot *(Grovesinia pyramidalis)*	4 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Tebucon® 3.6F Fungicide should be applied at 4 fl. oz. per acre in a tank-mix with the labeled rate of Super-Tin® in cover sprays. Follow label directions for the use of SuperTin. Do not add a surfactant to the spray solution when tank-mixing Tebucon® 3.6F Fungicide with SuperTin. Apply Tebucon® 3.6F Fungicide in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Tebucon® 3.6F Fungicide to full-size mature trees, and 4 to 6 fl. oz. per acre of Tebucon® 3.6F Fungicide to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE(S)**	**RATE OF TEBUCON® 3.6F Fungicide**
Soybean	Rust *(Phakopsora pachyrhizi)* Powdery Mildew *(Microsphaera diffusa)*	3 to 4 fl. oz. per acre
Use Directions: Apply Tebucon® 3.6F Fungicide as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant must be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment. Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season. Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Sunflower	Rust *(Puccinia helianthi)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of Tebucon® 3.6F Fungicide per acre per season or within 50 days of harvest.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Contact your state Extension Service or Repar representative for a list of approved surfactants. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Turnip (Application is limited to East of the Rockies	Cercospora leaf spot *(Cercospora brassicicola)*	4 to 7.2 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Wheat	Rusts leaf, stem, and stripe *(Puccinia spp.)* Head blight or scab *(Fusarium spp.)* - Suppression	4 fl. oz. per acre
	Notes: Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Tebucon® 3.6F Fungicide. Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn) For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.		
SEED LABELING: To meet U.S. Federal Seed Act requirements, all seed treated with Tebucon® 3.6F Fungicide must be labeled: **"TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with the fungicide Tebuconazole. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice."** **USE PRECAUTION:** When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.		
DISEASE	**RATE Fl Oz/CWT**	**DIRECTIONS FOR USE**
Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Tebucon® 3.6F Fungicide. The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut *(Sphacelotheca reilana)*	0.27 – 0.54	

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR CORP.

Sub label B

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on ornamentals and golf courses

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by

avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Chemigation: Do not apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- Do not apply within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.

Spray Drift Management

Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground.

Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Spray Volume: For best results Tebucon® 3.6F Fungicide may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are recommended depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1'/2 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label.

Mixing: Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that the Tebucon® 3.6F Fungicide is complete dispersed in the mixture.

Recommended Mixing Procedure:

1) **Water.** Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2) Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3) **Inductor.** If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA Bags.** Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water Dispersible Products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrates. when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable)
9) **Remaining quantity of water.**

Resistance Management Information

The active ingredient in Tebucon® 3.6F Fungicide is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and exhibits no known cross-resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. Because the speed and scope of resistant population development can not be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS AND PRECAUTIONS

For use on golf course turf only.

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.

Not for homeowner use.

Not for use on turf being grown for sale or commercial use as sod.

Do not use clippings for animal feed.

Do not exceed 3.6 fl.oz. of Tebucon® 3.6F Fungicide per 1,000 sq ft per year.

Do not apply more than 6 applications per year.

* Do not apply more than 3 applications per year in New York State.

Product Information

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Tebucon® 3.6F Fungicide is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to Tebucon® 3.6F Fungicide under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85 degrees F.

Tebucon® 3.6F Fungicide can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 21 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Tebucon® 3.6F Fungicide in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Tebucon® 3.6F Fungicide should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Dollar Spot (*Sclerotinia homoeocarpa*) Copper Spot (*Gloeocercospora sorghi*) Powdery Mildew (*Erysiphe graminis*) Corticium Red Thread (*Laetisaria fuciformis*) Rusts (*Puccinia* spp.)	0.6-1.1	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days.
Brown Patch/Rhizoctonia Blight, Large Patch (*Rhizoctonia solani*) Brown Ring Patch (*R.. circinata*)	0.6-1.1	
Anthracnose -Basal and Foliar (*Colletotrichum cereale*) Red Thread (*Laetisaria fuciformis*) Pink Patch (*Limonomyces rosipellis*)	0.6-1.1	
Bermuda Grass decline (*Gaeumannomyces graminis* var. *graminis*)	0.6-1.1	Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21 day intervals.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Take All Patch (*Gaeumannomyces graminis*)	0.6-1.1	For prevention, apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (*Pyricularia grisea*)	0.6-1.1	Apply when conditions are favorable for disease development at 21 day intervals. Under conditions favoring moderate to heavy disease pressure, Tebucon® 3.6F Fungicide can be tank mixed with a registered contact fungicide at the label rate.
Stipe Smut (*Ustilago striiformis*)	0.6-1.1	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot (*Leptosphaeria korrea, L. narmari, Ophiosphaerella herpotricha, Gaeumannomyces graminis*) Necrotic Ring Spot (*Leptosphaeria korrea*)	0.6-1.1	For prevention, apply in fall when soil temperature reach 65^0 F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (*Fusarium roseum*)	0.6-1.1	Apply first application in mid-June or 21 days prior to time this blight normally becomes evident. Make applications at no less than 21 day intervals.
Summer Patch (*Magnaporthe poae*)	0.6-1.1	Apply beginning in the spring. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 21 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

Gray Snow Mold/ Typhula Blight (*Typhula incarnate*) Pink Snow Mold/Microdochium Patch (*Microdochium nivalis*)	0.6-1.1	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that Tebucon® 3.6F Fungicide be tank-mixed with other registered snow mold products for best season long results.
Fairy Ring Chlorophyllum (Lepiota), Lycoperdon, Marasmius	0.6-1.1	For prevention in Cool season turf, make two applications at the low to medium rate in the spring when root zone soil temperatures reach 55-60°F. Make a second application using a 21 day interval. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. Do not use a wetting agent unless hydrophobic soil conditions exist. For curative treatment, use the medium to high rate. For Warm season turf breaking dormancy, do not make two consecutive applications of Tebucon® 3.6F Fungicide or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. For hydrophobic areas, use an appropriate wetting agent to effectively penetrate the hydrophobic zone commonly created with this disease.

NOTE: Apply the specified amount of Tebucon® 3.6F Fungicide in 10 to 30 gallons of water per 1000 sq ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl oz of Tebucon® 3.6F Fungicide per 1000 sq ft per year. Do not exceed 6 applications per year.

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS

For use on ornamental plants only; not for woodlands or forest management.
Not for homeowner use.
Do not apply more than 10 fl oz per acre in a single application.
Do not apply more than 0.31 gallons (40 fl oz) of Tebucon® 3.6F Fungicide (equal to 1.13 lbs of tebuconazole) per acre per year.
Do not make more than 4 applications per year at highest rate.
Do not apply to bearing fruit trees or vegetables

Tebucon® 3.6F Fungicide can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Tebucon® 3.6F Fungicide is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Tebucon® 3.6F Fungicide at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern and Geraniums.

Ornamentals Disease Control

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management.	
Crabapples (Ornamental),	Anthracnose Leaf Spot		

Dogwoods and Other Landscape (Ornamental) Trees	Powdery Mildew		
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust Southern Blight	Petal Blight - Apply 2 – 3 times per week into the flowers as the open and develop color.	
Ground Covers and Vines			
For small plantings: Add 1 teaspoon to 2.5 gallons of water.			

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR.

RE: FW: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

Robert Westin to: Madhu Mandava 01/29/2013 03:17 PM
Bcc: Robert Westin

From: Robert Westin/DC/USEPA/US
To: "Madhu Mandava" <Madhu@Mandava.com>
Bcc: Robert Westin/DC/USEPA/US

re: EPA Reg No 69361-11
OPP Decision Number 474152

Madhu:

The .pdf file version you sent me has the product name Revolution throughout. You must use the primary brand name in labels submitted to the Agency. Therefore, either change the product name to "Tebucon 3.6F Fungicide" and send me the corrected .pdf file or submit a formal request to change the primary brand name to Revolution. (would also need a revised CSF with the brand name Revolution).

In addition, please delete Welsh onion from the Grasses Grown for Seed section and add Welsh Onion to the Green Onion section on page 15.

What is the precedent (reason) for changing the application rate for grass from 0.6 to 0.6-1.1 and reducing the reapplication period to 21 days? Please let me know what accepted label I should look at to check these changes.

Thanks

Robert Westin

phone: 703-305-5721
FAX: 703-305-6920
email: westin.robert@epamail.epa.gov

"Madhu Mandava" Bob, 01/29/2013 01:15:23 PM

From: "Madhu Mandava" <Madhu@Mandava.com>
To: Robert Westin/DC/USEPA/US@EPA, "A Pennington" <Andrea@reparcorp.com>
Date: 01/29/2013 01:15 PM
Subject: RE: FW: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

Bob,

My apologies. Attached is Tebucon 3.6F Fungicide label.

Regards,

Madhu Mandava
Vice-President
Mandava Associates, LLC
6860 N Dallas Parkway
Suite 200
Plano, TX 75024

RE: FW: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide
Madhu Mandava
to:
Robert Westin, A Pennington
01/29/2013 01:15 PM
Hide Details
From: "Madhu Mandava" <Madhu@Mandava.com>

To: Robert Westin/DC/USEPA/US@EPA, "A Pennington" <Andrea@reparcorp.com>

2 Attachments

image001.gif Tebucon 3.6F Master Label+ A+B (01-2013).pdf

Bob,

My apologies. Attached is Tebucon 3.6F Fungicide label.

Regards,

Madhu Mandava
Vice-President
Mandava Associates, LLC
6860 N Dallas Parkway
Suite 200
Plano, TX 75024
Tel: (972) 265-7924
Fax: (972) 265-7942
www.mandava.com

Notice: This E-mail (including attachments) is covered by the Electronic Communications Privacy Act, 18 U.S.C. §§2510-2521, is confidential and may be legally privileged. If you are not the intended recipient, you are hereby notified that any retention, dissemination, distribution, or copying of this communication is strictly prohibited. Please reply to the sender that you have received the message in error and then delete it. Thank you.

From: Westin.Robert@epamail.epa.gov [mailto:Westin.Robert@epamail.epa.gov]
Sent: Tuesday, January 29, 2013 9:49 AM
To: A Pennington
Cc: Madhu Mandava
Subject: Re: FW: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

Only the Tebuconazole 3.6F Fungicide label came through. Please email me a .pdf file of the Tebucon 3.6F Fungicide label.

Thank you.

Robert Westin

phone: 703-305-5721
FAX: 703-305-6920
email: westin.robert@epamail.epa.gov

"A Pennington" ---01/28/2013 03:26:13 PM---Hello Mr. Westin Attached are the files you requested. If you need any additional information pleas

From: "A Pennington" <Andrea@reparcorp.com>
To: Robert Westin/DC/USEPA/US@EPA
Cc: "Madhu Mandava" <Madhu@Mandava.com>
Date: 01/28/2013 03:26 PM
Subject: FW: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

---

Hello Mr. Westin
Attached are the files you requested. If you need any additional information please let me know.

Thank you

Andrea Pennington
Product Manager
Mandava Associates, LLC
8070 Georgia Avenue
Suite 209
Silver Spring, MD 20910
Tel: (301) 562 - 7330
Cell: (903) 227-5499

From: Madhu Mandava
Sent: Mon 1/28/2013 12:35 PM
To: A Pennington; Andrea Pennington
Subject: FW: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

Please see message below.

Madhu Mandava
Vice-President
Mandava Associates, LLC

6860 N Dallas Parkway
Suite 200
Plano, TX 75024
Tel: (972) 265-7924
Fax: (972) 265-7942
www.mandava.com

Notice: This E-mail (including attachments) is covered by the Electronic Communications Privacy Act, 18 U.S.C. §§2510-2521, is confidential and may be legally privileged. If you are not the intended recipient, you are hereby notified that any retention, dissemination, distribution, or copying of this communication is strictly prohibited. Please reply to the sender that you have received the message in error and then delete it. Thank you.

From: Westin.Robert@epamail.epa.gov [mailto:Westin.Robert@epamail.epa.gov]
Sent: Friday, January 25, 2013 9:13 AM
To: Madhu Mandava
Subject: Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

Subject: EPA Reg No 69361-11, OPP Decision Number 474152
EPA Reg No 69361-27, OPP Decision Number 474153

Hi Madhu:

Please email me .pdf files of the two draft labels that you submitted as amendments on January 8, 2013.

Thank you.

Robert Westin

phone: 703-305-5721
FAX: 703-305-6920
email: westin.robert@epamail.epa.gov[attachment "Tebuconazole 3.6F Fungicide Master label 01-08-2013.pdf" deleted by Robert Westin/DC/USEPA/US]

Tebucon 3.6F Fungicide; Tebuconazole 3.6F Fungicide

Robert Westin to: madhu 01/25/2013 10:12 AM

Bcc: Robert Westin

From: Robert Westin/DC/USEPA/US

To: madhu@mandava.com

Bcc: Robert Westin/DC/USEPA/US

Subject: EPA Reg No 69361-11, OPP Decision Number 474152
EPA Reg No 69361-27, OPP Decision Number 474153

Hi Madhu:

Please email me .pdf files of the two draft labels that you submitted as amendments on January 8, 2013.

Thank you.

Robert Westin

phone: 703-305-5721
FAX: 703-305-6920
email: westin.robert@epamail.epa.gov

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

January 14, 2013

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

N. BHUSHAN MADAVA
MANDAVA ASSOCIATES, LLC
REPAR CORP
1050 CONN. AVE., N.W., SUITE 1000
WASHINGTON, DC 20036

PRODUCT NAME: TEBUCON 3.6F FUNGICIDE
COMPANY NAME: REPAR CORP
OPP IDENTIFICATION NUMBER:
EPA FILE SYMBOL: 69361-11
EPA RECEIPT DATE: 01/11/13 D 474152, S 929149

SUBJECT: RECEIPT OF AMENDMENT

DEAR REGISTRANT:

The Office of Pesticide Programs has received your application for an amendment and it has passed an administrative screen for completeness.

During the initial screen we determined that the application appears to qualify for fast track review. The package will now be forwarded to the Product Manager for review to determine its acceptability for fast track status.

If you have any questions, please contact Registration Division, Risk Management Team 21, at {Phone Number}.

Sincerely,

Front End Processing Staff
Information Services Branch
Information Technology & Resources Management Division

FAST-TRACK AMENDMENTS – Completeness Screening Checklist

Expert's In-Processing Signature: Schaible Date: 1/17/13 PM #: 21

EPA Reg. Number: 69361-11	EPA Receipt Date: 1/11/13

	Checklist Item	Yes	No	N/A
1	**Application Form** (EPA Form 8570-1) - signed?	✓		
2	**Confidential Statement of Formula** (EPA Form 8570-29) - signed?			✓
3	**Certification with Respect to Citation of Data** (EPA Form 8570-34) - signed?			✓
4	**Formulator's Exemption Statement** (EPA Form 8570-27) - signed?			✓
5	**Data Matrix** (EPA Form 8570-35) [Applicable for adding me-too uses] - signed?			✓
	a) Selective Method?			
	b) Cite-All Method?			
	c) Public copy of Matrix provided? See PR Notice 98-5			
6	**Is Label included?** (5 copies)	✓		
	a) Electronic Label submitted?		✓	

Comments:

Fee for Service

{929149J~

This package includes the following

- ○ New Registration
- ⊙ Amendment

☐ Studies? ☐ Fee Waiver?

☐ volpay % Reduction: ___

for Division

- ○ AD
- ○ BPPD
- ⊙ RD

Risk Mgr. 21

Receipt No. S-	929149
EPA File Symbol/Reg. No.	69361-11
Pin-Punch Date:	1/11/2013

☒ This item is NOT subject to FFS action.

Action Code:

Requested:

Granted:

Amount Due: $ _____

Parent/Child Decisions:

☐ Inert Cleared for Intended Use ☐ Uncleared Inert in Product

Reviewer: S. Hill Date: 1/14/13

Remarks:

MANDAVA ASSOCIATES, LLC

CONSULTANTS IN SCIENCE, TECHNOLOGY AND REGULATORY AFFAIRS

6860 N. DALLAS PARKWAY, SUITE 200, PLANO, TX 75024

Telephone: (972)-265-7924 · Fax: (972) 265-7942 · E-MAIL: Madhu@Mandava.com · www.Mandava.com

Via Federal Express

January 8, 2013

Document Processing Desk (AMEND)
Ms. Mary L Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)
U.S. Environmental Protection Agency
One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

SUBJECT: Tebucon 3.6F Fungicide EPA Registration Number: 69361-11
Label Amendment to change application rate and interval period

Dear Ms. Waller:

On behalf of Repar Corp. enclosed please find the submission of Label Amendment for Tebucon 3.6F Fungicide, EPA Registration Number, 69361-11.

In support of the application for registration amendment, please find the following materials:

1. One (1) Application for Pesticide Registration (EPA) Form 8570-1)
2. One (1) Copy of original EPA approved Label
3. Five (5) copies of Draft Labeling

Please send to my attention the stamped approved labels to the address listed above.

We look forward to receiving an expedited approval of the subject registration application amendment.

Sincerely yours,

M. Mandava

Madhu Mandava
Mandava Associates, LLC
Agent for Repar Corp

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95

EPA

United States
Environmental Protection Agency
Washington, DC 20460

- [ ] Registration
- [x] Amendment
- [ ] Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 69361-11	2. EPA Product Manager Mary Waller	3. Proposed Classification [x] None [ ] Restricted
4. Company/Product (Name) Repar Corporation/TEBUCON 3.6F FUNGICIDE	PM# 21	
5. Name and Address of Applicant *(Include ZIP Code)* Repar Corporation P.O. Box 4321 Silver Spring, MD. 20914 [ ] *Check if this is a new address*	6. **Expedited Reveiw.** In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

- [x] Amendment - Explain below.
- [ ] Resubmission in response to Agency letter dated _____
- [ ] Notification - Explain below.
- [ ] Final printed labels in repsonse to Agency letter dated _____
- [ ] "Me Too" Application.
- [ ] Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

1) Application for product registration Label Amendment to update the application rates and interval periods for TEBUCON 3.6F FOLIAR FUNGICIDE (EPA Reg. No. 69361-27). NON PRIA ACTION, No fee required
2) FAST TRACK and EXPEDITED REVIEW
3) Please direct all correspondence to: Madhu Mandava, Agent for Repar Corp., Mandava Associates, LLC; 6860 N. Dallas Parkway, Suite 200, Plano, TX 75024 Telephone: 972-265-7624, Fax: (202) 223 - 0141 E-mail address: madhu@mandava.com

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging	Unit Packaging	Water Soluble Packaging	2. Type of Container
[ ] Yes [x] No	[ ] Yes [x] No	[ ] Yes [x] No	[ ] Metal [x] Plastic [ ] Glass [ ] Paper [ ] Other (Specify) _____
Certification must be submitted	If "Yes" Unit Packaging wgt. / No. per container	If "Yes" Package wgt / No. per container	

3. Location of Net Contents Information	4. Size(s) Retail Container	5. Location of Label Directions
[x] Label [ ] Container	32 oz, 1.0, 2.5, & 270 gallons	[x]

6. Manner in Which Label is Affixed to Product: [x] Lithograph [ ] Paper glued [ ] Stenciled [ ] Other _____

Section - IV

1. Contact Point *(Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)*

Name	Title	Telephone No. (Include Area Code)
Madhu Mandava	Agent for Repar Corporation	972-265-7924

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received **(Stamped)**

2. Signature	3. Title
M. Man	Agent for Repar Corporation
4. Typed Name	**5. Date**
Madhu Mandava	January 8, 2013

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete. White - EPA File Copy (original) Yellow - Applicant Copy

Master Label

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

Sub label A: Agricultural Uses

A. Vegetable Crops including Asparagus, Beans (fresh and dry except succulent shelled), Cucurbit Vegetable Group, Dry Bulb Onion, Garlic, Great Headed (Elephant) Garlic, Welsh Onion, Shallot, Green Onion, Leek, Spring Onion, Scallion, Japanese Bunching Onion, Green Shallots, Green Eschalots, Garden Beet, Leafy Brassica Greens Group, Okra, Turnip.
B. Field Crops including Barley, Corn (sweet corn, field corn, field corn grown for seed and popcorn), Cotton, Grasses Grown for Seed, Peanuts, Soybeans, Sunflower, Wheat and Seed Treatment (sweet corn, field corn, field corn grown for seed and popcorn).
C. Fruit and Nut Crops including Lychee, and Pecan.
D. Miscellaneous Crops: Hops

Sub label B: Turf and Ornamental Uses

A. Disease Control in Golf Course Turf
B. Disease Control in Field, Nursery and Container Ornamentals and Commercial and Residential Landscapes including Roses, Flowers, Ornamental Crabapples, Dogwoods, and Other Landscape Trees, Azales, Camellias, Rhododendrons and Other Landscape Ornamental Shrubs, Ground Covers, Vines

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

Manufactured for:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

Not accepted
1-29-13

EPA Reg. No. 69361-11	NET CONTENTS: 2.5 GALLONS	EPA Est. No.

Sub label A

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on various agricultural crops

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%

OTHER INGREDIENTS: 61.3%

TOTAL.....100.0%

Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use

KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:

Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential

for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30

seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Spray Volume: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Chemigation: Apply Tebucon® 3.6F Fungicide through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.
Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add labeled amount of Tebucon® 3.6F Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Tebucon® 3.6F Fungicide should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

Compatibility Test for Mix Components:
Before mixing components, always perform a compatibility jar test. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water in a clear, clean, mixing jar. For other spray volumes, adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 2 teaspoons for each pound of dry product or 1 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions.
When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order:

1) **Water**. Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2) **Agitation**. Maintain constant agitation throughout mixing and application.
3) **Inductor**. If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA bags**. Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water-dispersible products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC), or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrate when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable).
9) **Remaining quantity of water.**

Maintain constant agitation during application.

SPRAY AND DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE Standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Additional requirements for aerial applications:
The boom length must not exceed 75% of the wingspan or rotor diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Applications greater than 10 feet above the canopy should be avoided.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications during temperature inversions.

Additional requirements for ground boom application:
Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

AERIAL DRIFT REDUCTION ADVISORY

This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply medium to large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

WIND

Do not apply when wind velocity exceeds 15 mph. Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HIMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Do not make applications during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions may be identified by temperatures that rise with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Resistance Management Statement

Tebucon® 3.6F Fungicide is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Tebucon® 3.6F Fungicide and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Repar Corporation encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Asparagus	Rust (*Puccinia* spp.)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Tebucon® 3.6F Fungicide per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).	
General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance. Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Barley	Rusts (*Puccinia* spp.) Head blight (*Fusarium* spp.) – Suppression	4 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Tebucon® 3.6F Fungicide. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Beans (fresh & dry except succulent shelled)	Rust (*Uromyces appendiculatus*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Beans, dry: Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (*Puccinia* spp.) Northern leaf blight (*Helminthosporium turcicum*) Southern leaf blight (*Helminthosporium maydis*) Northern leaf spot (*Helminthosporium carbonum*) Gray leaf spot (*Cercospora zeae-aydis*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Sweet corn: Tebucon® 3.6F Fungicide may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: Tebucon® 3.6F Fungicide may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) for sweet corn = 19 days. Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cotton	Southwestern cotton rust (*Puccinia cacabata*)	6 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Tebucon® 3.6F Fungicide may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew (*Sphaerotheca fuliginea / Podosphaera xanthii*) (*Erysiphe cichoracearum*)	4 to 6 fl. oz. per acre
	Gummy stem blight – suppression (*Didymella bryonae*) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Dry bulb onion Garlic Great-headed (elephant) garlic Shallot	White rot *(Sclerotium cepivorum)*	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii*)	4 to 6 fl. oz. per acre
	White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Tebucon® 3.6F Fungicide per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre. Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Tebucon® 3.6F Fungicide per acre per application. Repeat at an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 32.5 fl. oz. Tebucon® 3.6F Fungicide per acre per season if an in-furrow treatment is made. If Tebucon® 3.6F Fungicide is not applied as an in-furrow treatment then do not apply more than 12 fl oz. Tebucon® 3.6F Fungicide per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Grasses Grown For Seed Welsh onion	Rusts *(Puccinia* spp.*)*	4 to 8 fl. oz. per acre
	Apply the specified rate of Tebucon® 3.6F Fungicide as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl. oz. per acre
	Apply specified rate of Tebucon® 3.6F Fungicide when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
General Comments: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control. For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon® 3.6F Fungicide. A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Tebucon® 3.6F Fungicide may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application. Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (*Sclerotium cepivorum*) suppression only Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii)*	4 to 6 fl. oz. per acre
	For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 24 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Hops	Powdery mildew (*Sphaerotheca humuli/ Spharerotheca macularis)*	4 to 8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (*Cercospora brassicicola*) Powdery mildew (*Erysiphe cruciferarum*) Alternaria leaf spot (*Alternaria brassicicola*)	3 to 4 fl. oz. per acre
	Notes: Do not apply more than 16 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). Do not apply more often than once every 10 days.	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restriction: Application to turnip greens is limited to East of the Rockies.
Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Garden beet roots and tops (leaves)	Cercospora leaf spot (*Cercospora beticola*)	3 to 7.2 fl. oz. per acre
	Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Lychee	Anthracnose (*Colletotrichum gloesporioides)*	4 to 6 fl. oz. per acre
	Notes: Begin first application of Tebucon® 3.6F Fungicide as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Tebucon® 3.6F Fungicide per acre per season. Tebucon® 3.6F Fungicide can be applied up to and including the day of harvest (PHI = 0 days).	

General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 2 days.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Okra	Cercospora leaf spot *(Cercospora* spp.*)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Peanut	**SOILBORNE:** Sclerotium stem and pod rot (white mold, southern blight, southern stern rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) **FOLIAR:** Early leaf spot Late Leaf spot Leaf rust Web blotch (*Phoma*) Pepper spot (*Leptoshaerulina*)	7.2 fl. oz. per acre
	FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Tebucon® 3.6F Fungicide to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Tebucon® 3.6F Fungicide in the first advisory spray in July and continue Tebucon® 3.6F Fungicide applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.	

General Comments: For optimum control of the specified soilborne diseases, four consecutive applications of Tebucon® 3.6F Fungicide must be made at 14-day intervals.

A maximum of 28.8 fluid ounces of Tebucon® 3.6F Fungicide may be applied per crop season. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.

Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Tebucon® 3.6F Fungicide as a leaf spot resistance management strategy. A spray surfactant is not necessary when Tebucon® 3.6F Fungicide is tank mixed with chlorothalonil. Mixing or alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance.

Tebucon® 3.6F Fungicide must be carried by rainfall or irrigation into the root and pod zone for

control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Tebucon® 3.6F Fungicide against the root and pod rots.

Use Tebucon® 3.6F Fungicide in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.

Restricted-entry interval (REI) = 12 hours.

Timing of Tebucon® 3.6F Fungicide Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	TEBUCON® 3.6F Fungicide Application No.	Chlorothalonil Application No.
7 Applications	3,4,5 and 6	1,2 and 7

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCON® 3.6F Fungicide
Pecan	Brown leaf spot (*Sirosporium diffusium)* Downy spot *(Mycosphaerella caryigena)* Liver spot *(Gnomonia caryae)* Scab *(Cladosporium caryigenum)* Vein spot *(Gnomonia nerviseda)* Zonate leaf spot *(Grovesinia pyramidalis)*	4 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Tebucon® 3.6F Fungicide should be applied at 4 fl. oz. per acre in a tank-mix with the labeled rate of Super-Tin® in cover sprays. Follow label directions for the use of SuperTin. Do not add a surfactant to the spray solution when tank-mixing Tebucon® 3.6F Fungicide with SuperTin. Apply Tebucon® 3.6F Fungicide in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Tebucon® 3.6F Fungicide to full-size mature trees, and 4 to 6 fl. oz. per acre of Tebucon® 3.6F Fungicide to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE(S)**	**RATE OF TEBUCON® 3.6F Fungicide**
Soybean	Rust *(Phakopsora pachyrhizi)* Powdery Mildew *(Microsphaera diffusa)*	3 to 4 fl. oz. per acre

Use Directions: Apply Tebucon® 3.6F Fungicide as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant must be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.
Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Sunflower	Rust *(Puccinia helianthi)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of Tebucon® 3.6F Fungicide per acre per season or within 50 days of harvest.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Contact your state Extension Service or Repar representative for a list of approved surfactants. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Turnip (Application is limited to East of the Rockies	Cercospora leaf spot *(Cercospora brassicicola)*	4 to 7.2 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Wheat	Rusts leaf, stem, and stripe *(Puccinia spp.)* Head blight or scab *(Fusarium spp.)* - Suppression	4 fl. oz. per acre
	Notes: Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Tebucon® 3.6F Fungicide. Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn)
For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.

SEED LABELING: To meet U.S. Federal Seed Act requirements, all seed treated with Tebucon® 3.6F Fungicide must be labeled:
"TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with the fungicide Tebuconazole. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice."
USE PRECAUTION: When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

DISEASE	RATE Fl Oz/CWT	DIRECTIONS FOR USE
Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Tebucon® 3.6F Fungicide. The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut *(Sphacelotheca reilana)*	0.27 – 0.54	

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition

(approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR CORP.

Sub label B

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on ornamentals and golf courses

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS: 61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by

avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Chemigation: Do not apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- Do not apply within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.

Spray Drift Management
Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.
Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground.
Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Spray Volume: For best results Tebucon® 3.6F Fungicide may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are recommended depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:
Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1'/2 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable

compatibility agent and use the compatibility agent as directed on its label.

Mixing: Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that the Tebucon® 3.6F Fungicide is complete dispersed in the mixture.

Recommended Mixing Procedure:

1) **Water.** Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2) Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3) **Inductor.** If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA Bags.** Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water Dispersible Products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrates. when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable)
9) **Remaining quantity of water.**

Resistance Management Information
The active ingredient in Tebucon® 3.6F Fungicide is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and exhibits no known cross-resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. Because the speed and scope of resistant population development can not be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS AND PRECAUTIONS

For use on golf course turf only.
Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
Not for homeowner use.
Not for use on turf being grown for sale or commercial use as sod.
Do not use clippings for animal feed.
Do not exceed 3.6 fl.oz. of Tebucon® 3.6F Fungicide per 1,000 sq ft per year.
Do not apply more than 6 applications per year.
* Do not apply more than 3 applications per year in New York State.

Product Information

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Tebucon® 3.6F Fungicide is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to Tebucon® 3.6F Fungicide under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85 degrees F.

Tebucon® 3.6F Fungicide can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 21 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Tebucon® 3.6F Fungicide in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Tebucon® 3.6F Fungicide should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Dollar Spot (*Sclerotinia homoeocarpa*) Copper Spot (*Gloeocercospora sorghi*) Powdery Mildew (*Erysiphe graminis*) Corticium Red Thread (*Laetisaria fuciformis*) Rusts (*Puccinia* spp.)	0.6-1.1	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days.
Brown Patch/Rhizoctonia Blight, Large Patch (*Rhizoctonia solani*) Brown Ring Patch (*R.. circinata*)	0.6-1.1	
Anthracnose -Basal and Foliar (*Colletotrichum cereale*) Red Thread (*Laetisaria fuciformis*) Pink Patch (*Limonomyces rosipellis*)	0.6-1.1	
Bermuda Grass decline (*Gaeumannomyces graminis* var. *graminis*)	0.6-1.1	Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21 day intervals.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Take All Patch (*Gaeumannomyces graminis*)	0.6-1.1	For prevention, apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (*Pyricularia grisea*)	0.6-1.1	Apply when conditions are favorable for disease development at 21 day intervals. Under conditions favoring moderate to heavy disease pressure, Tebucon® 3.6F Fungicide can be tank mixed with a registered contact fungicide at the label rate.
Stipe Smut (*Ustilago striiformis*)	0.6-1.1	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot (*Leptosphaeria korrea, L. narmari, Ophiosphaerella herpotricha, Gaeumannomyces graminis*) Necrotic Ring Spot (*Leptosphaeria korrea*)	0.6-1.1	For prevention, apply in fall when soil temperature reach 65^0 F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (*Fusarium roseum*)	0.6-1.1	Apply first application in mid-June or 21 days prior to time this blight normally becomes evident. Make applications at no less than 21 day intervals.
Summer Patch (*Magnaporthe poae*)	0.6-1.1	Apply beginning in the spring. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 21 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Zoysia Patch, Large Patch of zoysia (*Rhizoctonia solani*)	0.6-1.1	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/ Typhula Blight (*Typhula incarnate*) Pink Snow Mold/Microdochium Patch *(Microdochium nivalis)*	0.6-1.1	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that Tebucon® 3.6F Fungicide be tank-mixed with other registered snow mold products for best season long results.
NOTE: Apply the specified amount of Tebucon® 3.6F Fungicide in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl. oz. of Tebucon® 3.6F Fungicide per 1000 sq. ft. per year. Do not exceed 6 applications per year.		

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS

For use on ornamental plants only; not for woodlands or forest management.
Not for homeowner use.
Do not apply more than 10 fl oz per acre in a single application.
Do not apply more than 0.31 gallons (40 fl oz) of Tebucon® 3.6F Fungicide (equal to 1.13 lbs of tebuconazole) per acre per year.
Do not make more than 4 applications per year at highest rate.
Do not apply to bearing fruit trees or vegetables

Tebucon® 3.6F Fungicide can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Tebucon® 3.6F Fungicide is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Tebucon® 3.6F Fungicide at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern and Geraniums.

Ornamentals Disease Control

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	**To Treat Existing Disease**
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management.	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthracnose Leaf Spot Powdery Mildew		
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust Southern Blight	Petal Blight - Apply 2 – 3 times per week into the flowers as the open and develop color.	
Ground Covers and Vines			
For small plantings: Add 1 teaspoon to 2.5 gallons of water.			

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR.

Material Sent for Data Extraction

Reg. # 69361-11

Description: Revised label- fixed sweet corn RDI

☒ Material(s) Sent to Data Extraction Contractors:

- ☒ New Stamped Label Dated 12-18-12
- ☐ Notification Dated _____
- ☐ New CSF(s) Dated _____
- ☐ Other: _____

☒ Decision #: 472982

☐ Other Action/Comments: _____

File this coversheet and attached materials in the jacket. It must be well organized and clipped together, NOT STAPLED. Then give the jacket with the coversheet and materials to staff in the Information Services Center (ISC) (Room S-4900). If a jacket is full or only available as an image, please file materials in a new jacket and bring it down to the (ISC). For further information please call 703-605-0716.

Reviewer: Robert Westin

Phone: 305-5721 Division: RD/FB

Date: Dec 19, 2012

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

DEC 18 2012

Repar Corporation
% N. Bhushan Mandava
Mandava Associates, LLC
1050 Connecticut Avenue, NW
Suite 1000
Washington, DC 20036

Product Name:	Tebucon 3.6F Fungicide
EPA Reg. No.:	69361-11
Subject:	Your amendment dated December 12, 2012
OPP Decision Number:	472982

Dear Dr. Mandava:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following changes are made:

1. On page 14 in the first column of the box delete "Welch onion"

2. On page 15 in the first column of the box for green onions add "Welsh onion"

One copy of the label stamped "Accepted with comments" is enclosed for your records. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,

Mary L. Waller

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7504P)

Enclosure

ACCEPTED
with COMMENTS
In EPA Letter Dated:
DEC 18 2012
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 69361-11

REPAR

Master Label

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

Sub label A: Agricultural Uses

A. Vegetable Crops including Asparagus, Beans (fresh and dry except succulent shelled), Cucurbit Vegetable Group, Dry Bulb Onion, Garlic, Great Headed (Elephant) Garlic, Welsh Onion, Shallot, Green Onion, Leek, Spring Onion, Scallion, Japanese Bunching Onion, Green Shallots, Green Eschalots, Garden Beet, Leafy Brassica Greens Group, Okra, Turnip.
B. Field Crops including Barley, Corn (sweet corn, field corn, field corn grown for seed and popcorn), Cotton, Grasses Grown for Seed, Peanuts, Soybeans, Sunflower, Wheat and Seed Treatment (sweet corn, field corn, field corn grown for seed and popcorn).
C. Fruit and Nut Crops including Lychee, and Pecan.
D. Miscellaneous Crops: Hops

Sub label B: Turf and Ornamental Uses

A. Disease Control in Golf Course Turf
B. Disease Control in Field, Nursery and Container Ornamentals and Commercial and Residential Landscapes including Roses, Flowers, Ornamental Crabapples, Dogwoods, and Other Landscape Trees, Azales, Camellias, Rhododendrons and Other Landscape Ornamental Shrubs, Ground Covers, Vines

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

Manufactured for:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11	NET CONTENTS: 2.5 GALLONS	EPA Est. No.

Sub label A

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on various agricultural crops

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%

OTHER INGREDIENTS: 61.3%

TOTAL.....100.0%

Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use

KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:

Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential

for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30

seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Spray Volume: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Chemigation: Apply Tebucon® 3.6F Fungicide through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.
Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add labeled amount of Tebucon® 3.6F Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Tebucon® 3.6F Fungicide should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

Compatibility Test for Mix Components:
Before mixing components, always perform a compatibility jar test. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water in a clear, clean, mixing jar. For other spray volumes, adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 2 teaspoons for each pound of dry product or 1 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions.
When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order:

1) **Water**. Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2) **Agitation**. Maintain constant agitation throughout mixing and application.
3) **Inductor**. If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA bags**. Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water-dispersible products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC), or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrate when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable).
9) **Remaining quantity of water.**

Maintain constant agitation during application.

SPRAY AND DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE Standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Additional requirements for aerial applications:
The boom length must not exceed 75% of the wingspan or rotor diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Applications greater than 10 feet above the canopy should be avoided.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications during temperature inversions.

Additional requirements for ground boom application:
Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

AERIAL DRIFT REDUCTION ADVISORY

This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply medium to large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

WIND

Do not apply when wind velocity exceeds 15 mph. Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HIMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Do not make applications during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions may be identified by temperatures that rise with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Resistance Management Statement

Tebucon® 3.6F Fungicide is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Tebucon® 3.6F Fungicide and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Repar Corporation encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Asparagus	Rust (*Puccinia* spp.)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Tebucon® 3.6F Fungicide per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).	
General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance. Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Barley	Rusts (*Puccinia* spp.)	4 fl. oz. per acre
	Head blight *(Fusarium* spp.*)* – Suppression	
	Notes: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Tebucon® 3.6F Fungicide. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Beans (fresh & dry except succulent shelled)	Rust (*Uromyces appendiculatus*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Beans, dry: Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (*Puccinia* spp.) Northern leaf blight (*Helminthosporium turcicum*) Southern leaf blight (*Helminthosporium maydis*) Northern leaf spot (*Helminthosporium carbonum*) Gray leaf spot (*Cercospora zeae-aydis*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Sweet corn: Tebucon® 3.6F Fungicide may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: Tebucon® 3.6F Fungicide may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) for sweet corn = 19 days. Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cotton	Southwestern cotton rust (*Puccinia cacabata*)	6 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Tebucon® 3.6F Fungicide may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew (*Sphaerotheca fuliginea* / *Podosphaera xanthii*) (*Erysiphe cichoracearum*)	4 to 6 fl. oz. per acre
	Gummy stem blight – suppression (*Didymella bryonae*) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot *(Sclerotium cepivorum)*	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii*)	4 to 6 fl. oz. per acre
	White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Tebucon® 3.6F Fungicide per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre. Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Tebucon® 3.6F Fungicide per acre per application. Repeat at an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 32.5 fl. oz. Tebucon® 3.6F Fungicide per acre per season if an in-furrow treatment is made. If Tebucon® 3.6F Fungicide is not applied as an in-furrow treatment then do not apply more than 12 fl oz. Tebucon® 3.6F Fungicide per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Grasses Grown For Seed	Rusts *(Puccinia* spp.*)*	4 to 8 fl. oz. per acre
	Apply the specified rate of Tebucon® 3.6F Fungicide as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl. oz. per acre
	Apply specified rate of Tebucon® 3.6F Fungicide when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	

General Comments: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.

For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon® 3.6F Fungicide.

A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Tebucon® 3.6F Fungicide may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (*Sclerotium cepivorum*) suppression only Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii)*	4 to 6 fl. oz. per acre
	For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 24 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Hops	Powdery mildew (*Sphaerotheca humuli/ Spharerotheca macularis)*	4 to 8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (*Cercospora brassicicola*) Powdery mildew (*Erysiphe cruciferarum*) Alternaria leaf spot (*Alternaria brassicicola*)	3 to 4 fl. oz. per acre
	Notes: Do not apply more than 16 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). Do not apply more often than once every 10 days.	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restriction: Application to turnip greens is limited to East of the Rockies.
Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Garden beet roots and tops (leaves)	Cercospora leaf spot (*Cercospora beticola*)	3 to 7.2 fl. oz. per acre
	Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Lychee	Anthracnose (*Colletotrichum gloesporioides)*	4 to 6 fl. oz. per acre
	Notes: Begin first application of Tebucon® 3.6F Fungicide as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Tebucon® 3.6F Fungicide per acre per season. Tebucon® 3.6F Fungicide can be applied up to and including the day of harvest (PHI = 0 days).	

General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 2 days.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Okra	Cercospora leaf spot *(Cercospora* spp.*)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Peanut	**SOILBORNE:** Sclerotium stem and pod rot (white mold, southern blight, southern stern rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) **FOLIAR:** Early leaf spot Late Leaf spot Leaf rust Web blotch (*Phoma*) Pepper spot (*Leptoshaerulina*)	7.2 fl. oz. per acre
	FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Tebucon® 3.6F Fungicide to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Tebucon® 3.6F Fungicide in the first advisory spray in July and continue Tebucon® 3.6F Fungicide applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.	

General Comments: For optimum control of the specified soilborne diseases, four consecutive applications of Tebucon® 3.6F Fungicide must be made at 14-day intervals.

A maximum of 28.8 fluid ounces of Tebucon® 3.6F Fungicide may be applied per crop season. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.

Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Tebucon® 3.6F Fungicide as a leaf spot resistance management strategy. A spray surfactant is not necessary when Tebucon® 3.6F Fungicide is tank mixed with chlorothalonil. Mixing or alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance.

Tebucon® 3.6F Fungicide must be carried by rainfall or irrigation into the root and pod zone for

control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Tebucon® 3.6F Fungicide against the root and pod rots.

Use Tebucon® 3.6F Fungicide in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.

Restricted-entry interval (REI) = 12 hours.

Timing of Tebucon® 3.6F Fungicide Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	TEBUCON® 3.6F Fungicide Application No.	Chlorothalonil Application No.
7 Applications	3,4,5 and 6	1,2 and 7

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCON® 3.6F Fungicide
Pecan	Brown leaf spot (*Sirosporium diffusium*) Downy spot *(Mycosphaerella caryigena)* Liver spot *(Gnomonia caryae)* Scab *(Cladosporium caryigenum)* Vein spot *(Gnomonia nerviseda)* Zonate leaf spot *(Grovesinia pyramidalis)*	4 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Tebucon® 3.6F Fungicide should be applied at 4 fl. oz. per acre in a tank-mix with the labeled rate of Super-Tin® in cover sprays. Follow label directions for the use of SuperTin. Do not add a surfactant to the spray solution when tank-mixing Tebucon® 3.6F Fungicide with SuperTin. Apply Tebucon® 3.6F Fungicide in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Tebucon® 3.6F Fungicide to full-size mature trees, and 4 to 6 fl. oz. per acre of Tebucon® 3.6F Fungicide to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE(S)**	**RATE OF TEBUCON® 3.6F Fungicide**
Soybean	Rust *(Phakopsora pachyrhizi)* Powdery Mildew *(Microsphaera diffusa)*	3 to 4 fl. oz. per acre

Use Directions: Apply Tebucon® 3.6F Fungicide as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant must be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.
Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Sunflower	Rust *(Puccinia helianthi)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of Tebucon® 3.6F Fungicide per acre per season or within 50 days of harvest.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Contact your state Extension Service or Repar representative for a list of approved surfactants. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Turnip (Application is limited to East of the Rockies	Cercospora leaf spot *(Cercospora brassicicola)*	4 to 7.2 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Wheat	Rusts leaf, stem, and stripe *(Puccinia spp.)* Head blight or scab *(Fusarium spp.)* - Suppression	4 fl. oz. per acre
	Notes: Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Tebucon® 3.6F Fungicide. Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn) For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.

SEED LABELING: To meet U.S. Federal Seed Act requirements, all seed treated with Tebucon® 3.6F Fungicide must be labeled:
"TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with the fungicide Tebuconazole. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice."
USE PRECAUTION: When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

DISEASE	RATE Fl Oz/CWT	DIRECTIONS FOR USE
Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Tebucon® 3.6F Fungicide. The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut *(Sphacelotheca reilana)*	0.27 – 0.54	

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition

(approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR CORP.

Sub label B

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on ornamentals and golf courses

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS: 61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by

avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Chemigation: Do not apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- Do not apply within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.

Spray Drift Management
Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.
Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground.
Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Spray Volume: For best results Tebucon® 3.6F Fungicide may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are recommended depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:
Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1'/2 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable

compatibility agent and use the compatibility agent as directed on its label.

Mixing: Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that the Tebucon® 3.6F Fungicide is complete dispersed in the mixture.

Recommended Mixing Procedure:

1) **Water.** Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2) Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3) **Inductor.** If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA Bags.** Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water Dispersible Products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrates. when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable)
9) **Remaining quantity of water.**

Resistance Management Information
The active ingredient in Tebucon® 3.6F Fungicide is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and exhibits no known cross-resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. Because the speed and scope of resistant population development can not be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS AND PRECAUTIONS
For use on golf course turf only.
Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
Not for homeowner use.
Not for use on turf being grown for sale or commercial use as sod.
Do not use clippings for animal feed.
Do not exceed 3.6 fl.oz. of Tebucon® 3.6F Fungicide per 1,000 sq ft per year.
Do not apply more than 6 applications per year.
* Do not apply more than 3 applications per year in New York State.

Product Information

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Tebucon® 3.6F Fungicide is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to Tebucon® 3.6F Fungicide under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85 degrees F.

Tebucon® 3.6F Fungicide can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Tebucon® 3.6F Fungicide in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Tebucon® 3.6F Fungicide should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Dollar Spot (*Sclerotinia homoeocarpa*) Copper Spot (*Gloeocercospora sorghi*) Powdery Mildew (*Erysiphe graminis*) Corticium Red Thread (*Laetisaria fuciformis*) Rusts (*Puccinia* spp.)	0.6	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.
Brown Patch/Rhizoctonia Blight, Large Patch (*Rhizoctonia solani*) Brown Ring Patch (*R.. circinata*)	0.6	
Anthracnose -Basal and Foliar (*Colletotrichum cereale*) Red Thread (*Laetisaria fuciformis*) Pink Patch (*Limonomyces rosipellis*)	0.6	
Bermuda Grass decline (*Gaeumannomyces graminis* var. *graminis*)	0.6	Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 28 day intervals.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Take All Patch (*Gaeumannomyces graminis*)	0.6	For prevention, apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (*Pyricularia grisea*)	0.6	Apply when conditions are favorable for disease development at 28 day intervals. Under conditions favoring moderate to heavy disease pressure, Tebucon® 3.6F Fungicide can be tank mixed with a registered contact fungicide at the label rate.
Stipe Smut (*Ustilago striiformis*)	0.6	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot (*Leptosphaeria korrea, L. narmari, Ophiosphaerella herpotricha, Gaeumannomyces graminis*) Necrotic Ring Spot (*Leptosphaeria korrea*)	0.6	For prevention, apply in fall when soil temperature reach 65^0 F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (*Fusarium roseum*)	0.6	Apply first application in mid-June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28 day intervals.
Summer Patch (*Magnaporthe poae*)	0.6	Apply beginning in the spring. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Zoysia Patch, Large Patch of zoysia (*Rhizoctonia solani*)	0.6	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/ Typhula Blight (*Typhula incarnate*) Pink Snow Mold/Microdochium Patch *(Microdochium nivalis)*	0.6	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that Tebucon® 3.6F Fungicide be tank-mixed with other registered snow mold products for best season long results.
NOTE: Apply the specified amount of Tebucon® 3.6F Fungicide in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl. oz. of Tebucon® 3.6F Fungicide per 1000 sq. ft. per year. Do not exceed 6 applications per year.		

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS

For use on ornamental plants only; not for woodlands or forest management.
Not for homeowner use.
Do not apply more than 10 fl oz per acre in a single application.
Do not apply more than 0.31 gallons (40 fl oz) of Tebucon® 3.6F Fungicide (equal to 1.13 lbs of tebuconazole) per acre per year.
Do not make more than 4 applications per year at highest rate.
Do not apply to bearing fruit trees or vegetables

Tebucon® 3.6F Fungicide can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Tebucon® 3.6F Fungicide is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Tebucon® 3.6F Fungicide at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern and Geraniums.

Ornamentals Disease Control

<table>
<tr><th>PLANTS</th><th>DISEASE</th><th colspan="2">APPLICATION</th></tr>
<tr><th></th><th></th><th>To Prevent Diseases</th><th>To Treat Existing Disease</th></tr>
<tr><td>Roses</td><td>Black Spot
Powdery Mildew
Rust</td><td>Apply every 14-21 days during the growing season, starting when leaves first appear.</td><td rowspan="5">Apply every 14 days for a total of 3 applications beginning at the first sign of disease.</td></tr>
<tr><td>Flowers</td><td>Leaf Spot
Powdery Mildew
Rust
Southern Blight</td><td rowspan="2">Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management.</td></tr>
<tr><td>Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees</td><td>Anthracnose
Leaf Spot
Powdery Mildew</td></tr>
<tr><td>Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs</td><td rowspan="2">Anthracnose
Black Spot
Leaf Spot
Petal Blight
Powdery Mildew
Rust
Southern Blight</td><td rowspan="2">Petal Blight - Apply 2 – 3 times per week into the flowers as the open and develop color.</td></tr>
<tr><td>Ground Covers and Vines</td></tr>
<tr><td colspan="4">For small plantings: Add 1 teaspoon to 2.5 gallons of water.</td></tr>
</table>

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR.

RE: Restricted Entry Interval Error on Your label - EPA Reg. No. 69361-11, -27
Srinivas Mandava, M.D.
to:
Mary Waller, Bhushan Mandava
12/12/2012 05:02 PM
Cc:
Cynthia Giles-Parker, Shaunta Hill, Heather Garvie, Robert Westin, Tamue Gibson
Hide Details
From: "Srinivas Mandava, M.D." Sort List...

To: Mary Waller/DC/USEPA/US@EPA, "Bhushan Mandava"

Cc: Cynthia Giles-Parker/DC/USEPA/US@EPA, Shaunta Hill/DC/USEPA/US@EPA, Heather Garvie/DC/USEPA/US@EPA, Robert Westin/DC/USEPA/US@EPA, Tamue Gibson/DC/USEPA/US@EPA

D472982
5928019

Dear Ms. Waller:

In response to your email from earlier today we are submitting the requested information and this is the 2nd of 2 emails for the 2 products of Tebucon 3.6F Fungicide (EPA Reg. No. 69361-11) and Tebuconazole 3.6F Fungicide (EPA Reg. No. 69361-27).

Please find enclosed with this email the following for Tebucon 3.6F Fungicide (EPA Reg. No. 69361-11) label amendment:

1) Application form
2) Highlighted PDF file with the corrections to the REI interval for sweet corn on page 12 of Sublabel A and the Agricultural Use Requirements Box statement on Sublabel B page 27
3) Clean label with the corrections on page 12 and 27.

Thank you for your email in bringing this to our attention.

Regards,

Srinivas Mandava, M.D.
Mandava Associates, LLC
1050 Connecticut Avenue, N.W.
Suite 1000
Washington, D.C. 20036
Tel: (202) 223 - 1424
Fax: (202) 223 – 0141

****
Notice: This E-mail (including attachments) is covered by the Electronic Communications Privacy Act, 18 U.S.C. §§2510-2521, is confidential and may be legally privileged. If you are not the intended recipient, you are hereby notified that any retention, dissemination, distribution, or copying of this communication is strictly prohibited. Please reply to the sender that you have received the message in error and then delete it. Thank you.
****

From: Waller.Mary@epamail.epa.gov [mailto:Waller.Mary@epamail.epa.gov]
Sent: Wednesday, December 12, 2012 2:56 PM
To: Bhushan Mandava; Srinivas Mandava, M.D.
Cc: Giles-Parker.Cynthia@epamail.epa.gov; Hill.Shaunta@epamail.epa.gov; Garvie.Heather@epamail.epa.gov; Westin.Robert@epamail.epa.gov; Gibson.Tamue@epamail.epa.gov
Subject: Restricted Entry Interval Error on Your label - EPA Reg. No. 69361-11, -27

Dear Dr. Bhushan and Srinivas Mandava:

It has come to my attention that your tebuconazole labels contain a serious error, and I am requesting that you submit an amendment to correct the error as soon as possible, or no later than 30 days from the date of this email. If you want to pdf an amendment to me (application form, highlighted label and clean label), you may skip mailing the amendment and email it to me directly. If you choose to email it, you must email it to me by no later than 12/17/12 as I am retiring and no one will have access to my email. If you cannot meet the 12/17/12 date, please mail your amendment in the usual fashion.

EPA Reg. No. 69361-11and 69361-27 (For both registrations -last accepted label dated 9/13/2012);

On page 12 of Sublabel A, the restricted entry interval is incorrect for sweet corn. It currently reads "19 hours" and must be revised to "19 days".

On page 27 of Sublabel B, the Agricultural Use Requirements Box statement indicates that the REI is listed with the individual crops, and it is not. Since the only use subject to the WPS is ornamentals, you must revise the Ag Use Requirements Box statement to read as follows: " Do not enter or allow worker entry into the treated area during the restricted entry interval (REI) of 12 hours."

Mary L. Waller
Product Manager 21
Fungicide Branch
Registration Division
Office of Pesticide Programs
U.S. Environmental Protection Agency
(703) 308-9354

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95

EPA

United States
Environmental Protection Agency
Washington, DC 20460

- [ ] Registration
- [x] Amendment
- [ ] Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 69361-11	2. EPA Product Manager Mary Waller	3. Proposed Classification [x] None [ ] Restricted
4. Company/Product (Name) Repar Corporation/TEBUCON 3.6F FUNGICIDE	PM# 21	
5. Name and Address of Applicant *(Include ZIP Code)* Repar Corporation P.O. Box 4321 Silver Spring, MD. 20914 [ ] *Check if this is a new address*	6. **Expedited Reveiw.** In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

- [x] Amendment - Explain below.
- [ ] Resubmission in response to Agency letter dated _____
- [ ] Notification - Explain below.
- [ ] Final printed labels in repsonse to Agency letter dated _____
- [ ] "Me Too" Application.
- [ ] Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

1) Label amendment for TEBUCON 3.6F FUNGICIDE (EPA Reg. No. 69361-11) in response to the Agency's request of December 12, 2012.
2) Please direct all correspondence to: Mandava Associates, LLC; 1050 Connecticut Avenue, NW / Suite 1000; Washington, D.C. 20036
3 The contact person fax number is (202) 223 - 0141 and e-mail address is: mandava@compuserve.com.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging	Unit Packaging	Water Soluble Packaging	2. Type of Container
[ ] Yes [x] No	[ ] Yes [x] No	[ ] Yes [x] No	[ ] Metal [x] Plastic [ ] Glass [ ] Paper [ ] Other (Specify) _____
* *Certification must be submitted*	If "Yes" Unit Packaging wgt. / No. per container	If "Yes" Package wgt / No. per container	

3. Location of Net Contents Information	4. Size(s) Retail Container	5. Location of Label Directions
[x] Label [ ] Container		[x]

6. Manner in Which Label is Affixed to Product: [ ] Lithograph [x] Paper glued [ ] Stenciled [ ] Other _____

Section - IV

1. Contact Point *(Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)*

Name	Title	Telephone No. (Include Area Code)
N. Bhushan Mandava	Agent for Repar Corporation	(202) 223-1424

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received (Stamped)

2. Signature	3. Title
N. Bhushan Mandava	Agent for Repar Corporation
4. Typed Name	**5. Date**
N. Bhushan Mandava	December 12, 2012

Redline Copy
pp 12, 27

Master Label

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

Sub label A: Agricultural Uses

A. Vegetable Crops including Asparagus, Beans (fresh and dry except succulent shelled), Cucurbit Vegetable Group, Dry Bulb Onion, Garlic, Great Headed (Elephant) Garlic, Welsh Onion, Shallot, Green Onion, Leek, Spring Onion, Scallion, Japanese Bunching Onion, Green Shallots, Green Eschalots, Garden Beet, Leafy Brassica Greens Group, Okra, Turnip.
B. Field Crops including Barley, Corn (sweet corn, field corn, field corn grown for seed and popcorn), Cotton, Grasses Grown for Seed, Peanuts, Soybeans, Sunflower, Wheat and Seed Treatment (sweet corn, field corn, field corn grown for seed and popcorn).
C. Fruit and Nut Crops including Lychee, and Pecan.
D. Miscellaneous Crops: Hops

Sub label B: Turf and Ornamental Uses

A. Disease Control in Golf Course Turf
B. Disease Control in Field, Nursery and Container Ornamentals and Commercial and Residential Landscapes including Roses, Flowers, Ornamental Crabapples, Dogwoods, and Other Landscape Trees, Azales, Camellias, Rhododendrons and Other Landscape Ornamental Shrubs, Ground Covers, Vines

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

Manufactured for:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11	NET CONTENTS: 2.5 GALLONS	EPA Est. No.

Sub label A

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on various agricultural crops

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%

OTHER INGREDIENTS:.....61.3%

TOTAL.....100.0%

Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use

KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:

Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential

for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30

seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Spray Volume: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Chemigation: Apply Tebucon® 3.6F Fungicide through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.
Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add labeled amount of Tebucon® 3.6F Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Tebucon® 3.6F Fungicide should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

Compatibility Test for Mix Components:
Before mixing components, always perform a compatibility jar test. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water in a clear, clean, mixing jar. For other spray volumes, adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 2 teaspoons for each pound of dry product or 1 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions.
When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order:

1) **Water**. Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2) **Agitation**. Maintain constant agitation throughout mixing and application.
3) **Inductor**. If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA bags**. Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water-dispersible products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC), or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrate when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable).
9) **Remaining quantity of water.**

Maintain constant agitation during application.

SPRAY AND DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE Standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Additional requirements for aerial applications:
The boom length must not exceed 75% of the wingspan or rotor diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Applications greater than 10 feet above the canopy should be avoided.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications during temperature inversions.

Additional requirements for ground boom application:
Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

AERIAL DRIFT REDUCTION ADVISORY

This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply medium to large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

WIND

Do not apply when wind velocity exceeds 15 mph. Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HIMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Do not make applications during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions may be identified by temperatures that rise with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Resistance Management Statement

Tebucon® 3.6F Fungicide is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Tebucon® 3.6F Fungicide and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Repar Corporation encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Asparagus	Rust (*Puccinia* spp.)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Tebucon® 3.6F Fungicide per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).	
General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance. Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Barley	Rusts (*Puccinia* spp.) Head blight *(Fusarium* spp.*)* – Suppression	4 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Tebucon® 3.6F Fungicide. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Beans (fresh & dry except succulent shelled)	Rust (*Uromyces appendiculatus*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Beans, dry: Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (*Puccinia* spp.) Northern leaf blight (*Helminthosporium turcicum*) Southern leaf blight (*Helminthosporium maydis*) Northern leaf spot (*Helminthosporium carbonum*) Gray leaf spot (*Cercospora zeae-aydis*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Sweet corn: Tebucon® 3.6F Fungicide may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: Tebucon® 3.6F Fungicide may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) for sweet corn = 19 days. Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cotton	Southwestern cotton rust (*Puccinia cacabata*)	6 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Tebucon® 3.6F Fungicide may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew (*Sphaerotheca fuliginea* / *Podosphaera xanthii*) (*Erysiphe cichoracearum*)	4 to 6 fl. oz. per acre
	Gummy stem blight – suppression (*Didymella bryonae*) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot *(Sclerotium cepivorum)*	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii*)	4 to 6 fl. oz. per acre
	White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Tebucon® 3.6F Fungicide per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre. Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Tebucon® 3.6F Fungicide per acre per application. Repeat at an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 32.5 fl. oz. Tebucon® 3.6F Fungicide per acre per season if an in-furrow treatment is made. If Tebucon® 3.6F Fungicide is not applied as an in-furrow treatment then do not apply more than 12 fl oz. Tebucon® 3.6F Fungicide per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Grasses Grown For Seed	Rusts *(Puccinia* spp.*)*	4 to 8 fl. oz. per acre
	Apply the specified rate of Tebucon® 3.6F Fungicide as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl. oz. per acre
	Apply specified rate of Tebucon® 3.6F Fungicide when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	

General Comments: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.
For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon® 3.6F Fungicide.
A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Tebucon® 3.6F Fungicide may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (*Sclerotium cepivorum*) suppression only Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii)*	4 to 6 fl. oz. per acre
	For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 24 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).
Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Hops	Powdery mildew (*Sphaerotheca humuli/ Spharerotheca macularis)*	4 to 8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (*Cercospora brassicicola*) Powdery mildew (*Erysiphe cruciferarum*) Alternaria leaf spot (*Alternaria brassicicola*)	3 to 4 fl. oz. per acre
	Notes: Do not apply more than 16 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). Do not apply more often than once every 10 days.	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restriction: Application to turnip greens is limited to East of the Rockies.
Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Garden beet roots and tops (leaves)	Cercospora leaf spot (*Cercospora beticola*)	3 to 7.2 fl. oz. per acre
	Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Lychee	Anthracnose (*Colletotrichum gloesporioides)*	4 to 6 fl. oz. per acre
	Notes: Begin first application of Tebucon® 3.6F Fungicide as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Tebucon® 3.6F Fungicide per acre per season. Tebucon® 3.6F Fungicide can be applied up to and including the day of harvest (PHI = 0 days).	
General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 2 days.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Okra	Cercospora leaf spot *(Cercospora* spp.*)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Peanut	**SOILBORNE:** Sclerotium stem and pod rot (white mold, southern blight, southern stern rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) **FOLIAR:** Early leaf spot Late Leaf spot Leaf rust Web blotch (*Phoma*) Pepper spot (*Leptoshaerulina*)	7.2 fl. oz. per acre
	FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Tebucon® 3.6F Fungicide to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Tebucon® 3.6F Fungicide in the first advisory spray in July and continue Tebucon® 3.6F Fungicide applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.	

General Comments: For optimum control of the specified soilborne diseases, four consecutive applications of Tebucon® 3.6F Fungicide must be made at 14-day intervals.

A maximum of 28.8 fluid ounces of Tebucon® 3.6F Fungicide may be applied per crop season. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.

Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Tebucon® 3.6F Fungicide as a leaf spot resistance management strategy. A spray surfactant is not necessary when Tebucon® 3.6F Fungicide is tank mixed with chlorothalonil. Mixing or alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance.

Tebucon® 3.6F Fungicide must be carried by rainfall or irrigation into the root and pod zone for

control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Tebucon® 3.6F Fungicide against the root and pod rots.

Use Tebucon® 3.6F Fungicide in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.

Restricted-entry interval (REI) = 12 hours.

Timing of Tebucon® 3.6F Fungicide Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	TEBUCON® 3.6F Fungicide Application No.	Chlorothalonil Application No.
7 Applications	3,4,5 and 6	1,2 and 7

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCON® 3.6F Fungicide
Pecan	Brown leaf spot (*Sirosporium diffusium)* Downy spot *(Mycosphaerella caryigena)* Liver spot *(Gnomonia caryae)* Scab *(Cladosporium caryigenum)* Vein spot *(Gnomonia nerviseda)* Zonate leaf spot *(Grovesinia pyramidalis)*	4 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Tebucon® 3.6F Fungicide should be applied at 4 fl. oz. per acre in a tank-mix with the labeled rate of Super-Tin® in cover sprays. Follow label directions for the use of SuperTin. Do not add a surfactant to the spray solution when tank-mixing Tebucon® 3.6F Fungicide with SuperTin. Apply Tebucon® 3.6F Fungicide in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Tebucon® 3.6F Fungicide to full-size mature trees, and 4 to 6 fl. oz. per acre of Tebucon® 3.6F Fungicide to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE(S)**	**RATE OF TEBUCON® 3.6F Fungicide**
Soybean	Rust *(Phakopsora pachyrhizi)* Powdery Mildew *(Microsphaera diffusa)*	3 to 4 fl. oz. per acre

Use Directions: Apply Tebucon® 3.6F Fungicide as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant must be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.
Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Sunflower	Rust *(Puccinia helianthi)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of Tebucon® 3.6F Fungicide per acre per season or within 50 days of harvest.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Contact your state Extension Service or Repar representative for a list of approved surfactants. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Turnip (Application is limited to East of the Rockies	Cercospora leaf spot *(Cercospora brassicicola)*	4 to 7.2 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Wheat	Rusts leaf, stem, and stripe *(Puccinia spp.)* Head blight or scab *(Fusarium spp.)* - Suppression	4 fl. oz. per acre
	Notes: Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Tebucon® 3.6F Fungicide. Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn)
For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.

SEED LABELING: To meet U.S. Federal Seed Act requirements, all seed treated with Tebucon® 3.6F Fungicide must be labeled:
"TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with the fungicide Tebuconazole. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice."
USE PRECAUTION: When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

DISEASE	RATE Fl Oz/CWT	DIRECTIONS FOR USE
Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Tebucon® 3.6F Fungicide. The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut *(Sphacelotheca reilana)*	0.27 – 0.54	

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition

(approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR CORP.

Sub label B

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on ornamentals and golf courses

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS: 61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by

avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Chemigation: Do not apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- Do not apply within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.

Spray Drift Management
Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.
Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground.
Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Spray Volume: For best results Tebucon® 3.6F Fungicide may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are recommended depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:
Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1'/2 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable

compatibility agent and use the compatibility agent as directed on its label.

Mixing: Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that the Tebucon® 3.6F Fungicide is complete dispersed in the mixture.

Recommended Mixing Procedure:

1) **Water.** Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2) Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3) **Inductor.** If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA Bags.** Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water Dispersible Products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrates. when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable)
9) **Remaining quantity of water.**

Resistance Management Information
The active ingredient in Tebucon® 3.6F Fungicide is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and exhibits no known cross-resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. Because the speed and scope of resistant population development can not be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS AND PRECAUTIONS

For use on golf course turf only.
Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
Not for homeowner use.
Not for use on turf being grown for sale or commercial use as sod.
Do not use clippings for animal feed.
Do not exceed 3.6 fl.oz. of Tebucon® 3.6F Fungicide per 1,000 sq ft per year.
Do not apply more than 6 applications per year.
* Do not apply more than 3 applications per year in New York State.

Product Information

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Tebucon® 3.6F Fungicide is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to Tebucon® 3.6F Fungicide under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85 degrees F.

Tebucon® 3.6F Fungicide can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Tebucon® 3.6F Fungicide in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Tebucon® 3.6F Fungicide should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Dollar Spot (*Sclerotinia homoeocarpa*) Copper Spot (*Gloeocercospora sorghi*) Powdery Mildew (*Erysiphe graminis*) Corticium Red Thread (*Laetisaria fuciformis*) Rusts (*Puccinia* spp.)	0.6	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.
Brown Patch/Rhizoctonia Blight, Large Patch (*Rhizoctonia solani*) Brown Ring Patch (*R.. circinata*)	0.6	
Anthracnose -Basal and Foliar (*Colletotrichum cereale*) Red Thread (*Laetisaria fuciformis*) Pink Patch (*Limonomyces rosipellis*)	0.6	
Bermuda Grass decline (*Gaeumannomyces graminis* var. *graminis*)	0.6	Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 28 day intervals.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Take All Patch (*Gaeumannomyces graminis*)	0.6	For prevention, apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (*Pyricularia grisea*)	0.6	Apply when conditions are favorable for disease development at 28 day intervals. Under conditions favoring moderate to heavy disease pressure, Tebucon® 3.6F Fungicide can be tank mixed with a registered contact fungicide at the label rate.
Stipe Smut (*Ustilago striiformis*)	0.6	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot (*Leptosphaeria korrea, L. narmari, Ophiosphaerella herpotricha, Gaeumannomyces graminis*) Necrotic Ring Spot (*Leptosphaeria korrea*)	0.6	For prevention, apply in fall when soil temperature reach 65^0 F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (*Fusarium roseum*)	0.6	Apply first application in mid-June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28 day intervals.
Summer Patch (*Magnaporthe poae*)	0.6	Apply beginning in the spring. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Zoysia Patch, Large Patch of zoysia (*Rhizoctonia solani*)	0.6	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/ Typhula Blight (*Typhula incarnate*) Pink Snow Mold/Microdochium Patch *(Microdochium nivalis)*	0.6	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that Tebucon® 3.6F Fungicide be tank-mixed with other registered snow mold products for best season long results.
NOTE: Apply the specified amount of Tebucon® 3.6F Fungicide in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl. oz. of Tebucon® 3.6F Fungicide per 1000 sq. ft. per year. Do not exceed 6 applications per year.		

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS

For use on ornamental plants only; not for woodlands or forest management.
Not for homeowner use.
Do not apply more than 10 fl oz per acre in a single application.
Do not apply more than 0.31 gallons (40 fl oz) of Tebucon® 3.6F Fungicide (equal to 1.13 lbs of tebuconazole) per acre per year.
Do not make more than 4 applications per year at highest rate.
Do not apply to bearing fruit trees or vegetables

Tebucon® 3.6F Fungicide can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Tebucon® 3.6F Fungicide is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Tebucon® 3.6F Fungicide at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern and Geraniums.

Ornamentals Disease Control

<table>
<tr><th>PLANTS</th><th>DISEASE</th><th colspan="2">APPLICATION</th></tr>
<tr><th></th><th></th><th>To Prevent Diseases</th><th>To Treat Existing Disease</th></tr>
<tr><td>Roses</td><td>Black Spot
Powdery Mildew
Rust</td><td>Apply every 14-21 days during the growing season, starting when leaves first appear.</td><td rowspan="5">Apply every 14 days for a total of 3 applications beginning at the first sign of disease.</td></tr>
<tr><td>Flowers</td><td>Leaf Spot
Powdery Mildew
Rust
Southern Blight</td><td rowspan="2">Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management.</td></tr>
<tr><td>Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees</td><td>Anthracnose
Leaf Spot
Powdery Mildew</td></tr>
<tr><td>Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs</td><td rowspan="2">Anthracnose
Black Spot
Leaf Spot
Petal Blight
Powdery Mildew
Rust
Southern Blight</td><td rowspan="2">Petal Blight - Apply 2 – 3 times per week into the flowers as the open and develop color.</td></tr>
<tr><td>Ground Covers and Vines</td></tr>
<tr><td colspan="4">For small plantings: Add 1 teaspoon to 2.5 gallons of water.</td></tr>
</table>

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR.

Restricted Entry Interval Error on Your label - EPA Reg. No. 69361-11, -27

Mary Waller to: Bhushan Mandava, Srinivas Mandava, M.D. 12/12/2012 02:55 PM
Cc: Cynthia Giles-Parker, Shaunta Hill, Heather Garvie, Robert Westin, Tamue Gibson

From: Mary Waller/DC/USEPA/US
To: Bhushan Mandava <bhushan@Mandava.com>, "Srinivas Mandava, M.D." <Srini@Mandava.com>
Cc: Cynthia Giles-Parker/DC/USEPA/US@EPA, Shaunta Hill/DC/USEPA/US@EPA, Heather Garvie/DC/USEPA/US@EPA, Robert Westin/DC/USEPA/US@EPA, Tamue Gibson/DC/USEPA/US@EPA

Dear Dr. Bhushan and Srinivas Mandava:

It has come to my attention that your tebuconazole labels contain a serious error, and I am requesting that you submit an amendment to correct the error as soon as possible, or no later than 30 days from the date of this email. If you want to pdf an amendment to me (application form, highlighted label and clean label), you may skip mailing the amendment and email it to me directly. If you choose to email it, you must email it to me by no later than 12/17/12 as I am retiring and no one will have access to my email. If you cannot meet the 12/17/12 date, please mail your amendment in the usual fashion.

EPA Reg. No. 69361-11and 69361-27 (For both registrations -last accepted label dated 9/13/2012);

On page 12 of Sublabel A, the restricted entry interval is incorrect for sweet corn. It currently reads "19 hours" and must be revised to "19 days".

On page 27 of Sublabel B, the Agricultural Use Requirements Box statement indicates that the REI is listed with the individual crops, and it is not. Since the only use subject to the WPS is ornamentals, you must revise the Ag Use Requirements Box statement to read as follows: " Do not enter or allow worker entry into the treated area during the restricted entry interval (REI) of 12 hours."

Mary L. Waller
Product Manager 21
Fungicide Branch
Registration Division
Office of Pesticide Programs
U.S. Environmental Protection Agency
(703) 308-9354

Material Sent for Data Extraction

Reg. # 69361-11

Description: Newly stamped label

☒ Material(s) Sent to Data Extraction Contractors:

- ☒ New Stamped Label Dated 9/13/12
- ☐ Notification Dated _____
- ☐ New CSF(s) Dated _____
- ☐ Other: _____

☐ Decision #: 466424

☐ Other Action/Comments: _____

File this coversheet and attached materials in the jacket. It must be well organized and clipped together, NOT STAPLED. Then give the jacket with the coversheet and materials to staff in the Information Services Center (ISC) (Room S-4900). If a jacket is full or only available as an image, please file materials in a new jacket and bring it down to the (ISC). For further information please call 703-605-0716.

Reviewer: H. GARVIE

Phone: 308-0034 Division: RD

Date: 9/18/12

jacket

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

N. Bhushan Mandava
Agent for Repar Corporation
Mandava Associates, LLC
1050 Connecticut Avenue, N.W. Suite 1000
Washington, DC 20036

SEP 13 2012

Subject: Tebucon® 3.6F Fungicide
EPA Reg. No. 69361-11
Your resubmission dated September 5, 2012
EPA Decision Number: 466424

Dear Dr. Mandava:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable. The Agency acknowledges the alternate brand name Tebucon 3.6F Foliar Fungicide.

One copy of the label stamped "Accepted" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label before the product is released for shipment. If you have any questions, please contact Heather Garvie by phone at: 703-308-0034 or via email at: garvie.heather@epa.gov .

Sincerely,

Mary L. Waller

Mary Waller
Product Manager 21
Fungicide Branch
Registration Division (7505P)

Enclosure: Stamped label "Accepted"

Master Label

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

Sub label A: Agricultural Uses

A. Vegetable Crops including Asparagus, Beans (fresh and dry except succulent shelled), Cucurbit Vegetable Group, Dry Bulb Onion, Garlic, Great Headed (Elephant) Garlic, Welsh Onion, Shallot, Green Onion, Leek, Spring Onion, Scallion, Japanese Bunching Onion, Green Shallots, Green Eschalots, Garden Beet, Leafy Brassica Greens Group, Okra, Turnip.
B. Field Crops including Barley, Corn (sweet corn, field corn, field corn grown for seed and popcorn), Cotton, Grasses Grown for Seed, Peanuts, Soybeans, Sunflower, Wheat and Seed Treatment (sweet corn, field corn, field corn grown for seed and popcorn).
C. Fruit and Nut Crops including Lychee, and Pecan.
D. Miscellaneous Crops: Hops

Sub label B: Turf and Ornamental Uses

A. Disease Control in Golf Course Turf
B. Disease Control in Field, Nursery and Container Ornamentals and Commercial and Residential Landscapes including Roses, Flowers, Ornamental Crabapples, Dogwoods, and Other Landscape Trees, Azales, Camellias, Rhododendrons and Other Landscape Ornamental Shrubs, Ground Covers, Vines

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

ACCEPTED
9/13/2012
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 69361-11

Manufactured for:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11	NET CONTENTS: 2.5 GALLONS	EPA Est. No.

Sub label A

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on various agricultural crops

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN

CAUTION

For product use information call 1-866-248-7426

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378	

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart..

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential

for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30

seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Spray Volume: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Chemigation: Apply Tebucon® 3.6F Fungicide through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.
Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add labeled amount of Tebucon® 3.6F Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Tebucon® 3.6F Fungicide should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

Compatibility Test for Mix Components:
Before mixing components, always perform a compatibility jar test. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water in a clear, clean, mixing jar. For other spray volumes, adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 2 teaspoons for each pound of dry product or 1 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions.
When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order:

1) **Water**. Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2) **Agitation**. Maintain constant agitation throughout mixing and application.
3) **Inductor**. If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA bags**. Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water-dispersible products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC), or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrate when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable).
9) **Remaining quantity of water.**

Maintain constant agitation during application.

SPRAY AND DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE Standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Additional requirements for aerial applications:
The boom length must not exceed 75% of the wingspan or rotor diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Applications greater than 10 feet above the canopy should be avoided.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications during temperature inversions.

Additional requirements for ground boom application:
Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

AERIAL DRIFT REDUCTION ADVISORY

This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply medium to large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

WIND

Do not apply when wind velocity exceeds 15 mph. Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HIMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Do not make applications during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions may be identified by temperatures that rise with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Resistance Management Statement

Tebucon® 3.6F Fungicide is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Tebucon® 3.6F Fungicide and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Repar Corporation encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Asparagus	Rust (*Puccinia* spp.)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Tebucon® 3.6F Fungicide per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).	
General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance. Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Barley	Rusts (*Puccinia* spp.) Head blight *(Fusarium* spp.*)* – Suppression	4 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Tebucon® 3.6F Fungicide. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Beans (fresh & dry except succulent shelled)	Rust (*Uromyces appendiculatus*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Beans, dry: Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (*Puccinia* spp.) Northern leaf blight (*Helminthosporium turcicum*) Southern leaf blight (*Helminthosporium maydis*) Northern leaf spot (*Helminthosporium carbonum*) Gray leaf spot (*Cercospora zeae-aydis*)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Sweet corn: Tebucon® 3.6F Fungicide may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: Tebucon® 3.6F Fungicide may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) for sweet corn = 19 hours. Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cotton	Southwestern cotton rust (*Puccinia cacabata*)	6 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Tebucon® 3.6F Fungicide may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew (*Sphaerotheca fuliginea* / *Podosphaera xanthii*) (*Erysiphe cichoracearum*)	4 to 6 fl. oz. per acre
	Gummy stem blight – suppression (*Didymella bryonae*) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot *(Sclerotium cepivorum)*	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii*)	4 to 6 fl. oz. per acre
	White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Tebucon® 3.6F Fungicide per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre. Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Tebucon® 3.6F Fungicide per acre per application. Repeat at an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 32.5 fl. oz. Tebucon® 3.6F Fungicide per acre per season if an in-furrow treatment is made. If Tebucon® 3.6F Fungicide is not applied as an in-furrow treatment then do not apply more than 12 fl oz. Tebucon® 3.6F Fungicide per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Grasses Grown For Seed	Rusts *(Puccinia* spp.*)*	4 to 8 fl. oz. per acre
	Apply the specified rate of Tebucon® 3.6F Fungicide as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl. oz. per acre
	Apply specified rate of Tebucon® 3.6F Fungicide when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	

General Comments: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.

For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon® 3.6F Fungicide.

A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Tebucon® 3.6F Fungicide may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (*Sclerotium cepivorum*) suppression only Rust (*Puccinia allii, Puccinia porri*) Purple blotch (*Alternaria porii)*	4 to 6 fl. oz. per acre
	For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Tebucon® 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 24 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Hops	Powdery mildew (*Sphaerotheca humuli/ Spharerotheca macularis)*	4 to 8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (*Cercospora brassicicola*) Powdery mildew (*Erysiphe cruciferarum*) Alternaria leaf spot (*Alternaria brassicicola*)	3 to 4 fl. oz. per acre
	Notes: Do not apply more than 16 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). Do not apply more often than once every 10 days.	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restriction: Application to turnip greens is limited to East of the Rockies.
Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Garden beet roots and tops (leaves)	Cercospora leaf spot (*Cercospora beticola*)	3 to 7.2 fl. oz. per acre
	Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. Tebucon® 3.6F Fungicide per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Lychee	Anthracnose (*Colletotrichum gloesporioides*)	4 to 6 fl. oz. per acre
	Notes: Begin first application of Tebucon® 3.6F Fungicide as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Tebucon® 3.6F Fungicide per acre per season. Tebucon® 3.6F Fungicide can be applied up to and including the day of harvest (PHI = 0 days).	

General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 2 days.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Okra	Cercospora leaf spot *(Cercospora* spp.*)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Tebucon® 3.6F Fungicide per acre per season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Peanut	**SOILBORNE:** Sclerotium stem and pod rot (white mold, southern blight, southern stern rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) **FOLIAR:** Early leaf spot Late Leaf spot Leaf rust Web blotch (*Phoma*) Pepper spot (*Leptoshaerulina*)	7.2 fl. oz. per acre
	FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Tebucon® 3.6F Fungicide to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Tebucon® 3.6F Fungicide in the first advisory spray in July and continue Tebucon® 3.6F Fungicide applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.	

General Comments: For optimum control of the specified soilborne diseases, four consecutive applications of Tebucon® 3.6F Fungicide must be made at 14-day intervals.

A maximum of 28.8 fluid ounces of Tebucon® 3.6F Fungicide may be applied per crop season. Tebucon® 3.6F Fungicide may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.

Tebucon® 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Tebucon® 3.6F Fungicide as a leaf spot resistance management strategy. A spray surfactant is not necessary when Tebucon® 3.6F Fungicide is tank mixed with chlorothalonil. Mixing or alternating Tebucon® 3.6F Fungicide with other DMI fungicides may lead to resistance.

Tebucon® 3.6F Fungicide must be carried by rainfall or irrigation into the root and pod zone for

control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Tebucon® 3.6F Fungicide against the root and pod rots.

Use Tebucon® 3.6F Fungicide in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.

Restricted-entry interval (REI) = 12 hours.

Timing of Tebucon® 3.6F Fungicide Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	TEBUCON® 3.6F Fungicide Application No.	Chlorothalonil Application No.
7 Applications	3,4,5 and 6	1,2 and 7

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCON® 3.6F Fungicide
Pecan	Brown leaf spot (*Sirosporium diffusium*) Downy spot (*Mycosphaerella caryigena*) Liver spot (*Gnomonia caryae*) Scab (*Cladosporium caryigenum*) Vein spot (*Gnomonia nerviseda*) Zonate leaf spot (*Grovesinia pyramidalis*)	4 to 8 fl. oz. per acre
	Notes: Apply Tebucon® 3.6F Fungicide in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Tebucon® 3.6F Fungicide should be applied at 4 fl. oz. per acre in a tank-mix with the labeled rate of Super-Tin® in cover sprays. Follow label directions for the use of SuperTin. Do not add a surfactant to the spray solution when tank-mixing Tebucon® 3.6F Fungicide with SuperTin. Apply Tebucon® 3.6F Fungicide in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Tebucon® 3.6F Fungicide to full-size mature trees, and 4 to 6 fl. oz. per acre of Tebucon® 3.6F Fungicide to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE(S)**	**RATE OF TEBUCON® 3.6F Fungicide**
Soybean	Rust *(Phakopsora pachyrhizi)* Powdery Mildew *(Microsphaera diffusa)*	3 to 4 fl. oz. per acre

Use Directions: Apply Tebucon® 3.6F Fungicide as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant must be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.
Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Sunflower	Rust *(Puccinia helianthi)*	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon® 3.6F Fungicide at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of Tebucon® 3.6F Fungicide per acre per season or within 50 days of harvest.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Contact your state Extension Service or Repar representative for a list of approved surfactants. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Turnip (Application is limited to East of the Rockies	Cercospora leaf spot *(Cercospora brassicicola)*	4 to 7.2 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Tebucon® 3.6F Fungicide may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Tebucon® 3.6F Fungicide per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	**DISEASE**	**RATE OF TEBUCON® 3.6F Fungicide**
Wheat	Rusts leaf, stem, and stripe *(Puccinia spp.)* Head blight or scab *(Fusarium spp.)* - Suppression	4 fl. oz. per acre
	Notes: Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Tebucon® 3.6F Fungicide may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Tebucon® 3.6F Fungicide. Apply Tebucon® 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air. Application timing directions: Rusts: Apply Tebucon® 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon® 3.6F Fungicide for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon® 3.6F Fungicide. Tebucon® 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon® 3.6F Fungicide will be resistant to weathering. Tebucon® 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn)
For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.

SEED LABELING: To meet U.S. Federal Seed Act requirements, all seed treated with Tebucon® 3.6F Fungicide must be labeled:
"TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with the fungicide Tebuconazole. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice."
USE PRECAUTION: When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

DISEASE	RATE Fl Oz/CWT	DIRECTIONS FOR USE
Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Tebucon® 3.6F Fungicide. The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut *(Sphacelotheca reilana)*	0.27 – 0.54	

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition

(approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR CORP.

Sub label B

TEBUCON® 3.6F Fungicide*

*** Alternate Brand Name: Tebucon® 3.6F Foliar Fungicide**

For control of specified diseases on ornamentals and golf courses

ACTIVE INGREDIENT:
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%
OTHER INGREDIENTS:.....61.3%
TOTAL.....100.0%
Contains 3.6 pounds Tebuconazole per gallon

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN
CAUTION

For product use information call 1-866-248-7426

FIRST AID

If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in the eyes	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency assistance, call the National Pesticide Information Center 1-800-858-7378

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

EPA Reg. No. 69361-11 NET CONTENTS: 2.5 GALLONS EPA Est. No.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by

avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Nonrefillable containers 5 gallons or less:

Container disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use of disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or less:

Container disposal: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of this container.

Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers of 5 gallons or larger:

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Chemigation: Do not apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- Do not apply within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.

Spray Drift Management

Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground.

Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Spray Volume: For best results Tebucon® 3.6F Fungicide may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are recommended depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1'/2 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable

compatibility agent and use the compatibility agent as directed on its label.

Mixing: Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that the Tebucon® 3.6F Fungicide is complete dispersed in the mixture.

Recommended Mixing Procedure:

1) **Water.** Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2) Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3) **Inductor.** If an inductor is used, rinse it thoroughly after each component has been added.
4) **Products in PVA Bags.** Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5) **Water Dispersible Products.** Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6) **Water-soluble products.**
7) **Emulsifiable concentrates** (such as oil concentrates. when applicable).
8) **Water soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable)
9) **Remaining quantity of water.**

Resistance Management Information

The active ingredient in Tebucon® 3.6F Fungicide is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and exhibits no known cross-resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. Because the speed and scope of resistant population development can not be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS AND PRECAUTIONS

For use on golf course turf only.
Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
Not for homeowner use.
Not for use on turf being grown for sale or commercial use as sod.
Do not use clippings for animal feed.
Do not exceed 3.6 fl.oz. of Tebucon® 3.6F Fungicide per 1,000 sq ft per year.
Do not apply more than 6 applications per year.
* Do not apply more than 3 applications per year in New York State.

Product Information

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Tebucon® 3.6F Fungicide is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to Tebucon® 3.6F Fungicide under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85 degrees F.

Tebucon® 3.6F Fungicide can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Tebucon® 3.6F Fungicide in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Tebucon® 3.6F Fungicide should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Dollar Spot (*Sclerotinia homoeocarpa*) Copper Spot (*Gloeocercospora sorghi*) Powdery Mildew (*Erysiphe graminis*) Corticium Red Thread (*Laetisaria fuciformis*) Rusts (*Puccinia* spp.)	0.6	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.
Brown Patch/Rhizoctonia Blight, Large Patch (*Rhizoctonia solani*) Brown Ring Patch (*R.. circinata*)	0.6	
Anthracnose -Basal and Foliar (*Colletotrichum cereale*) Red Thread (*Laetisaria fuciformis*) Pink Patch (*Limonomyces rosipellis*)	0.6	
Bermuda Grass decline (*Gaeumannomyces graminis* var. *graminis*)	0.6	Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 28 day intervals.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Take All Patch (*Gaeumannomyces graminis*)	0.6	For prevention, apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (*Pyricularia grisea*)	0.6	Apply when conditions are favorable for disease development at 28 day intervals. Under conditions favoring moderate to heavy disease pressure, Tebucon® 3.6F Fungicide can be tank mixed with a registered contact fungicide at the label rate.
Stipe Smut (*Ustilago striiformis*)	0.6	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot (*Leptosphaeria korrea, L. narmari, Ophiosphaerella herpotricha, Gaeumannomyces graminis*) Necrotic Ring Spot (*Leptosphaeria korrea*)	0.6	For prevention, apply in fall when soil temperature reach 65^0 F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (*Fusarium roseum*)	0.6	Apply first application in mid-June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28 day intervals.
Summer Patch (*Magnaporthe poae*)	0.6	Apply beginning in the spring. Do not make two consecutive applications of Tebucon® 3.6F Fungicide. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE of TEBUCON® 3.6F Fungicide (Fl. oz./1000 Sq Ft)	NOTES
Zoysia Patch, Large Patch of zoysia (*Rhizoctonia solani*)	0.6	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/ Typhula Blight (*Typhula incarnate*) Pink Snow Mold/Microdochium Patch *(Microdochium nivalis)*	0.6	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that Tebucon® 3.6F Fungicide be tank-mixed with other registered snow mold products for best season long results.
NOTE: Apply the specified amount of Tebucon® 3.6F Fungicide in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl. oz. of Tebucon® 3.6F Fungicide per 1000 sq. ft. per year. Do not exceed 6 applications per year.		

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS

For use on ornamental plants only; not for woodlands or forest management.
Not for homeowner use.
Do not apply more than 10 fl oz per acre in a single application.
Do not apply more than 0.31 gallons (40 fl oz) of Tebucon® 3.6F Fungicide (equal to 1.13 lbs of tebuconazole) per acre per year.
Do not make more than 4 applications per year at highest rate.
Do not apply to bearing fruit trees or vegetables

Tebucon® 3.6F Fungicide can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Tebucon® 3.6F Fungicide is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Tebucon® 3.6F Fungicide at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern and Geraniums.

Ornamentals Disease Control

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management.	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthracnose Leaf Spot Powdery Mildew		
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust Southern Blight	Petal Blight - Apply 2 – 3 times per week into the flowers as the open and develop color.	
Ground Covers and Vines			
For small plantings: Add 1 teaspoon to 2.5 gallons of water.			

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

CONDITIONS OF SALE AND LIMITED WARRANTY:

REPAR CORPORATION warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of REPAR CORPORATION (REPAR) or the SELLER. To the extent consistent with applicable law, Repar Corporation shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, REPAR makes no warranties, guarantees, or representations of any kind, either express or impaired, or by usage of trade, statutory or otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

REPAR and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of REPAR.

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 1.25 hours per response for registration and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460.
Do not send the completed form to this address.

Certification with Respect to Citation of Data

Applicant's/Registrant's Name, Address, and Telephone Number Repar Corporation P.O. Box 4321 Silver Spring, MD 20914; (202) 223-1424	EPA Registration Number/File Symbol 69361-11
Active Ingredient(s) and/or representative test compound(s) Tebuconazole (CAS# 107534-96-3)	Date July 9, 2012
General Use Pattern(s) (list all those claimed for this product using 40 CFR Part 158) Fungicide, Terrestrial, Food and Non-food Crops	Product Name TEBUCON 3.6F FUNGICIDE

NOTE: If your product is a 100% repackaging of another purchased EPA-registered product labeled for all the same uses on your label, you do not need to submit this form. You must submit the Formulator's Exemption Statement (EPA Form 8570-27).

☐ I am responding to a Data-Call-In Notice, and have included with this form a list of companies sent offers of compensation (the Data Matrix form should be used for this purpose).

SECTION I: METHOD OF DATA SUPPORT (Check one method only)

☐ I am using the cite-all method of support, and have included with this form a list of companies sent offers of compensation (the Data Matrix form should be used for this purpose).

☑ I am using the selective method of support (or cite-all option under the selective method), and have included with this form a completed list of data requirements (the Data Matrix form must be used).

SECTION II: GENERAL OFFER TO PAY

[Required if using the cite-all method or when using the cite-all option under the selective method to satisfy one or more data requirements]

☑ I hereby offer and agree to pay compensation, to other persons, with regard to the approval of this application, to the extent required by FIFRA.

SECTION III: CERTIFICATION

I certify that this application for registration, this form for reregistration, or this Data-Call-In response is supported by all data submitted or cited in the application for registration, the form for reregistration, or the Data-Call-In response. In addition, if the cite-all option or cite-all option under the selective method is indicated in Section I, this application is supported by all data in the Agency's files that (1) concern the properties or effects of this product or an identical or substantially similar product, or one or more of the ingredients in this product; and (2) is a type of data that would be required to be submitted under the data requirements in effect on the date of approval of this application if the application sought the initial registration of a product of identical or similar composition and uses .

I certify that for each exclusive use study cited in support of this registration or reregistration, that I am the original data submitter or that I have obtained the written permission of the original data submitter to cite that study.

I certify that for each study cited in support of this registration or reregistration that is not an exclusive use study, either: (a) I am the original data submitter; (b) I have obtained the permission of the original data submitter to use the study in support of this application; (c) all periods of eligibility for compensation have expired for the study; (d) the study is in the public literature; or (e) I have notified in writing the company that submitted the study and have offered (I) to pay compensation to the extent required by sections 3(c)(1)(F) and/or 3(c)(2)(B) of FIFRA; and (ii) to commence negotiations to determine the amount and terms of compensation, if any, to be paid for the use of the study.

I certify that in all instances where an offer of compensation is required, copies of all offers to pay compensation and evidence of their delivery in accordance with sections 3(c)(1)(F) and/or 3(c)(2)(B) of FIFRA are available and will be submitted to the Agency upon request. Should I fail to produce such evidence to the Agency upon request, I understand that the Agency may initiate action to deny, cancel or suspend the registration of my product in conformity with FIFRA.

I certify that the statements I have made on this form and all attachments to it are true, accurate, and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

Signature	Date	Typed or Printed Name and Title
N. Bhushan Mandava	July 9, 2012	N. Bhushan Mandava, Agent for Repar Corporation

EPA Form 8570-34 (9-97) Electronic and Paper versions available. Submit only Paper version.

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 14 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	
Ingredient Tebuconazole (CAS# 107534-96-3)		

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Generic Endangered Species Task Force	Pay	
			4110 136th Street, Northwest		
			Gig Harbor, WA 98332 (Company #84653)		
			Phibro Wood, LLC	Pay	
			65 Challenger Road, Third Floor		
			Ridgefield Park, NJ 07660 (Company # 84661)		
			Trustchem Co., Ltd.	Pay	
			89 Hanzhang Road, D2, 23rd Floor		
			Nanjing 210019 (Company # 86296)		
			Berkem Developpement	Pay	
			149 East 36th Street		
			New York, NY 10016 (Company # 86505)		
			Willowood Propiconazole, LLC	Pay	
			4110 136th Street, Northwest		
			Gig Harbor, WA 98332 (Company #87284)		

Signature : N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 13 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	

Ingredient Tebuconazole (CAS# 107534-96-3)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Sharda Worldwide Exports Pvt., Ltd.	Pay	
			P.O. Box 640		
			Hockessin, DE 19707 (Company #82633)		
			Rotam Agrochemical Company Limited	Pay	
			4032 Crockers Lake Blvd., Suite 818		
			Sarasota, FL 43238 (Company # 83100)		
			Direct AG Source, LLC	Pay	
			115 Obtuse Hill Road		
			Brookfield, CT 06804 (Company # 83222)		
			Sharda USA LLC	Pay	
			P.O. Box 640		
			Hockessin, DE 19707 (Company # 83529)		
			Amtide, LLC	Pay	
			21 Hubble		
			Irvine, CA 92618 (Company # 83851)		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 12 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	

Ingredient Tebuconazole (CAS# 107534-96-3)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Arch Treatment Technologies, Inc.	Pay	
			5660 New Northside Drive, Suite 1100		
			Atlanta, GA 30328 (Company #75506)		
			U.S. Triazole Task Force	Pay	
			600 13th Street, NW, 12th Floor		
			Washington, DC 20005 (Company # 75576)		
			Gro-Pro, LLC	Pay	
			4110 136th Street, Northwest		
			Gig Harbor, WA 98332 (Company # 79676)		
			Rotam Limited	Pay	
			4032 Crockers Lake Blvd., Suite 818		
			Sarasota, FL 43238 (Company # 81598)		
			Source Dynamics, LLC	Pay	
			10039 East Troon North Drive		
			Scottsdale, AZ 85262 (Company # 82542)		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 11 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	
Ingredient Tebuconazole (CAS# 107534-96-3)		

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Punjab Chemicals & Crop Protection, Ltd.	Pay	
			10039 East Troon North Drive		
			Scottsdale, AZ 85262 (Company #74280)		
			Osmose New Zealand	Pay	
			980 Ellicott Street		
			Buffalo, NY 14209 (Company # 74405)		
			Arch Wood Protection (NZ) Limited	Pay	
			5660 New Northside Drive, Ste 1100		
			Atlanta, GA 30328 (Company # 75101)		
			Agricultural Handlers Exposure Task Force	Pay	
			P.O. Box 509/1720 Prospect Drive		
			Macon, MO 63552 (Company # 75234)		
			Osmose Utilities Services, Inc.	Pay	
			980 Ellicott Street		
			Buffalo, NY 14209 (Company # 75341)		

Signature	Name and Title	Date
N. Bhushan Mandava	N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 10 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	
Ingredient Tebuconazole (CAS# 107534-96-3)		

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Agricultural Reentry Task Force	Pay	
			1350 I Street, N.W.		
			Washington, DC 20005 (Company #71755)		
			Bayer Advanced P.O. Box 12014	Pay	
			2 T.W. Alexander Drive		
			RTP, NC 27709 (Company # 72155)		
			Nisso TM LLC	Pay	
			45 Broadway, Suite 2120		
			New York, NY 10006 (Company # 73545)		
			FIFRA Endangered Species Task Force, LLC	Pay	
			1350 I Street, N.W.		
			Washington, DC 20005 (Company # 73989)		
			Irvita Plant Protection N.V.	Pay	
			4515 Falls of Neuse Rd., Suite 300		
			Raleigh, NC 27609 (Company # 74054)		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 9 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	

Ingredient Tebuconazole (CAS# 107534-96-3)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Spray Drift Task Force	Pay	
			1900 K Street, NW		
			Washington, DC 20006 (Company #66607)		
			Cheminova, Inc.	Pay	
			1600 Wilson Blvd., Suite 700		
			Arlington, VA 22209 (Company # 67760)		
			United Phosphorus, Inc.	Pay	
			630 Freedom Business Center, Suite 402		
			King of Prussia, PA 19406 (Company # 70506)		
			Sulphur Mills, Ltd.	Pay	
			1629 Pollasky Avenue, Suite 111		
			Clovis, CA 93612 (Company # 70905)		
			Outdoor Residential Exposure Task Force	Pay	
			1350 I Street, N.W.		
			Washington, DC 20005 (Company # 71754)		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 15 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	

Ingredient Tebuconazole (CAS# 107534-96-3)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Willowood, LLC	Pay	
			4110 136th Street, N.W		
			Gig Harbor, WA 98332 (Company #87290)		
			Willowood Tebuconazole, LLC	Pay	
			4110 136th Street, N.W		
			Gig Harbor, WA 98332 (Company #87811)		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 1 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	
Ingredient Tebuconazole (CAS# 107534-96-3)		

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
[redacted]			Repar Corporation	OWN	
			P.O. Box 4321 Silver Spring, MD 20914		
			Repar Corporation	OWN	
			Repar Corporation		
			Repar Corporation	OWN	
			Repar Corporation	OWN	
			Repar Corporation	OWN	
			Repar Corporation	OWN	
			Repar Corporation	OWN	
			Repar Corporation	OWN	
			Repar Corporation	OWN	
			Repar Corporation	OWN	
			Repar Corporation		
			Repar Corporation		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Public File Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 8 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	
Ingredient Tebuconazole (CAS# 107534-96-3)		

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			KOP-COAT, Inc.	Pay	
			436 Seventh Avenue		
			Pittsburgh, PA 15219 (Company #60061)		
			Sipcam Agro USA, Inc.	Pay	
			2520 Meridian Pkwy., Suite 525		
			Durham, NC 27713 (Company # 60063)		
			Arch Wood Protection, Inc.	Pay	
			5660 New Northside Drive, Suite 1100		
			Atlanta, GA 30328 (Company # 62190)		
			Makhteshim-Agan of North America, Inc.	Pay	
			4515 Falls of Neuse Rd., Suite 300		
			Raleigh, NC 27609 (Company # 66222)		
			Arysta Lifescience North America, LLC	Pay	
			15401 Weston Parkway, Suite 150		
			Cary, NC 27513 (Company # 66330)		

Signature	Name and Title	Date
N. Bhushan Mandava	N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 7 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	

Ingredient Tebuconazole (CAS# 107534-96-3)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Lanxess Corporation	Pay	
			111 RIDC Park West Drive		
			Pittsburgh, PA 15275 (Company #39967)		
			Luxembourg-Pamol, Inc.	Pay	
			3647 Willowbend Blvd., Suite 810		
			Houston, TX 77054 (Company # 42519)		
			Albaugh, Inc.	Pay	
			1525 NE 36th Street		
			Ankeny, IA 50021 (Company # 42750)		
			BASF Sparks, LLC	Pay	
			P.O. Box 13528		
			RTP, NC 27709 (Company # 51036)		
			NuFarm Americas, Inc.	Pay	
			150 Harvester Drive, Suite 200		
			Burr Ridge, IL 60527 (Company # 55146)		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 6 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	
Ingredient Tebuconazole (CAS# 107534-96-3)		

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Chemical Specialities, Inc.	Pay	
			P.O. Box 1330		
			Harrisburg, NC 28075 (Company #10465)		
			CP Chemicals, Inc.	Pay	
			2395 Cains Mill Road		
			Sumter, SC 29154 (Company # 11435)		
			Makhteshim Chemical Works, LTD.	Pay	
			4515 Falls of Neuse Road, Suite 300		
			Raleigh, NC 27609 (Company # 11678)		
			Loveland Products, Inc.	Pay	
			P.O. Box 1286		
			Greeley, CO 80632 (Company # 34704)		
			Oxon Italia S.P.A.	Pay	
			2520 Meridian Pkwy., Suite 525		
			Durham, NC 27713 (Company # 35915)		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012			EPA Reg No./File Symbol 69361-11		Page 5 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914			Product TEBUCON 3.6F FUNGICIDE		
Ingredient Tebuconazole (CAS# 107534-96-3)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Bayer Corp. P.O. Box 4913	Pay	
			8400 Hawthorn Road		
			Kansas City, MO 64120 (Company #3125)		
			Cerexagri, Inc.	Pay	
			630 Freedom Business Center, Suite 402		
			King of Prussia, PA 19406 (Company # 4581)		
			Cheminova A/S	Pay	
			1600 Wilson Blvd., Suite 700		
			Arlington, VA 22209 (Company # 4787)		
			Gustafson, LLC	Pay	
			P.O. Box 660065		
			Dallas, TX 75266 (Company # 7501)		
			J.J. Mauget Co.	Pay	
			12733 Director's Loop		
			Woodbridge, VA 22192 (Company # 7946)		
Signature N. Bhushan Mandava			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation		Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 4 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	

Ingredient Tebuconazole (CAS# 107534-96-3)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL for Generic Data			Bayer CropScience, LP P.O. Box 12014	Pay	
			2 T.W. Alexander Drive		
			Research Triangle Park, NC 27709 (CO.#264)		
			FMC CORP. Agricultural Products Group	Pay	
			1735 Market St., RM 1978		
			Philadelphia, PA 19103 (Company # 279)		
			Bayer Envrionmental Science P.O. Box 12014	Pay	
			2 T.W. Alexander Drive		
			Research Triangle Park, NC 27709 (CO.#432)		
			Cleary Chemicals, LLC	Pay	
			178 Ridge Road, Suite A		
			Dayton, NJ 08810 (Company # 1001)		
			Osmose, Inc.	Pay	
			980 Ellicott Street		
			Buffalo, NY 14209 (Company # 3008)		

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 3 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	
Ingredient Tebuconazole (CAS# 107534-96-3)		

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
830.7560	Partition Coefficient (n-octanol/water),		Repar Corporation		Not Required
	Generator Column Method				
830.7570	Partition Coefficient (n-octanol/water),		Repar Corporation		See 830.7550
	Estimation by Liquid Chromatography				
830.7840	Water Solubility: Column Elution Method;	47016901	Repar Corporation	OWN	
	Shake Flask Method				
830.7860	Water Solubility, Generator Column Method		Repar Corporation		See 830.7840
830.7950	Vapor Pressure		Repar Corporation		Not Required

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 2 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	

Ingredient Tebuconazole (CAS# 107534-96-3)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
830.6315	Flammability		Repar Corporation		Not Required
830.6316	Explodability		Repar Corporation		Not Required
830.6317	Storage Stability	47016901	Repar Corporation	OWN	
830.6319	Miscibility	47016901	Repar Corporation	OWN	
830.6320	Corrosion Characteristics	47016901	Repar Corporation	OWN	
830.6321	Dielectric Breakdown Voltage		Repar Corporation		Not Required
830.7000	pH	47016901	Repar Corporation	OWN	
830.7050	UV/Visible Absorption		Repar Corporation		Not Required
830.7100	Viscosity	47016901	Repar Corporation	OWN	
830.7200	Melting Point/Melting Range	47016901	Repar Corporation	OWN	
830.7220	Boiling Point/Boiling Range		Repar Corporation		Not Required
830.7300	Density/Relative Density/Bulk Density	47016901	Repar Corporation	OWN	
830.7370	Dissociation Constants in Water		Repar Corporation		Not Required
830.7550	Partition Coefficient (n-octanol/water),		Repar Corporation		Not Required
	Shake Flask Method				

Signature	Name and Title	Date
N. Bhushan Mandava	N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

Form Approved OMB No. 2070-0060

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date July 9, 2012	EPA Reg No./File Symbol 69361-11	Page 1 of 15
Applicant's/Registrant's Name & Address Repar Corporation P.O. Box 4321 Silver Spring, MD 20914	Product TEBUCON 3.6F FUNGICIDE	

Ingredient Tebuconazole (CAS# 107534-96-3)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
830.1550	Product Identity and Composition	47016901	Repar Corporation	OWN	
			P.O. Box 4321 Silver Spring, MD 20914		
830.1600	Description of Materials Used to Produce the Product	47016901	Repar Corporation	OWN	
830.1620	Description of Production Process		Repar Corporation		Not Required
830.1650	Description of Formulation Process	47016901	Repar Corporation	OWN	
830.1670	Discussion of Formation of Impurities	47016901	Repar Corporation	OWN	
830.1700	Preliminary Analysis	47016901	Repar Corporation	OWN	
830.1750	Certified Limits	47016901	Repar Corporation	OWN	
830.1800	Enforcement Analytical Method	47016901	Repar Corporation	OWN	
830.6302	Color	47016901	Repar Corporation	OWN	
830.6303	Physical State	47016901	Repar Corporation	OWN	
830.6304	Odor	47016901	Repar Corporation	OWN	
830.6313	Stability to Normal & Elevated Temperatures, Metals		Repar Corporation		Not Required
	and Metal Ions				
830.6314	Oxidation/Reduction: Chemical Incompatibility		Repar Corporation		Not Required

Signature N. Bhushan Mandava	Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar Corporation	Date July 9, 2012

EPA Form 8570-35 (9-97) Electronic and Paper versions available. Submit only Paper version.

Agency Internal Use Copy

MANDAVA ASSOCIATES, LLC

CONSULTANTS IN SCIENCE, TECHNOLOGY AND REGULATORY AFFAIRS
1050 Connecticut Avenue, N.W., Suite 1000, WASHINGTON, DC 20036-4510
Telephone: (202)-223-1424/1747 · Fax: (202)-223-0141 · E-MAIL: mandava@compuserve.com

HAND DELIVERED

July 11, 2012

Ms. Mary L. Waller
Product Manager (PM 21)
Fungicide Branch
Registration Division (7504P)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Washington, DC 20460

SUBJECT:	Submission:	Label Amendment
	Product Name:	Tebucon® 3.6F Fungicide
	Active Ingredient:	Tebuconazole
	Company:	Repar Corporation
	EPA Reg. No.:	69361-11

Dear Ms. Waller:

This is to notify the EPA that we are submitting the Label Amendment for Tebucon® 3.6F Fungicide (EPA Reg. No.: 69361-11). At this time we are incorporating the approved labels for Tebucon® 3.6F Fungicide (EPA Reg. No. 69361-11) label for Agricultural Crop uses and the Tebuconazole 3.6F Fungicide (EPA Reg. No. 69361-27) label for Turf and Ornamental uses into a single master label with 2 sub labels.

I. This label amendment contains a Master Label for Tebucon® 3.6F Fungicide with:
   1) Sub Label A: For Use in Agricultural Crops
   2) Sub Label B: For Use in Ornamentals and Golf Courses

II. At this time, we are also notifying the Agency for an **alternate brand name** of Tebucon® 3.6F Foliar Fungicide for this product as part of this submission.

Please find the following included with this submission:

1. Application for Pesticide (EPA Form 8570-1)
2. Certification with Respect to Citation of Data (EPA Form 8570-34)
3. Data Matrix (EPA Form 8570-35)
4. 5 Label copies for Tebucon® 3.6F Fungicide (EPA Reg. 69361-11)

We request the Agency to please approve the label amendment for Tebucon® 3.6F Fungicide at the earliest and your assistance in this matter would be greatly appreciated.

Thank you for your comments and assistance on this subject matter.

Sincerely,

N. Bhushan Mandava

N. Bhushan Mandava, Ph.D.
Agent for Repar Corporation

Enclosures

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95

EPA

United States
Environmental Protection Agency
Washington, DC 20460

- [ ] Registration
- [x] Amendment
- [ ] Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number	2. EPA Product Manager	3. Proposed Classification
69361-11	Mary Waller	[x] None [ ] Restricted
4. Company/Product (Name) Repar Corporation/TEBUCON 3.6F FUNGICIDE	**PM#** 21	

5. Name and Address of Applicant *(Include ZIP Code)*
Repar Corporation
P.O. Box 4321
Silver Spring, MD. 20914

[ ] *Check if this is a new address*

6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to:
EPA Reg. No. _____
Product Name _____

Section - II

- [x] Amendment - Explain below.
- [ ] Resubmission in response to Agency letter dated _____
- [ ] Notification - Explain below.
- [ ] Final printed labels in repsonse to Agency letter dated _____
- [ ] "Me Too" Application.
- [ ] Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

1) This application is for the amended label for additional uses for TEBUCON 3.6F Fungicide (EPA Reg. No. 69361-11).
2) Submittal of the Certification with Respect to Citation of Data (EPA Form 8750 -34) and Data Matrix (EPA Form 8570 - 35).
3) FAST TRACK and EXPEDITED REVIEW
4) Please direct all correspondence to: Mandava Associates, LLC; 1050 Connecticut Avenue, NW / Suite 1000; Washington, D.C. 20036
5) The contact person fax number is (202) 223 - 0141 and e-mail address is: mandava@compuserve.com.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging	Unit Packaging	Water Soluble Packaging	2. Type of Container
[ ] Yes [x] No	[ ] Yes [x] No	[ ] Yes [x] No	[ ] Metal [x] Plastic [ ] Glass [ ] Paper [ ] Other (Specify) _____
Certification must be submitted	If "Yes" Unit Packaging wgt. / No. per container	If "Yes" Package wgt / No. per container	

3. Location of Net Contents Information	4. Size(s) Retail Container	5. Location of Label Directions
[x] Label [ ] Container		[x]

6. Manner in Which Label is Affixed to Product [x] Lithograph [ ] Paper glued [ ] Stenciled [ ] Other _____

Section - IV

1. Contact Point *(Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)*

Name	Title	Telephone No. (Include Area Code)
N. Bhushan Mandava	Agent for Repar Corporation	(202) 223-1424

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received **(Stamped)**

2. Signature	3. Title
N. Bhushan Mandava	Agent for Repar Corporation
4. Typed Name	**5. Date**
N. Bhushan Mandava	July 9, 2012

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete. White - EPA File Copy (original) Yellow - Applicant Copy

RE: tebuconazole label amendment comments
Srinivas Mandava, M.D.
to:
Heather Garvie, Bhushan Mandava
09/05/2012 05:53 PM
Hide Details
From: "Srinivas Mandava, M.D." <Srini@Mandava.com>

To: Heather Garvie/DC/USEPA/US@EPA, "Bhushan Mandava" <mandava@compuserve.com>

4 Attachments

Tebucon 3.6F Master Label+ A+B (09-05-12).pdf Tebucon 3.6F Master Label+ A+B (09-05-12).doc Tebuconazole 3.6F Master Label+ A+B (09-05-12).doc

Tebuconazole 3.6F Master Label+ A+B (09-05-12).pdf

Dear Heather:

Thank you for your email for the label amendments for Tebucon 3.6F Fungicide (EPA Reg. No. 69361-11) and Tebuconazole 3.6F Fungicide (EPA Reg. No. 69361-27). We have made the suggested changes to both the product labels and are sending them to you in both WORD and PDF file formats.

Again thank you for your assistance in this matter.

Regards,

Srinivas Mandava, M.D.
Mandava Associates, LLC
1050 Connecticut Avenue, N.W.
Suite 1000
Washington, D.C. 20036
Tel: (202) 223 - 1424
Fax: (202) 223 – 0141

****
Notice: This E-mail (including attachments) is covered by the Electronic Communications Privacy Act, 18 U.S.C. §§2510-2521, is confidential and may be legally privileged. If you are not the intended recipient, you are hereby notified that any retention, dissemination, distribution, or copying of this communication is strictly prohibited. Please reply to the sender that you have received the message in error and then delete it. Thank you.
****

From: Heather Garvie [mailto:Garvie.Heather@epamail.epa.gov]
Sent: Wednesday, September 05, 2012 1:41 PM
To: Bhushan Mandava
Cc: Srinivas Mandava, M.D.
Subject: tebuconazole label amendment comments

Dr. Mandava:

Mary Waller has had the opportunity to go over your three tebuconazole labels that you submitted as amendments and has a few comments that need to be addressed before these can be approved. Please submit revised labels to me via email at your earliest convenience. Comments are as follows:

69361-26: no comments

69361-11:

1. If you want to keep the current format of this label, then you'll need to add in all of the WPS required language. So on page 27 you'll need to add the WPS

tebuconazole label amendment comments

Heather Garvie to: Bhushan Mandava 09/05/2012 01:40 PM

Cc: "Srinivas Mandava, M.D."

From: Heather Garvie/DC/USEPA/US

To: "Bhushan Mandava" <mandava@compuserve.com>

Cc: "Srinivas Mandava, M.D." <Srini@Mandava.com>

Dr. Mandava:

Mary Waller has had the opportunity to go over your three tebuconazole labels that you submitted as amendments and has a few comments that need to be addressed before these can be approved. Please submit revised labels to me via email at your earliest convenience. Comments are as follows:

69361-26: no comments

69361-11:

1. If you want to keep the current format of this label, then you'll need to add in all of the WPS required language. So on page 27 you'll need to add the WPS box and the language immediately above the Non-Agricultural Use Requirements box and language. This is because ornamentals are subject to WPS.

2. page 1: change the word ornaments to ornamentals under Sub label B heading.

3. page 1: change the word welch onion to welsh onion under sub label A.

4. page 1: move the alternate brand name asterisk and information immediately under the product name and above the line that is above "Sub label A: Agricultural Uses". Do the same on page 25.

5. page 3: in the User Safety Recommendations box, add a third bullet that reads, "Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing." Be sure to add the same language to page 26 as well.

6. page 5: under the "Spray Volume" heading, change the first sentence to read, "Apply Tebucon 3.6F Fungicide in a minimum of....."

7. page 7: item number 8; spell out AMS and UAN. Do the same on page 29.

8. page 20 in the Application Directions box under "Notes": change the word "recommended" to "labeled" in the second sentence.

9. page 23 under Seed Labeling; revise the second sentence to read, "Treated Seed. Do not use for Food, Feed or Oil Purposes. Treated with the fungicide Tebuconazole. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice." This information all must be in quotations.

10. page 33; under Ornamental Use Restrictions and Precautions; update the third statement to read, "Do not apply more than 10 fl oz per acre in a single application. Also revise the fourth statement to read, " "Do not apply more than 0.31 gallons (40 fl oz) of Tebucon 3.6F Fungicide (equal to 1.13 lbs of tebuconazole) per acre per year".

69361-27: same comments as 69361-11.

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

June 19, 2012

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

N. BHUSHAN MADAVA
MANDAVA ASSOCIATES, LLC
REPAR CORP
1050 CONN. AVE., N.W., SUITE 1000
WASHINGTON, DC 20036

PRODUCT NAME: TEBUCON 3.6F FUNGICIDE
COMPANY NAME: REPAR CORP
OPP IDENTIFICATION NUMBER:
EPA FILE SYMBOL: 69361-11
EPA RECEIPT DATE: 06/15/12

SUBJECT: RECEIPT OF AMENDMENT

DEAR REGISTRANT:

The Office of Pesticide Programs has received your application for an amendment and it has passed an administrative screen for completeness.

During the initial screen we determined that the application appears to qualify for fast track review. The package will now be forwarded to the Product Manager for review to determine its acceptability for fast track status.

If you have any questions, please contact Registration Division, Risk Management Team 21, at (703) 308-9354.

Sincerely,

Front End Processing Staff
Information Services Branch
Information Technology & Resources Management Division

Fee for Service

{918644;~

This package includes the following

- ○ New Registration
- ⊙ Amendment

☐ Studies? ☐ Fee Waiver?

☐ volpay % Reduction: ____

for Division

- ○ AD
- ○ BPPD
- ⊙ RD

Risk Mgr. 21

Receipt No.	S- 918644
EPA File Symbol/Reg. No.	69361-11
Pin-Punch Date:	6/15/2012

☒ This item is NOT subject to FFS action.

Action Code:

Requested:

Granted:

Amount Due: $ _____

Parent/Child Decisions:

☐ Inert Cleared for Intended Use ☐ Uncleared Inert in Product

Reviewer: Jennifer Haines Date: 6/19/12

Remarks:

21

MANDAVA ASSOCIATES, LLC

CONSULTANTS IN SCIENCE, TECHNOLOGY AND REGULATORY AFFAIRS
1730 M STREET, N.W., SUITE 906, WASHINGTON, DC 20036-4510
Telephone: (202)-223-1424/1747 · Fax: (202)-223-0141 · E-MAIL: Mandava@compuserve.com

October 23, 2009

Mary L. Waller
Product Manager 21
Fungicide Branch
Registration Division
Office of Pesticide Programs
Washington, DC 20460

SUBJECT: Official Name Change for Tebucon 3.6 F Fungicide
EPA Reg. No.: 69361-11

Company: Repar Corporation

Dear Ms. Waller:

On September 28, 2008, EPA accepted a label for the subject product with the product name "Tebucon 3.6 F Foliar Fungicide". The listed official name of the product is "Tebucon 3.6F Fungicide".

By means of this letter, we request the Agency to approve the change of the official name for Tebucon 3.6 F Fungicide (EPA Reg. No.: 69361-11).

Listed Official Name for the Product: Tebucon 3.6 F Fungicide
New Official Name for the Product: Tebucon 3.6 F **Foliar** Fungicide

We request you to approve the name change for the subject product registration.

Sincerely,

N. Bhushan Mandava

N. Bhushan Mandava
Agent for Repar Corporation

RE: Product Name: Repar Corp. Tebucon 3.6F EPA Reg No 69361-11
Bhushan Mandava to: Robert Westin 10/23/2009 03:25 PM
Cc: Mary Waller

Dear Bob:
Attached, please find a copy of the letter requesting the Agency to change the product official name from "Tebucon 3.6 F Fungicide" to "Tebucon 3.6F Foliar Fungicide".
The original letter (hard copy) is sent to you by U.S. Mail.

Best Regards,

N. Bhushan Mandava
Mandava Associates, LLC
1050 Connecticut Avenue, N.W.
Suite 1000
Washington, D.C. 20036
(202) 223 - 1424
(202) 223 – 0141

****
Notice: This E-mail (including attachments) is covered by the Electronic Communications Privacy Act, 18 U.S.C. §§2510-2521, is confidential and may be legally privileged. If you are not the intended recipient, you are hereby notified that any retention, dissemination, distribution, or copying of this communication is strictly prohibited. Please reply to the sender that you have received the message in error and then delete it. Thank you.
****

From: Westin.Robert@epamail.epa.gov [mailto:Westin.Robert@epamail.epa.gov]
Sent: Wednesday, October 21, 2009 10:11 AM
To: Bhushan Mandava
Subject: Product Name: Repar Corp. Tebucon 3.6F EPA Reg No 69361-11

Hi Bushan:

On September 28 EPA accepted a label for the subject product with the product name "Tebucon 3.6 F Foliar Fungicide". The listed official name of the product is "Tebucon 3.6F Fungicide". I need a letter from you requesting that the official name be changed to "Tebucon 3.6 F Foliar Fungicide".

Please give me a call if you have any questions.

Robert Westin

PPIS Deficiency Notice

PM NUMBER: 21 REGISTRATION NUMBER: 69361 - 11

Transferred from: Senedu Alemu Date Processed: 10/15/09

This jacket has been found to be deficient according to established EPA data processing Guidelines for the following reasons:

1. Current accepted label dated ____/____/_____
   ___ is missing from the jacket. ____ is missing acceptance stamp.
   ___ Acceptance stamp is missing date and/or registration number.
   ___ Is missing the following information: _____

2. Acceptance letter dated 9/28/09 and associated label are inconsistent concerning: ___ Acceptance date. X Product name. ___ Registration number.

3. Comments letter associated with label dated ____/____/_____ is missing from the jacket.

4. Accepted (Basic / Alternate) CSF dated ____/____/_____
   ___ Is missing from the jacket. ____ Is misfiled or has wrong registration number.
   ___ Sole active ingredient source product _____ - _____ has been cancelled.

5. ___ Active and inert chemical percentages do not equal 100%.

6. ✓ Other information missing / inconsistent within the jacket. See below for details.

No supporting document for addition or change product name.
Please review and return to contractors for data processing

Thanks!

REGISTRATION JACKET INTRA-TRANSFER REQUEST

(PLEASE FILL OUT FORM COMPLETELY)

Transfer File To: Mary Waller (PM21) Phone_____

Reason For Transfer:

- ☐ RSB CHEMISTRY REVIEW
- ☐ CODING (SIG)
- ☐ ACTION COMPLETED
- ☒ PRECAUTIONARY LABEL REVIEW
- ☐ WITHDRAW PENDING APPLICATION
- ☐ OTHER _____

TRANSFERRED FROM: **Senedu Alemu (SIG)** Phone: 305-6178

DATE REQUEST: 10/15/09 PM TEAM: _____ (If Applicable)

REGISTRATION NUMBER (S) OF JACKET (S) TO BE TRANSFERRED
(PLEASE INDICATE VOLUME NUMBER IF APPLICABLE)

REGISTRATION #	COMMENTS	LOCATION	OTHER INFORMATION
009361 - 11			
-			
-			
-			
-			
-			
-			
-			
-			

SPECIAL COMMENTS:

FOR FILE ROOM STAFF USE NOTIFY REQUESTOR:

DATE TIME INITIALS

Product Name: Repar Corp. Tebucon 3.6F EPA Reg No 69361-11

Robert Westin to: N. Bushan Mandava 10/21/2009 10:10 AM

Bcc: Robert Westin

Hi Bushan:

On September 28 EPA accepted a label for the subject product with the product name "Tebucon 3.6 F Foliar Fungicide". The listed official name of the product is "Tebucon 3.6F Fungicide". I need a letter from you requesting that the official name be changed to "Tebucon 3.6 F Foliar Fungicide".

Please give me a call if you have any questions.

Robert Westin

phone: 703-305-5721
FAX: 703-305-6920
email: westin.robert@epa.gov